

Multiconclusion Rules, Hypersequent Syntax and Step Frames

N. Bezhanishvili¹ and **S. Ghilardi**²

¹University of Amsterdam

²Università degli Studi di Milano

TOLO IV, Tbilisi, June 24, 2014

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Step-by-step constructions of free modal algebras have longstanding tradition:

- they can traced back to Fine normal forms (1975);
- Abramsky (1988, published much later in 2005);
- Ghilardi (1995; for Heyting case 1992, for S4 case 2010);
- coalgebraic literature (N. Bezhanishvili, Kurz, Kupke, Gehrke, Pattinson, Schröder, Venema, ... 2000→);
- recent advances in a paper by Coumans-vanGool (JLC 2013) followed by (N. Bezhanishvili, Ghilardi, Jibladze, in press);
- diagonalizable algebras case in (vanGool, AiML 2014).

20 Years in Free Modal Algebras Constructions

Coumans-van Gool emphasis is on partial algebras; a light reformulation of their point of view uses *two-sorted structures*.

These are called step-algebras and step-frames in (N. Bezhanishvili-Ghilardi-Jibladze, in press).

We review step-algebras and step-frames and then discuss ongoing work *concerning proof theoretic applications*

We outline the kind of proof theoretic aspects we want to investigate.

20 Years in Free Modal Algebras Constructions

Coumans-van Gool emphasis is on partial algebras; a light reformulation of their point of view uses *two-sorted structures*.

These are called step-algebras and step-frames in (N. Bezhanishvili-Ghilardi-Jibladze, in press).

We review step-algebras and step-frames and then discuss ongoing work *concerning proof theoretic applications*

We outline the kind of proof theoretic aspects we want to investigate.

20 Years in Free Modal Algebras Constructions

Coumans-van Gool emphasis is on partial algebras; a light reformulation of their point of view uses *two-sorted structures*.

These are called step-algebras and step-frames in (N. Bezhanishvili-Ghilardi-Jibladze, in press).

We review step-algebras and step-frames and then discuss ongoing work *concerning proof theoretic applications*

We outline the kind of proof theoretic aspects we want to investigate.

20 Years in Free Modal Algebras Constructions

Coumans-van Gool emphasis is on partial algebras; a light reformulation of their point of view uses *two-sorted structures*.

These are called step-algebras and step-frames in (N. Bezhanishvili-Ghilardi-Jibladze, in press).

We review step-algebras and step-frames and then discuss ongoing work *concerning proof theoretic applications*

We outline the kind of proof theoretic aspects we want to investigate.

Logics and Inference Systems

A **logic** L is a set of formulae containing tautologies, Aristotle's law and closed under necessitation, modus ponens and uniform substitution.

An **inference system** Ax for L is a set of inference rules

$$\frac{\phi_1(\underline{x}), \dots, \phi_n(\underline{x})}{\psi(\underline{x})} \quad (1)$$

satisfying the following conditions

Logics and Inference Systems

A **logic** L is a set of formulae containing tautologies, Aristotle's law and closed under necessitation, modus ponens and uniform substitution.

An **inference system** Ax for L is a set of inference rules

$$\frac{\phi_1(\underline{x}), \dots, \phi_n(\underline{x})}{\psi(\underline{x})} \quad (1)$$

satisfying the following conditions

Logics and Inference Systems

- all rules in Ax are derivable from L ;
- from the rules in Ax all formulae in L are provable;
- all formulae occurring as premises or as conclusions of rules in Ax have modal degree at most 1.

Rules satisfying the last condition are called *reduced*; non-reduced rules can be equivalently (for our purposes) replaced by reduced ones.

Logics and Inference Systems

- all rules in Ax are derivable from L ;
- from the rules in Ax all formulae in L are provable;
- all formulae occurring as premises or as conclusions of rules in Ax have modal degree at most 1.

Rules satisfying the last condition are called *reduced*; non-reduced rules can be equivalently (for our purposes) replaced by reduced ones.

Logics and Inference Systems

- all rules in Ax are derivable from L ;
- from the rules in Ax all formulae in L are provable;
- all formulae occurring as premises or as conclusions of rules in Ax have modal degree at most 1.

Rules satisfying the last condition are called *reduced*; non-reduced rules can be equivalently (for our purposes) replaced by reduced ones.

Logics and Inference Systems

- all rules in Ax are derivable from L ;
- from the rules in Ax all formulae in L are provable;
- all formulae occurring as premises or as conclusions of rules in Ax have modal degree at most 1.

Rules satisfying the last condition are called *reduced*; non-reduced rules can be equivalently (for our purposes) replaced by reduced ones.

Logics and Inference Systems

For every L one can find an Ax which is suitable for L (there are also canonical methods for that), but ...

... not all equivalent inference systems are proof-theoretically equally good ...

... for some of them proof search may become intricicated ...

... we want at least the following: *to prove a formula ϕ , only formulae up to the modal degree of ϕ are needed.*

Since the above property leads to decidability, it won't be possible to get it in general, but we want to have some criteria to recognize good systems.

Logics and Inference Systems

For every L one can find an Ax which is suitable for L (there are also canonical methods for that), but ...

... not all equivalent inference systems are proof-theoretically equally good ...

... for some of them proof search may become intricicated ...

... we want at least the following: *to prove a formula ϕ , only formulae up to the modal degree of ϕ are needed.*

Since the above property leads to decidability, it won't be possible to get it in general, but we want to have some criteria to recognize good systems.

Logics and Inference Systems

For every L one can find an Ax which is suitable for L (there are also canonical methods for that), but ...

... not all equivalent inference systems are proof-theoretically equally good ...

... for some of them proof search may become intricately ...

... we want at least the following: *to prove a formula ϕ , only formulae up to the modal degree of ϕ are needed.*

Since the above property leads to decidability, it won't be possible to get it in general, but we want to have some criteria to recognize good systems.

Logics and Inference Systems

For every L one can find an Ax which is suitable for L (there are also canonical methods for that), but ...

... not all equivalent inference systems are proof-theoretically equally good ...

... for some of them proof search may become intricicated ...

... we want at least the following: *to prove a formula ϕ , only formulae up to the modal degree of ϕ are needed.*

Since the above property leads to decidability, it won't be possible to get it in general, but we want to have some criteria to recognize good systems.

Logics and Inference Systems

For every L one can find an Ax which is suitable for L (there are also canonical methods for that), but ...

... not all equivalent inference systems are proof-theoretically equally good ...

... for some of them proof search may become intricicated ...

... we want at least the following: *to prove a formula ϕ , only formulae up to the modal degree of ϕ are needed.*

Since the above property leads to decidability, it won't be possible to get it in general, but we want to have some criteria to recognize good systems.

Logics and Inference Systems

As an example, take GL-system. It can be axiomatized by the single axiom $\Box(\Box x \rightarrow x) \rightarrow \Box x$.

An inference system for L consists of transitivity and Gödel-Löb rules

$$\frac{\Box^+ y \rightarrow x}{\Box y \rightarrow \Box x} \quad \frac{\Box x \rightarrow x}{x} \quad (2)$$

An alternative one (the good one!) consists of the rule

$$\frac{\Box^+ x \wedge \Box y \rightarrow y}{\Box x \rightarrow \Box y} \quad (3)$$

Logics and Inference Systems

Given a logic L , for a finite set of formulae Γ and for a formula ϕ , we write

$$\Gamma \vdash_L \phi \quad (4)$$

for the *global consequence relation*: this means that ϕ has a proof from axioms and rules of L with premises Γ (uniform substitution applies only to formulae in L , not to formulae in Γ). For transitive systems, $\Gamma \vdash_L \phi$ is the same as $\Box^+(\bigwedge \Gamma) \rightarrow \phi \in L$.

For axioms systems Ax , we have a similar notion

$$\Gamma \vdash_{Ax} \phi \quad (5)$$

Notice that if Ax is an inference system for L , the relations (4) and (5) are equivalent.

Logics and Inference Systems

Definition

An inference system Ax has the *bounded proof property* (bpp) iff whenever $\Gamma \vdash_{Ax} \phi$ holds, there is a proof in Ax of ϕ with premises Γ in which formulae not exceeding the modal degree of formulae in Γ, ϕ occur.

Definition

A logic L has the *finite model property* (fmp) iff whenever $\Gamma \vdash_{Ax} \phi$ does not hold, then there is a Kripke model based on a finite frame for L where the Γ are everywhere true and ϕ is not.

Logics and Inference Systems

Definition

An inference system Ax has the *bounded proof property* (bpp) iff whenever $\Gamma \vdash_{Ax} \phi$ holds, there is a proof in Ax of ϕ with premises Γ in which formulae not exceeding the modal degree of formulae in Γ, ϕ occur.

Definition

A logic L has the *finite model property* (fmp) iff whenever $\Gamma \vdash_{Ax} \phi$ does not hold, then there is a Kripke model based on a finite frame for L where the Γ are everywhere true and ϕ is not.

Outline

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

Outline

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

One-Step Modal Algebras

Definition

- 1 A *one-step modal algebra* is a quadruple $(A_0, A_1, i_0, \diamond_0)$, where A_0, A_1 are Boolean algebras, $i_0 : A_0 \rightarrow A_1$ is a Boolean morphism, and $\diamond_0 : A_0 \rightarrow A_1$ is a semilattice morphism. The algebras A_0, A_1 are called the *source* and the *target* Boolean algebras of the one-step modal algebra $(A_0, A_1, i_0, \diamond_0)$.
- 2 A *one-step extension* of the one-step modal algebra $(A_0, A_1, i_0, \diamond_0)$ is a one-step modal algebra $(A_1, A_2, i_1, \diamond_1)$ satisfying $i_1 \diamond_0 = \diamond_1 i_0$.

One-Step Modal Algebras

Definition

- 1 A *one-step modal algebra* is a quadruple $(A_0, A_1, i_0, \diamond_0)$, where A_0, A_1 are Boolean algebras, $i_0 : A_0 \rightarrow A_1$ is a Boolean morphism, and $\diamond_0 : A_0 \rightarrow A_1$ is a semilattice morphism. The algebras A_0, A_1 are called the *source* and the *target* Boolean algebras of the one-step modal algebra $(A_0, A_1, i_0, \diamond_0)$.
- 2 A *one-step extension* of the one-step modal algebra $(A_0, A_1, i_0, \diamond_0)$ is a one-step modal algebra $(A_1, A_2, i_1, \diamond_1)$ satisfying $i_1 \diamond_0 = \diamond_1 i_0$.

Applying Duality

To get a better understanding of the situation, it is nice to apply duality. Since we are interested in finite algebras, duality needs not topological machinery and is easy.

Definition

- 1 A *one-step frame* is a quadruple (W_1, W_0, f, R) , where W_0, W_1 are sets, $f : W_1 \rightarrow W_0$ is a function and $R \subseteq W_1 \times W_0$ is a relation.
- 2 A *one-step extension* of the one-step frame (W_1, W_0, f, R) is a one-step frame (W_2, W_1, g, S) satisfying $f \circ S = g \circ R$.

Applying Duality

To get a better understanding of the situation, it is nice to apply duality. Since we are interested in finite algebras, duality needs not topological machinery and is easy.

Definition

- 1 A *one-step frame* is a quadruple (W_1, W_0, f, R) , where W_0, W_1 are sets, $f : W_1 \rightarrow W_0$ is a function and $R \subseteq W_1 \times W_0$ is a relation.
- 2 A *one-step extension* of the one-step frame (W_1, W_0, f, R) is a one-step frame (W_2, W_1, g, S) satisfying $f \circ S = g \circ R$.

Step Correspondence Theory

We need to define what it means for for (the dual of) a one-step frame (W_1, W_0, f, R) to validate an inference rule. The definition is almost straightforward; we illustrate it via the example of the **K4** rule

$$\frac{\Box^+ y \rightarrow x}{\Box y \rightarrow \Box x}$$

This happens iff

$$\forall x, y \subseteq W_0 (f^*(x) \cap \Box_R x \subseteq f^*(y) \Rightarrow \Box_R x \subseteq \Box_R y). \quad (6)$$

The idea is to apply the step version of correspondence theory in order to eliminate the second order quantifiers via Ackermann rule.

Step Correspondence Theory

We need to define what it means for for (the dual of) a one-step frame (W_1, W_0, f, R) to validate an inference rule. The definition is almost straightforward; we illustrate it via the example of the **K4** rule

$$\frac{\Box^+ y \rightarrow x}{\Box y \rightarrow \Box x}$$

This happens iff

$$\forall x, y \subseteq W_0 (f^*(x) \cap \Box_R x \subseteq f^*(y) \Rightarrow \Box_R x \subseteq \Box_R y). \quad (6)$$

The idea is to apply the step version of correspondence theory in order to eliminate the second order quantifiers via Ackermann rule.

Step Correspondence Theory

We need to define what it means for for (the dual of) a one-step frame (W_1, W_0, f, R) to validate an inference rule. The definition is almost straightforward; we illustrate it via the example of the **K4** rule

$$\frac{\Box^+ y \rightarrow x}{\Box y \rightarrow \Box x}$$

This happens iff

$$\forall x, y \subseteq W_0 (f^*(x) \cap \Box_R x \subseteq f^*(y) \Rightarrow \Box_R x \subseteq \Box_R y). \quad (6)$$

The idea is to apply the step version of correspondence theory in order to eliminate the second order quantifiers via Ackermann rule.

Step Correspondence Theory

Applying adjunction $\exists_f \dashv f^*$, Ackermann rule, set-theoretic definition of \subseteq , definition of \square_R and Ackermann rule again we get:

$$\forall x, y \subseteq W_0. \quad f^*(x) \cap \square_R x \subseteq f^*(y) \Rightarrow \square_R x \subseteq \square_R y$$

$$\forall x, y \subseteq W_0. \quad \exists_f(f^*(x) \cap \square_R x) \subseteq y \Rightarrow \square_R x \subseteq \square_R y$$

$$\forall x \subseteq W_0. \quad \square_R x \subseteq \square_R \exists_f(f^*(x) \cap \square_R x)$$

$$\forall x \subseteq W_0. \quad \square_R x \subseteq \square_R \exists_f(\square_R x)$$

$$\forall x \subseteq W_0 \forall w \in W_0. \quad w \in \square_R x \Rightarrow w \in \square_R \exists_f(\square_R x).$$

$$\forall x \subseteq W_0 \forall w \in W_0. \quad R(w) \subseteq x \Rightarrow w \in \square_R \exists_f(\square_R x)$$

$$\forall w \in W_0. \quad w \in \square_R \exists_f(\square_R R(w)).$$

Step Correspondence Theory

Last condition reads as:

$$\forall w \forall v (R(w, v) \rightarrow \exists w_1 (f(w_1) = v \ \& \ R(w_1) \subseteq R(w))) . \quad (7)$$

The possibility of getting a first order condition is subject to sufficient syntactic conditions very similar to those of Sahlqvist theorem.

Step Correspondence Theory

Last condition reads as:

$$\forall w \forall v (R(w, v) \rightarrow \exists w_1 (f(w_1) = v \ \& \ R(w_1) \subseteq R(w))) . \quad (7)$$

The possibility of getting a first order condition is subject to sufficient syntactic conditions very similar to those of Sahlqvist theorem.

1 Ordinary Rules

- Step algebras and step frames
- **The Step Embedding Theorem**
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

The Step Embedding Theorem

Definition

A one-step frame (W_1, W_0, f, R) is *conservative* iff (i) f is surjective and (ii) for every $w_1, w_2 \in W_1$ we have that

$$f(w_1) = f(w_2) \ \& \ R(w_1) = R(w_2) \ \Rightarrow \ w_1 = w_2. \quad (8)$$

Dually, a one-step modal algebra (A_0, A_1, i, \diamond) is conservative iff (i) i is injective and (ii) the set

$$\{i(a) \mid a \in A_0\} \cup \{\diamond a \mid a \in A_0\}$$

generate A_1 as a Boolean algebra.

The Step Embedding Theorem

Definition

A one-step frame (W_1, W_0, f, R) is *conservative* iff (i) f is surjective and (ii) for every $w_1, w_2 \in W_1$ we have that

$$f(w_1) = f(w_2) \ \& \ R(w_1) = R(w_2) \ \Rightarrow \ w_1 = w_2. \quad (8)$$

Dually, a one-step modal algebra (A_0, A_1, i, \diamond) is conservative iff (i) i is injective and (ii) the set

$$\{i(a) \mid a \in A_0\} \cup \{\diamond a \mid a \in A_0\}$$

generate A_1 as a Boolean algebra.

The Step Embedding Theorem

Theorem

Let Ax be an inference system for a logic L . Then every finite conservative step-frame validating Ax is a p -morphic image of a finite Kripke frame for L iff Ax has the bpp and L has the fmp.

Kripke frames are here seen as step-frames with identical step transition function f ; the notion of a p -morphism between step-frames is the obvious one.

The Step Embedding Theorem

A p -morphism between step frames $\mathcal{F}' = (W'_1, W'_0, f', R')$ and $\mathcal{F} = (W_1, W_0, f, R)$ is a pair of surjective maps $\mu : W'_1 \rightarrow W_1$, $\nu : W'_0 \rightarrow W_0$ such that

$$f \circ \mu = \nu \circ f' \quad \text{and} \quad R \circ \mu = \nu \circ R' . \quad (9)$$

$$\begin{array}{ccc} W'_1 & \xrightarrow{\mu} & W_1 \\ f' \downarrow & & \downarrow f \\ W'_0 & \xrightarrow{\nu} & W_0 \end{array}$$

$$\begin{array}{ccc} W'_1 & \xrightarrow{\mu} & W_1 \\ R' \downarrow & & \downarrow R \\ W'_0 & \xrightarrow{\nu} & W_0 \end{array}$$

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- **An Example**
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

An example

Using the above theorem, it is easy to prove fmp and bpp for simple logics like K , $K4$, T , $S4$, \dots . As an example, let us consider the density axiom $\Box\Box x \rightarrow \Box x$ (we add this axiom to K).

First, we turn the axiom into a rule; there is a default naive method for that giving

$$\frac{y \rightarrow \Box x}{\Box y \rightarrow \Box x}. \quad (10)$$

Second, applying step correspondence, we get the following first-order characterization for validation of (10) in step frames:

$$\forall w \forall v (wRv \Rightarrow \exists k (wRf(k) \ \& \ kRv)) \quad (11)$$

An example

Using the above theorem, it is easy to prove fmp and bpp for simple logics like K , $K4$, T , $S4$, \dots . As an example, let us consider the density axiom $\Box\Box x \rightarrow \Box x$ (we add this axiom to K).

First, we turn the axiom into a rule; there is a default naive method for that giving

$$\frac{y \rightarrow \Box x}{\Box y \rightarrow \Box x}. \quad (10)$$

Second, applying step correspondence, we get the following first-order characterization for validation of (10) in step frames:

$$\forall w \forall v (wRv \Rightarrow \exists k (wRf(k) \ \& \ kRv)) \quad (11)$$

An example

Using the above theorem, it is easy to prove fmp and bpp for simple logics like K , $K4$, T , $S4$, \dots . As an example, let us consider the density axiom $\Box\Box x \rightarrow \Box x$ (we add this axiom to K).

First, we turn the axiom into a rule; there is a default naive method for that giving

$$\frac{y \rightarrow \Box x}{\Box y \rightarrow \Box x}. \quad (10)$$

Second, applying step correspondence, we get the following first-order characterization for validation of (10) in step frames:

$$\forall w \forall v (wRv \Rightarrow \exists k (wRf(k) \ \& \ kRv)) \quad (11)$$

An example

Third, we fix a finite conservative step-frame $\mathcal{S} = (W_1, W_0, f, R)$ satisfying (11); we must find a finite frame $\mathfrak{F} = (V, S)$ which dense in the standard sense

$$\forall w \forall v (wSv \Rightarrow \exists k (kSv \ \& \ wSk)). \quad (12)$$

and a surjective map $\mu : V \longrightarrow W_1$ such that $R \circ \mu = f \circ \mu \circ S$.

The idea is to take $V := W_1$ and $\mu := id_{W_1}$, so that we need to check

$$\forall w \forall v (wRv \Leftrightarrow \exists w' (wSw' \ \& \ f(w') = v)). \quad (13)$$

An example

Third, we fix a finite conservative step-frame $\mathcal{S} = (W_1, W_0, f, R)$ satisfying (11); we must find a finite frame $\mathfrak{F} = (V, S)$ which dense in the standard sense

$$\forall w \forall v (wSv \Rightarrow \exists k (kSv \ \& \ wSk)). \quad (12)$$

and a surjective map $\mu : V \longrightarrow W_1$ such that $R \circ \mu = f \circ \mu \circ S$.

The idea is to take $V := W_1$ and $\mu := id_{W_1}$, so that we need to check

$$\forall w \forall v (wRv \Leftrightarrow \exists w' (wSw' \ \& \ f(w') = v)). \quad (13)$$

An example

Third, we fix a finite conservative step-frame $\mathcal{S} = (W_1, W_0, f, R)$ satisfying (11); we must find a finite frame $\mathfrak{F} = (V, S)$ which dense in the standard sense

$$\forall w \forall v (wSv \Rightarrow \exists k (kSv \ \& \ wSk)). \quad (12)$$

and a surjective map $\mu : V \longrightarrow W_1$ such that $R \circ \mu = f \circ \mu \circ S$.

The idea is to take $V := W_1$ and $\mu := id_{W_1}$, so that we need to check

$$\forall w \forall v (wRv \Leftrightarrow \exists w' (wSw' \ \& \ f(w') = v)). \quad (13)$$

An example

Some ingenuity is needed in the general case to find the appropriate S but there are templates. Our template is

$$\forall w \forall w' (wSw' \Leftrightarrow \exists v (wRv \ \& \ f(w') = v)). \quad (14)$$

Thus, taking into consideration that f is also surjective (because S is conservative)

$$\forall v \exists w f(w) = v, \quad (15)$$

we need the validity of the implication

$$(15) \ \& \ (11) \ \& \ (14) \Rightarrow (13) \ \& \ (12).$$

An example

Some ingenuity is needed in the general case to find the appropriate S but there are templates. Our template is

$$\forall w \forall w' (wSw' \Leftrightarrow \exists v (wRv \ \& \ f(w') = v)). \quad (14)$$

Thus, taking into consideration that f is also surjective (because S is conservative)

$$\forall v \exists w f(w) = v, \quad (15)$$

we need the validity of the implication

$$(15) \ \& \ (11) \ \& \ (14) \ \Rightarrow \ (13) \ \& \ (12).$$

An example

Some ingenuity is needed in the general case to find the appropriate S but there are templates. Our template is

$$\forall w \forall w' (wSw' \Leftrightarrow \exists v (wRv \ \& \ f(w') = v)). \quad (14)$$

Thus, taking into consideration that f is also surjective (because S is conservative)

$$\forall v \exists w f(w) = v, \quad (15)$$

we need the validity of the implication

$$(15) \ \& \ (11) \ \& \ (14) \ \Rightarrow \ (13) \ \& \ (12).$$

An example

The implication

$$(15) \ \& \ (11) \ \& \ (14) \ \Rightarrow \ (13) \ \& \ (12).$$

can be shown to be valid manually or by a prover (SPASS solves the problem in less than half a second; the superposition proof it finds takes 47 lines).

As a consequence of this we get altogether bpp, fmp and first order definability (decidability and canonicity also follows by general reasons).

An example

The implication

$$(15) \ \& \ (11) \ \& \ (14) \ \Rightarrow \ (13) \ \& \ (12).$$

can be shown to be valid manually or by a prover (SPASS solves the problem in less than half a second; the superposition proof it finds takes 47 lines).

As a consequence of this we get altogether bpp, fmp and first order definability (decidability and canonicity also follows by general reasons).

An example

Let's summarize the three steps:

- **First:** produce the inference rules (there are automatic methods, not always they give the good rules).
- **Second:** apply correspondence theory (this is automatic).
- **Third:** produce p-morphic extensions to standard frames (not automatic, but there are templates); provers can discharge the final proof obligation.

An example

Let's summarize the three steps:

- **First:** produce the inference rules (there are automatic methods, not always they give the good rules).
- **Second:** apply correspondence theory (this is automatic).
- **Third:** produce p-morphic extensions to standard frames (not automatic, but there are templates); provers can discharge the final proof obligation.

An example

Let's summarize the three steps:

- **First:** produce the inference rules (there are automatic methods, not always they give the good rules).
- **Second:** apply correspondence theory (this is automatic).
- **Third:** produce p-morphic extensions to standard frames (not automatic, but there are templates); provers can discharge the final proof obligation.

An example

Let's summarize the three steps:

- **First**: produce the inference rules (there are automatic methods, not always they give the good rules).
- **Second**: apply correspondence theory (this is automatic).
- **Third**: produce p-morphic extensions to standard frames (not automatic, but there are templates); provers can discharge the final proof obligation.

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- **Some Case Studies**

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

Some case studies

We wonder to which extent the above mechanization of the metatheory can be pushed.

We analyzed some more significant cases. The first case is **GL** system axiomatized by the single axiom $\Box(\Box x \rightarrow x) \rightarrow \Box x$.

First Step can be driven so that to obtain a rule which is equivalent (for our purposes) to the well-known rule

$$\frac{\Box^+ x \wedge \Box y \rightarrow y}{\Box x \rightarrow \Box y}$$

Some case studies

We wonder to which extent the above mechanization of the metatheory can be pushed.

We analyzed some more significant cases. The first case is **GL** system axiomatized by the single axiom $\Box(\Box x \rightarrow x) \rightarrow \Box x$.

First Step can be driven so that to obtain a rule which is equivalent (for our purposes) to the well-known rule

$$\frac{\Box^+ x \wedge \Box y \rightarrow y}{\Box x \rightarrow \Box y}$$

Some case studies

We wonder to which extent the above mechanization of the metatheory can be pushed.

We analyzed some more significant cases. The first case is **GL** system axiomatized by the single axiom $\Box(\Box x \rightarrow x) \rightarrow \Box x$.

First Step can be driven so that to obtain a rule which is equivalent (for our purposes) to the well-known rule

$$\frac{\Box^+ x \wedge \Box y \rightarrow y}{\Box x \rightarrow \Box y}$$

Some case studies

Second Step (via Ackermann rule applied to fixpoint logic) gives

$$\forall w R(w) \subseteq \mu(Y, w) \exists_f (f^*(R(w)) \cap \Box_R R(w) \cap \Box_R Y). \quad (16)$$

In *finite* one-step frames this simplifies to

$$\forall w (R(w) \subseteq \{f(w') \mid R(w') \subset R(w)\}). \quad (17)$$

Notice that \subset is strict inclusion, so the above condition is a 'step' irreflexivity.

Some case studies

Second Step (via Ackermann rule applied to fixpoint logic) gives

$$\forall w R(w) \subseteq \mu(Y, w) \exists_f (f^*(R(w)) \cap \Box_R R(w) \cap \Box_R Y). \quad (16)$$

In *finite* one-step frames this simplifies to

$$\forall w (R(w) \subseteq \{f(w') \mid R(w') \subset R(w)\}). \quad (17)$$

Notice that \subset is strict inclusion, so the above condition is a ‘step’ irreflexivity.

Some case studies

Third Step is not difficult, but is not fully automatic. We can use the template S for transitive systems, but then the resulting Kripke frame is not irreflexive, so one needs to take the disjoint union of the irreflexive subframes satisfying (13).

It should be noticed that, if we do the same analysis for the system axiomatized by transitivity and Löb rule, we get a weaker condition than (16). Using the fact that the condition is too weak, it is possible to prove formally that bpp fails.

Some case studies

Third Step is not difficult, but is not fully automatic. We can use the template S for transitive systems, but then the resulting Kripke frame is not irreflexive, so one needs to take the disjoint union of the irreflexive subframes satisfying (13).

It should be noticed that, if we do the same analysis for the system axiomatized by transitivity and Löb rule, we get a weaker condition than (16). Using the fact that the condition is too weak, it is possible to prove formally that bpp fails.

Some case studies

Our second case study is the system **S4.3** axiomatized via **S4** reflexivity and transitivity axioms plus $\Box(\Box x \rightarrow y) \vee \Box(\Box y \rightarrow x)$.

First step: the inference rule extracted automatically from the axiom is not good (bpp fails). Instead, we use Goré infinitely many rules:

$$\frac{\dots \Box y \rightarrow x_j \vee \bigvee_{j \neq i} \Box x_i \dots}{\Box y \rightarrow \bigvee_{i=1}^n \Box x_i} \quad (18)$$

The rules are indexed by n and the n -th rule has n premises, according to the values $j = 1, \dots, n$.

Some case studies

Our second case study is the system **S4.3** axiomatized via **S4** reflexivity and transitivity axioms plus $\Box(\Box x \rightarrow y) \vee \Box(\Box y \rightarrow x)$.

First step: the inference rule extracted automatically from the axiom is not good (bpp fails). Instead, we use Goré infinitely many rules:

$$\frac{\dots \Box y \rightarrow x_j \vee \bigvee_{j \neq i} \Box x_i \dots}{\Box y \rightarrow \bigvee_{i=1}^n \Box x_i} \quad (18)$$

The rules are indexed by n and the n -th rule has n premises, according to the values $j = 1, \dots, n$.

Some case studies

Second Step: correspondence theory applies to these rules.
Interpreting the results in finite frames one gets

$$\forall w \forall S \subseteq R(w) \exists v \in S \exists w' (f(w') = v \ \& \ S \subseteq R(w') \subseteq R(w)). \quad (19)$$

Third Step: the same method used in **GL** case shows that one-step frames satisfying (19) are p-morphic images of Kripke frames for **S4.3**. This establishes bpp and fmp.

Some case studies

Second Step: correspondence theory applies to these rules.
Interpreting the results in finite frames one gets

$$\forall w \forall S \subseteq R(w) \exists v \in S \exists w' (f(w') = v \ \& \ S \subseteq R(w') \subseteq R(w)). \quad (19)$$

Third Step: the same method used in **GL** case shows that one-step frames satisfying (19) are p-morphic images of Kripke frames for **S4.3**.
This establishes bpp and fmp.

Some case studies

As a further case study let us consider **S5**.

First Step The following rule has been proposed in the literature:

$$\frac{\Box\Gamma \Rightarrow y, \Box\Delta}{\Box\Gamma \Rightarrow \Box y, \Box\Delta}. \quad (20)$$

In the resulting system, cuts cannot be completely eliminated, but can be limited to subformulae of the sequent to be proved. This ‘analytic’ cut-elimination property is sufficient to imply the bpp, and thus we should be able to get the bpp directly by our methods.

Some case studies

As a further case study let us consider **S5**.

First Step The following rule has been proposed in the literature:

$$\frac{\Box\Gamma \Rightarrow y, \Box\Delta}{\Box\Gamma \Rightarrow \Box y, \Box\Delta} . \quad (20)$$

In the resulting system, cuts cannot be completely eliminated, but can be limited to subformulae of the sequent to be proved. This ‘analytic’ cut-elimination property is sufficient to imply the bpp, and thus we should be able to get the bpp directly by our methods.

Some case studies

Second Step Correspondence theory gives

$$\forall w \forall v (wRv \rightarrow \exists \tilde{w} (f(\tilde{w}) = v \ \& \ R(w) = R(\tilde{w}))). \quad (21)$$

Third Step Step frames satisfying the above property are easily seen to be p-morphic images of reflexive, transitive, symmetric Kripke frames; this establishes bpp and fmp.

Some case studies

Second Step Correspondence theory gives

$$\forall w \forall v (wRv \rightarrow \exists \tilde{w} (f(\tilde{w}) = v \ \& \ R(w) = R(\tilde{w}))). \quad (21)$$

Third Step Step frames satisfying the above property are easily seen to be p-morphic images of reflexive, transitive, symmetric Kripke frames; this establishes bpp and fmp.

Some case studies

As a final case study consider the system obtained by adding to **K** the axiom $\Box\Box x \leftrightarrow \Box x$. This is **density+transitivity**; we can join the rules we already used for density and transitivity. This is not a good idea: **bpp fails!**

Instead, we use the following couple of rules suggested to us by G. Mints:

$$\frac{\Box + \Gamma \rightarrow \alpha}{\Box \Gamma \rightarrow \Box \alpha} \quad \frac{\Gamma, \Box \Delta \Rightarrow \Box \alpha}{\Box \Gamma, \Box \Delta \Rightarrow \Box \alpha} \quad (22)$$

Some case studies

As a final case study consider the system obtained by adding to **K** the axiom $\Box\Box x \leftrightarrow \Box x$. This is **density+transitivity**; we can join the rules we already used for density and transitivity. This is not a good idea: **bpp fails!**

Instead, we use the following couple of rules suggested to us by G. Mints:

$$\frac{\Box^+ \Gamma \rightarrow \alpha}{\Box \Gamma \rightarrow \Box \alpha} \quad \frac{\Gamma, \Box \Delta \Rightarrow \Box \alpha}{\Box \Gamma, \Box \Delta \Rightarrow \Box \alpha} \quad (22)$$

Some case studies

Second Step Correspondence theory gives, besides step transitivity (7), the condition

$$\forall w \forall v (wRv \rightarrow \exists w' (w'Rv \ \& \ \{f(w')\} \cup R(w') \subseteq R(w))). \quad (23)$$

Third Step Step frames satisfying the above property are easily seen to be p-morphic images of transitive and dense Kripke frames; this establishes bpp and fmp.

Notice that the fact that (11)+ (7) do not imply (23) is a formal argument proving that bpp fails if we adopt the old rule (10) in a transitive context. Thus, at least in principle, *model finders* can be used as automatic supports for showing that bpp fails.

Some case studies

Second Step Correspondence theory gives, besides step transitivity (7), the condition

$$\forall w \forall v (wRv \rightarrow \exists w' (w'Rv \ \& \ \{f(w')\} \cup R(w') \subseteq R(w))). \quad (23)$$

Third Step Step frames satisfying the above property are easily seen to be p-morphic images of transitive and dense Kripke frames; this establishes bpp and fmp.

Notice that the fact that (11)+ (7) do not imply (23) is a formal argument proving that bpp fails if we adopt the old rule (10) in a transitive context. Thus, at least in principle, *model finders* can be used as automatic supports for showing that bpp fails.

Some case studies

Second Step Correspondence theory gives, besides step transitivity (7), the condition

$$\forall w \forall v (wRv \rightarrow \exists w' (w'Rv \ \& \ \{f(w')\} \cup R(w') \subseteq R(w))). \quad (23)$$

Third Step Step frames satisfying the above property are easily seen to be p-morphic images of transitive and dense Kripke frames; this establishes bpp and fmp.

Notice that the fact that (11)+ (7) do not imply (23) is a formal argument proving that bpp fails if we adopt the old rule (10) in a transitive context. Thus, at least in principle, *model finders* can be used as automatic supports for showing that bpp fails.

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

Why many conclusions?

A *multiple-conclusion rule* is a pair of finite sets of formulae $\langle \Gamma, S \rangle$.

If $\Gamma = \{\gamma_1, \dots, \gamma_n\}$, $S = \{\delta_1, \dots, \delta_m\}$, we write the rule $\langle \Gamma, S \rangle$ as Γ/S or as

$$\frac{\gamma_1, \dots, \gamma_n}{\delta_1 \mid \dots \mid \delta_m} (R)$$

The formulae $\Gamma = \{\gamma_1, \dots, \gamma_n\}$ are said to be the *premises* of the rule (R) and the formulae $S = \{\delta_1, \dots, \delta_m\}$ are said to be the *conclusions* of the rule (R) .

The rule (R) is *valid* in a modal algebra (A, \Box) iff for every valuation V

$$V(\gamma_1) = 1 \ \& \ \dots \ \& \ V(\gamma_n) = 1 \quad \Rightarrow \quad V(\delta_1) = 1 \ \text{or} \ \dots \ \text{or} \ V(\delta_m) = 1 .$$

Thus *rule validity defines a universal class* (not a variety!).

Why many conclusions?

A *multiple-conclusion rule* is a pair of finite sets of formulae $\langle \Gamma, S \rangle$.

If $\Gamma = \{\gamma_1, \dots, \gamma_n\}$, $S = \{\delta_1, \dots, \delta_m\}$, we write the rule $\langle \Gamma, S \rangle$ as Γ/S or as

$$\frac{\gamma_1, \dots, \gamma_n}{\delta_1 \mid \dots \mid \delta_m} (R)$$

The formulae $\Gamma = \{\gamma_1, \dots, \gamma_n\}$ are said to be the *premises* of the rule (R) and the formulae $S = \{\delta_1, \dots, \delta_m\}$ are said to be the *conclusions* of the rule (R) .

The rule (R) is *valid* in a modal algebra (A, \Box) iff for every valuation V

$$V(\gamma_1) = 1 \ \& \ \dots \ \& \ V(\gamma_n) = 1 \quad \Rightarrow \quad V(\delta_1) = 1 \ \text{or} \ \dots \ \text{or} \ V(\delta_m) = 1 .$$

Thus *rule validity defines a universal class* (not a variety!).

Why many conclusions?

Multiple-conclusion rules recently gained attention in the literature from many points of view.

From an algebraic and a semantic point of view, (Kracht 07, Jerabek 09, N. & G. Bezhanishvili & Iemhoff 2014), they constitute an essential tool for investigating classes of algebras beyond varieties and they supply nice canonical formulae axiomatizations.

From a completely different research perspective, *the proof-theoretic oriented community* (since Avron 96) realized that standard sequent formalisms are insufficient to handle complex logics and moved to more expressive hypersequent calculi.

Why many conclusions?

Multiple-conclusion rules recently gained attention in the literature from many points of view.

From an algebraic and a semantic point of view, (Kracht 07, Jerabek 09, N. & G. Bezhanishvili & Iemhoff 2014), they constitute an essential tool for investigating classes of algebras beyond varieties and they supply nice canonical formulae axiomatizations.

From a completely different research perspective, *the proof-theoretic oriented community* (since Avron 96) realized that standard sequent formalisms are insufficient to handle complex logics and moved to more expressive hypersequent calculi.

Why many conclusions?

Multiple-conclusion rules recently gained attention in the literature from many points of view.

From an algebraic and a semantic point of view, (Kracht 07, Jerabek 09, N. & G. Bezhanishvili & Iemhoff 2014), they constitute an essential tool for investigating classes of algebras beyond varieties and they supply nice canonical formulae axiomatizations.

From a completely different research perspective, *the proof-theoretic oriented community* (since Avron 96) realized that standard sequent formalisms are insufficient to handle complex logics and moved to more expressive hypersequent calculi.

Why many conclusions?

Compare e.g. the simplicity of the hypersequent rule

$$\frac{\tilde{\Gamma}, \Box\Gamma, \Box\Gamma' \Rightarrow \Delta \quad \tilde{\Gamma}', \Box\Gamma', \Box\Gamma \Rightarrow \Delta'}{\tilde{\Gamma}, \Box\Gamma \Rightarrow \Delta \mid \tilde{\Gamma}', \Box\Gamma' \Rightarrow \Delta'} \quad (\text{Dich})$$

for **S4.3** with the above Goré rules.

Rule (Dich) can be rewritten as multiconclusion rule to

$$\frac{\Box\gamma \wedge \Box\gamma' \rightarrow \delta \quad \Box\gamma' \wedge \Box\gamma \rightarrow \delta'}{\Box\gamma \rightarrow \delta \mid \Box\gamma' \rightarrow \delta'}$$

Notice however that (Dich) does not define the variety of **S4.3** algebras but a universal class of algebras

$$\forall x \forall y (\Box x \leq \Box y \text{ or } \Box y \leq \Box x)$$

generating it.

Why many conclusions?

A similar observation applies to Avron's **S5** hypersequent rule

$$\frac{\Box\Gamma, \Gamma' \Rightarrow \Box\Delta, \Delta'}{\Box\Gamma \Rightarrow \Box\Delta \mid \Gamma' \Rightarrow \Delta'} \text{ (S5)}$$

axiomatizing the universal class of algebras where we have

$$\forall x (\Box x = 1 \text{ or } \Box x = 0)$$

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- Stable Classes

Derived Rules

Let K be a set of multiple-conclusion rules; a multiple-conclusion rule Γ/S is *derivable from K* - written $K \vdash \Gamma/S$ iff every modal algebra validating all rules in K validates also Γ/S .

In the terminology of modal rule systems (Jerabek 09, N. & G. Bezhaniashvili & Iemhoff 2014), it can be proved that this equivalently means that Γ/S belongs to the smallest modal rule system including K .

What we want to build here is a *Hilbert style* calculus for recognizing $K \vdash \Gamma/S$. This calculus will consequently be complete also for global consequence relation in K .

Derived Rules

Let K be a set of multiple-conclusion rules; a multiple-conclusion rule Γ/S is *derivable from K* - written $K \vdash \Gamma/S$ iff every modal algebra validating all rules in K validates also Γ/S .

In the terminology of modal rule systems (Jerabek 09, N. & G. Bezhanišvili & Iemhoff 2014), it can be proved that this equivalently means that Γ/S belongs to the smallest modal rule system including K .

What we want to build here is a *Hilbert style* calculus for recognizing $K \vdash \Gamma/S$. This calculus will consequently be complete also for global consequence relation in K .

Derived Rules

Let K be a set of multiple-conclusion rules; a multiple-conclusion rule Γ/S is *derivable from K* - written $K \vdash \Gamma/S$ iff every modal algebra validating all rules in K validates also Γ/S .

In the terminology of modal rule systems (Jerabek 09, N. & G. Bezhnashvili & Iemhoff 2014), it can be proved that this equivalently means that Γ/S belongs to the smallest modal rule system including K .

What we want to build here is a *Hilbert style* calculus for recognizing $K \vdash \Gamma/S$. This calculus will consequently be complete also for global consequence relation in K .

Hyperformulae and derivations

Our calculus will manipulate hyperformulae, seen as disjunctions of global assertions (this is the shape of conclusions of our multi-conclusion rules).

A *hyperformula* is a finite set of propositional formulae written in the form

$$\alpha_1 \mid \cdots \mid \alpha_n. \quad (24)$$

We use letters S, S_1, S', \dots for hyperformulae; the notation $S \mid S'$ means set union and $S \mid \alpha$ and $\alpha \mid S$ stand for $S \mid \{\alpha\}$ and $\{\alpha\} \mid S$, respectively.

Hyperformulae and derivations

Our calculus will manipulate hyperformulae, seen as disjunctions of global assertions (this is the shape of conclusions of our multi-conclusion rules).

A *hyperformula* is a finite set of propositional formulae written in the form

$$\alpha_1 \mid \cdots \mid \alpha_n. \quad (24)$$

We use letters S, S_1, S', \dots for hyperformulae; the notation $S \mid S'$ means set union and $S \mid \alpha$ and $\alpha \mid S$ stand for $S \mid \{\alpha\}$ and $\{\alpha\} \mid S$, respectively.

Hyperformulae and derivations

Our calculus will manipulate hyperformulae, seen as disjunctions of global assertions (this is the shape of conclusions of our multi-conclusion rules).

A *hyperformula* is a finite set of propositional formulae written in the form

$$\alpha_1 \mid \cdots \mid \alpha_n. \quad (24)$$

We use letters S, S_1, S', \dots for hyperformulae; the notation $S \mid S'$ means set union and $S \mid \alpha$ and $\alpha \mid S$ stand for $S \mid \{\alpha\}$ and $\{\alpha\} \mid S$, respectively.

Hyperformulae and derivations

Definition

Let Γ be a set of propositional modal formulae and let K be a set of multiple-conclusion rules. A *K-hyperproof* (or a *K-derivation* or just a *derivation*) under assumptions Γ is a finite list of hyperformulae S_1, \dots, S_n such that each S_i in it matches one of the following requirements:

- (i) S_i is of the kind $\alpha \mid S$, where $\alpha \in \Gamma$ or α is a tautology or α is an instance of the **K** distribution axiom;
- (ii) S_i is obtained from hyperformulae preceding it by applying a rule from K or the necessitation rule or the modus ponens rule.

We write $\Gamma \vdash_K S$ to mean that there is a *K-derivation* ending with S .

Hyperformulae and derivations

An important remark is in order for (ii): when we say that S_i is obtained by applying an inference rule, *we include uniform substitution and weakening in the application of the rule*. Thus, if the rule is

$$\frac{\gamma_1, \dots, \gamma_n}{\delta_1 \mid \dots \mid \delta_m} (R)$$

when we say that S_i is obtained from (R) , we mean that there are a hyperformula S and a substitution σ such that S_i is of the kind $S \mid \delta_1\sigma \mid \dots \mid \delta_m\sigma$ and that there are $j_1, \dots, j_n < i$ such that S_{j_1} is of the kind $S \mid \gamma_1\sigma$, and \dots and S_{j_n} is of the kind $S \mid \gamma_n\sigma$.

Hyperformulae and derivations

In other words, when rule (R) is used, we apply a substitution to its contextual form

$$\frac{\gamma_1 \mid S, \dots, \gamma_n \mid S}{\delta_1 \mid \dots \mid \delta_m \mid S} (R)$$

Proposition

We have $K \vdash \Gamma / S$ iff there is a K -derivation under assumptions Γ ending in S .

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- **Step Frame Characterizations**
- Stable Classes

The Step embedding theorem (hyper version)

It is routine to define what it means for a modal calculus K (seen as a set of reduced multiconclusion rules) to enjoy fmp and bpp.

It is also routine to define validation of a reduced multiconclusion rule in a step algebra and in a step frame. We have

Theorem

Let K be a modal calculus. Then K enjoys both bpp and fmp iff every finite conservative step-frame validating K is a p -morphic image of a finite Kripke frame validating K .

As examples you can take the calculi obtained by translating hypersequent rules for **S4.3**, **S5** into multiconclusion rules (these systems axiomatize the class of corresponding *prime* algebras).

The Step embedding theorem (hyper version)

It is routine to define what it means for a modal calculus K (seen as a set of reduced multiconclusion rules) to enjoy fmp and bpp.

It is also routine to define validation of a reduced multiconclusion rule in a step algebra and in a step frame. We have

Theorem

Let K be a modal calculus. Then K enjoys both bpp and fmp iff every finite conservative step-frame validating K is a p -morphic image of a finite Kripke frame validating K .

As examples you can take the calculi obtained by translating hypersequent rules for **S4.3**, **S5** into multiconclusion rules (these systems axiomatize the class of corresponding *prime* algebras).

The Step embedding theorem (hyper version)

It is routine to define what it means for a modal calculus K (seen as a set of reduced multiconclusion rules) to enjoy fmp and bpp.

It is also routine to define validation of a reduced multiconclusion rule in a step algebra and in a step frame. We have

Theorem

Let K be a modal calculus. Then K enjoys both bpp and fmp iff every finite conservative step-frame validating K is a p -morphic image of a finite Kripke frame validating K .

As examples you can take the calculi obtained by translating hypersequent rules for **S4.3**, **S5** into multiconclusion rules (these systems axiomatize the class of corresponding *prime* algebras).

1 Ordinary Rules

- Step algebras and step frames
- The Step Embedding Theorem
- An Example
- Some Case Studies

2 Multiconclusion Rules

- A Hilbert calculus for hyperformulae
- Step Frame Characterizations
- **Stable Classes**

Homomorphic Images and Stability

A *stable embedding* of a modal algebra $\mathfrak{A} = (A, \diamond)$ into a modal algebra $\mathfrak{B} = (B, \diamond)$ is an injective Boolean morphism $\mu : A \rightarrow B$ such that we have $\diamond\mu(x) \leq \mu(\diamond x)$ for all $x \in A$.

A class \mathcal{C} of modal algebras is said to be *stable* iff whenever $\mathfrak{B} \in \mathcal{C}$ and \mathfrak{A} has a stable embedding into \mathfrak{B} , then $\mathfrak{A} \in \mathcal{C}$ too.

We have dual notions for frames. $\mathfrak{F} = (W, R)$ is a *homomorphic image* of $\mathfrak{F}' = (W', R')$ iff there is a surjective map $f : W' \rightarrow W$ such that $xR'y$ implies $f(x)Rf(y)$ for all $x, y \in W'$ (in case $\mathfrak{F}, \mathfrak{F}'$ are descriptive, f is asked to be continuous too).

A class of (ordinary or descriptive) frames is said to be *stable* iff it is closed under homomorphic images.

Homomorphic Images and Stability

A *stable embedding* of a modal algebra $\mathfrak{A} = (A, \diamond)$ into a modal algebra $\mathfrak{B} = (B, \diamond)$ is an injective Boolean morphism $\mu : A \rightarrow B$ such that we have $\diamond\mu(x) \leq \mu(\diamond x)$ for all $x \in A$.

A class \mathcal{C} of modal algebras is said to be *stable* iff whenever $\mathfrak{B} \in \mathcal{C}$ and \mathfrak{A} has a stable embedding into \mathfrak{B} , then $\mathfrak{A} \in \mathcal{C}$ too.

We have dual notions for frames. $\mathfrak{F} = (W, R)$ is a *homomorphic image* of $\mathfrak{F}' = (W', R')$ iff there is a surjective map $f : W' \rightarrow W$ such that $xR'y$ implies $f(x)Rf(y)$ for all $x, y \in W'$ (in case $\mathfrak{F}, \mathfrak{F}'$ are descriptive, f is asked to be continuous too).

A class of (ordinary or descriptive) frames is said to be *stable* iff it is closed under homomorphic images.

Homomorphic Images and Stability

A *stable embedding* of a modal algebra $\mathfrak{A} = (A, \diamond)$ into a modal algebra $\mathfrak{B} = (B, \diamond)$ is an injective Boolean morphism $\mu : A \rightarrow B$ such that we have $\diamond\mu(x) \leq \mu(\diamond x)$ for all $x \in A$.

A class \mathcal{C} of modal algebras is said to be *stable* iff whenever $\mathfrak{B} \in \mathcal{C}$ and \mathfrak{A} has a stable embedding into \mathfrak{B} , then $\mathfrak{A} \in \mathcal{C}$ too.

We have dual notions for frames. $\mathfrak{F} = (W, R)$ is a *homomorphic image* of $\mathfrak{F}' = (W', R')$ iff there is a surjective map $f : W' \rightarrow W$ such that $xR'y$ implies $f(x)Rf(y)$ for all $x, y \in W'$ (in case $\mathfrak{F}, \mathfrak{F}'$ are descriptive, f is asked to be continuous too).

A class of (ordinary or descriptive) frames is said to be *stable* iff it is closed under homomorphic images.

Homomorphic Images and Stability

A *stable embedding* of a modal algebra $\mathfrak{A} = (A, \diamond)$ into a modal algebra $\mathfrak{B} = (B, \diamond)$ is an injective Boolean morphism $\mu : A \rightarrow B$ such that we have $\diamond\mu(x) \leq \mu(\diamond x)$ for all $x \in A$.

A class \mathcal{C} of modal algebras is said to be *stable* iff whenever $\mathfrak{B} \in \mathcal{C}$ and \mathfrak{A} has a stable embedding into \mathfrak{B} , then $\mathfrak{A} \in \mathcal{C}$ too.

We have dual notions for frames. $\mathfrak{F} = (W, R)$ is a *homomorphic image* of $\mathfrak{F}' = (W', R')$ iff there is a surjective map $f : W' \rightarrow W$ such that $xR'y$ implies $f(x)Rf(y)$ for all $x, y \in W'$ (in case $\mathfrak{F}, \mathfrak{F}'$ are descriptive, f is asked to be continuous too).

A class of (ordinary or descriptive) frames is said to be *stable* iff it is closed under homomorphic images.

Homomorphic Images and Stability

A modal calculus K is *stable* iff so is the class of modal algebras validating it (equivalently: the class of descriptive frames validating it).

The following Theorem is proved in (N. & G. Bezhanishvili & Iemhoff 2014):

Theorem

- (i) *A modal calculus K is stable iff it is axiomatizable via stable characteristic rules.*
- (ii) *A stable modal calculus enjoys fmp.*

Homomorphic Images and Stability

A modal calculus K is *stable* iff so is the class of modal algebras validating it (equivalently: the class of descriptive frames validating it).

The following Theorem is proved in (N. & G. Bezhanishvili & Iemhoff 2014):

Theorem

- (i) *A modal calculus K is stable iff it is axiomatizable via stable characteristic rules.*
- (ii) *A stable modal calculus enjoys fmp.*

Stable Characteristic Rules

Stable characteristic rules are the rules associated with finite Kripke frames in the following way.

Let $\mathfrak{F} = (F, R_F)$ be a finite frame. For every $a \in F$ we introduce a new propositional variable x_a . The *modal stable rule* of \mathfrak{F} is

$$\frac{\bigvee_{i=1}^n x_{a_i}, \bigwedge_{i \neq j} \neg(x_{a_i} \wedge x_{a_j}), \bigwedge_{i=1}^n (x_{a_i} \rightarrow \Box \bigvee_{b \in R_F(a_i)} x_b)}{\neg x_{a_1} \mid \cdots \mid \neg x_{a_n}} (r_{\mathfrak{F}})$$

where we suppose that $F = \{a_1, \dots, a_n\}$.

Stable Characteristic Rules

The following proposition is proved in (N. & G. Bezhanishvili & Iemhoff 2014):

Proposition

Let $\mathfrak{A} = (A, \diamond)$ be a modal algebra. Then

- 1 \mathfrak{A} does not validate $(r_{\mathfrak{F}})$ iff there is a stable embedding of \mathfrak{F}^* into \mathfrak{A} .
- 2 \mathfrak{A} does not validate $(r_{\mathfrak{F}})$ iff there is a surjective stable map from \mathfrak{A}_* onto \mathfrak{F} .

Bpp for Stable Calculi

To get bpp however we need to modify rules (r_{\boxtimes}^-) as shown below.

$$\frac{\bigvee_{i=1}^n x_{a_i}, \bigwedge_{i \neq j} \neg(x_{a_i} \wedge x_{a_j}), \bigwedge_{i=1}^n (x_{a_i} \rightarrow \Box r_{a_i}), \bigwedge_{i=1}^n (r_{a_i} \rightarrow \bigvee_{b \in R_F(a_i)} x_b)}{\neg x_{a_1} \mid \cdots \mid \neg x_{a_n}}$$

Lemma

Rules (r_{\boxtimes}^+) and (r_{\boxtimes}^-) are inter-derivable.

Bpp for Stable Calculi

To get bpp however we need to modify rules ($r_{\mathfrak{F}}$) as shown below.

$$\frac{\bigvee_{i=1}^n x_{a_i}, \quad \bigwedge_{i \neq j} \neg(x_{a_i} \wedge x_{a_j}), \quad \bigwedge_{i=1}^n (x_{a_i} \rightarrow \Box r_{a_i}), \quad \bigwedge_{i=1}^n (r_{a_i} \rightarrow \bigvee_{b \in R_F(a_i)} x_b)}{\neg x_{a_1} \mid \cdots \mid \neg x_{a_n}}$$

Lemma

Rules ($r_{\mathfrak{F}}^+$) and ($r_{\mathfrak{F}}$) are inter-derivable.

Bpp for Stable Calculi

Theorem

Any modal calculus axiomatized by rules of the kind $(r_{\mathfrak{F}}^+)$ enjoys bpp and fmp.

Corollary

Let \mathcal{C} be a stable class of (ordinary) Kripke frames such that membership of a finite frame to \mathcal{C} is decidable. Then validity of a formula (more generally, of a rule) in \mathcal{C} is decidable as well.

Bpp for Stable Calculi

Theorem

Any modal calculus axiomatized by rules of the kind $(r_{\mathfrak{F}}^+)$ enjoys bpp and fmp.

Corollary

Let \mathcal{C} be a stable class of (ordinary) Kripke frames such that membership of a finite frame to \mathcal{C} is decidable. Then validity of a formula (more generally, of a rule) in \mathcal{C} is decidable as well.

Conclusions

- step methods seem to be quite effective in jointly proving bpp and fmp;
- in simple cases the application of the methods is fully automatic (in a sense we are *mechanizing the metatheory* of modal logic!);
- in more complex cases some ingenuity is needed, still uniform arguments often work;
- entire classes of logics can be covered (see the above results on stable classes);
- the scalability of the methods is to be tested for more complicated logics arising in computer science applications.

Conclusions

- step methods seem to be quite effective in jointly proving bpp and fmp;
- in simple cases the application of the methods is fully automatic (in a sense we are *mechanizing the metatheory* of modal logic!);
- in more complex cases some ingenuity is needed, still uniform arguments often work;
- entire classes of logics can be covered (see the above results on stable classes);
- the scalability of the methods is to be tested for more complicated logics arising in computer science applications.

Conclusions

- step methods seem to be quite effective in jointly proving bpp and fmp;
- in simple cases the application of the methods is fully automatic (in a sense we are *mechanizing the metatheory* of modal logic!);
- in more complex cases some ingenuity is needed, still uniform arguments often work;
- entire classes of logics can be covered (see the above results on stable classes);
- the scalability of the methods is to be tested for more complicated logics arising in computer science applications.

Conclusions

- step methods seem to be quite effective in jointly proving bpp and fmp;
- in simple cases the application of the methods is fully automatic (in a sense we are *mechanizing the metatheory* of modal logic!);
- in more complex cases some ingenuity is needed, still uniform arguments often work;
- entire classes of logics can be covered (see the above results on stable classes);
- the scalability of the methods is to be tested for more complicated logics arising in computer science applications.

Conclusions

- step methods seem to be quite effective in jointly proving bpp and fmp;
- in simple cases the application of the methods is fully automatic (in a sense we are *mechanizing the metatheory* of modal logic!);
- in more complex cases some ingenuity is needed, still uniform arguments often work;
- entire classes of logics can be covered (see the above results on stable classes);
- the scalability of the methods is to be tested for more complicated logics arising in computer science applications.