




# Quantitative results on continuity of the spectral factorization mapping in the scalar case

Lasha Ephremidze<sup>1,3</sup> · Eugene Shargorodsky<sup>2</sup> ·  
Ilya Spitkovsky<sup>1,4</sup>

Received: 1 January 2016 / Revised: 18 February 2016 / Accepted: 26 February 2016 /  
Published online: 8 April 2016  
© The Author(s) 2016. This article is published with open access at Springerlink.com

**Abstract** In the scalar case, the spectral factorization mapping  $f \rightarrow f^+$  puts a nonnegative integrable function  $f$  having an integrable logarithm in correspondence with an outer analytic function  $f^+$  such that  $f = |f^+|^2$  is almost everywhere. The main question addressed here is to what extent  $\|f^+ - g^+\|_{H_2}$  is controlled by  $\|f - g\|_{L_1}$  and  $\|\log f - \log g\|_{L_1}$ .

**Keywords** Spectral factorization · Paley–Wiener condition · Convergence rate



To Sergei Grudsky on the occasion of his 60th birthday.

Lasha Ephremidze was partially supported by the Shota Rustaveli National Science Foundation Grant (Contract Numbers: 31/47) and Ilya Spitkovsky was supported in part by Faculty Research funding from the Division of Science and Mathematics, New York University Abu Dhabi, and by Plumeri Award for Faculty Excellence from the College of William and Mary.

✉ Eugene Shargorodsky  
eugene.shargorodsky@kcl.ac.uk

- 1 Division of Science and Mathematics, New York University Abu Dhabi (NYUAD), Saadiyat Island, P.O. Box 129188, Abu Dhabi, United Arab Emirates
- 2 Department of Mathematics, Kings College London, Strand, London WC2R 2LS, UK
- 3 A. Razmadze Mathematical Institute, I. Javakishvili Tbilisi State University, 6, Tamarashvili st., Tbilisi 0177, Georgia
- 4 Department of Mathematics, College of William and Mary, Williamsburg, VA 23187, USA

## Mathematics Subject Classification 47A68

### 1 Introduction

Let  $f$  be a nonnegative integrable function on the unit circle in the complex plain,  $0 \leq f \in L_1(\mathbb{T})$ , satisfying the Paley–Wiener condition

$$\log f \in L_1(\mathbb{T}). \quad (1)$$

Then it admits a spectral factorization

$$f(t) = f^+(t)f^-(t) \quad \text{a.e. on } \mathbb{T}, \quad (2)$$

where  $f^+$  is a function analytic inside the unit circle,  $f^+ \in \mathcal{A}(\mathbb{T}_+)$ , and  $f^-(z) = \overline{f^+(1/\bar{z})}$ , which is analytic outside the unit circle including the infinity,  $f^- \in \mathcal{A}(\mathbb{T}_-)$ . More specifically,  $f^+$  belongs to the Hardy space  $H_2(\mathbb{D})$ ; therefore, its boundary values  $f^+(t) = f^+(e^{i\theta}) = \lim_{r \rightarrow 1} f^+(re^{i\theta})$  exist a.e. and the Eq. (2) holds for these boundary values. Note also that  $f^+ = \overline{f^-}$  a.e. on  $\mathbb{T}$  and therefore (2) is equivalent to

$$f(t) = |f^+(t)|^2 \quad \text{a.e. on } \mathbb{T}.$$

Condition (1) is necessary for factorization (2) to exist. It also plays an important role in the linear prediction theory of stationary stochastic processes, one of the historically first applications of spectral factorization (see [16,21]). Namely, let  $\dots, X_{-1}, X_0, X_1, \dots$  be a stationary stochastic process with the spectral measure  $d\mu = f dt + d\mu_s$ . In a different but equivalent language,  $\{X_n\}_{n \in \mathbb{Z}}$  is a sequence in a Hilbert space and  $\langle X_n, X_k \rangle = \frac{1}{2\pi} \int_{\mathbb{T}} e^{i(n-k)\theta} d\mu(\theta)$ . The process is *deterministic* if  $X_{n+1}$  can be represented as the limit of linear combinations of vectors from  $\{\dots, X_{n-1}, X_n\}$ , i.e.  $X_{n+1} \in \overline{\text{Span}\{\dots, X_{n-1}, X_n\}}$ . As it happens (see e.g., [16]), condition (1) is necessary and sufficient for the process to be non-deterministic.

Starting with the original applications in the prediction theory of stochastic processes, spectral factorization procedure appeared in such seemingly distant areas as singular integral equations [4,13], linear estimation [15], quadratic and  $H_\infty$  control [2,4,12], communications [11], filter design [8,19,20], etc.

If we require  $f^+$  to be an outer analytic function, then the factorization (2) is unique up to a constant factor  $c$  with absolute value 1,  $|c| = 1$ . The unique spectral factor which is positive at the origin can be a priori written as

$$f^+(z) = \exp\left(\frac{1}{4\pi} \int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} \log f(e^{i\theta}) d\theta\right). \quad (3)$$

In most applications, a spectral factor  $f^+$  in (2) is not explicitly required to be outer and instead is subject to certain extremal conditions called, in various works, minimal phase or maximal energy, optimal, etc. In mathematical terms, however, they amount to  $f^+$  being outer, so seeking the solution (3) is natural.

From the practical point of view, it is important to study the continuity properties of the spectral factorization map

$$f \mapsto f^+ \tag{4}$$

defined by (3). Namely, we are interested in knowing how close  $g^+$  is to  $f^+$  when a spectral density  $g$  is close to  $f$ . The reason why we study this question is that usually an estimated spectral density function  $\hat{f}$  being dealt with is constructed empirically from observations and is only an approximation to the theoretically existing spectral density  $f$ . Therefore, we need to know how close  $\hat{f}^+$  remains to  $f^+$  under such approximation.

An answer to the above question depends on norms we use as a measurement of the accuracy in the spaces of functions and of their spectral factors. Since the boundary values of the function (3) can be expressed as

$$f^+(t) = \sqrt{f(t)} \exp\left(\frac{i}{2} \widetilde{\log f(t)}\right),$$

where  $\sim$  stands for the harmonic conjugation operator

$$\tilde{h}(e^{i\tau}) = (P) \frac{1}{2\pi} \int_0^{2\pi} h(e^{i\theta}) \cot\left(\frac{\tau - \theta}{2}\right) d\theta,$$

and the conjugation is not a bounded operator on  $L_\infty$  or  $C(\mathbb{T})$ , it is not surprising that the map (4) is not continuous in these spaces [1]. Furthermore, it is shown in [5] that every continuous function on  $\mathbb{T}$  is a discontinuity point of the spectral factorization mapping in the uniform norm, whereas in [14] it was shown that on a large class of function spaces (the so-called decomposing Banach algebras) the spectral factorization mapping is continuous.

The spectral factorization of a trigonometric polynomial

$$f(t) = \sum_{k=-N}^N c_k t^k, \tag{5}$$

which is non-negative on  $\mathbb{T}$ , has the form

$$f(t) = \sum_{k=0}^N a_k t^k \sum_{k=0}^N \overline{a_k} t^{-k};$$

i.e., the spectral factor  $f^+$  is a polynomial of the same degree  $N$ . This result is known as the Fejér-Riesz lemma (see, e.g., [8]). The spectral factor can also be expressed in terms of zeros of polynomial (5), and therefore the map (4) is continuous on  $\mathcal{P}_N$ , the set of all functions of the form (5). Papers [6,7] are devoted to estimating the constant  $C_N$  in the inequality

$$\|\phi^+ - \psi^+\|_{L_\infty} \leq C_N \|\phi - \psi\|_{L_\infty}, \quad \phi, \psi \in \mathcal{P}_N,$$

and it is shown there that  $C_N \sim \log N$  asymptotically, under the condition that the values of functions  $\phi$  and  $\psi$  are bounded away from 0.

Moving to Lebesgue spaces, the map (4) is not continuous in the  $L_1$  norm in general, since a small change of values of function  $f$ , if these values are close to 0, may cause a significant change of  $\log f$ . However,

$$\|f_n - f\|_{L_1} \rightarrow 0 \text{ and } \|\log f_n - \log f\|_{L_1} \rightarrow 0 \implies \|f_n^+ - f^+\|_{H_2} \rightarrow 0. \quad (6)$$

A proof of an analog of (6) for more general matrix case can be found in [3] or [10]. In the present paper, we discuss quantitative estimates of the rate in the above convergence. Firstly, we look for estimates of  $\|g^+ - f^+\|_{H_2}$  in terms of  $\|g - f\|_{L_1}$  and  $\|\log g - \log f\|_{L_1}$ . It turns out that, in general, there is no such estimate. Namely, there is no function  $\Pi : [0, +\infty)^2 \rightarrow [0, +\infty)$  such that  $\lim_{s,t \rightarrow 0} \Pi(s, t) = 0$  for which the estimate

$$\|g^+ - f^+\|_{H_2}^2 \leq \Pi(\|g - f\|_{L_1}, \|\log g - \log f\|_{L_1})$$

holds for all  $f, g \geq 0$  with  $\|f\|_{L_1}, \|g\|_{L_1} \leq 1$ .

**Theorem 1** *There exist functions  $f_n, g_n \geq 0, n \in \mathbb{N}$ , such that*

$$\|f_n\|_{L_1}, \|g_n\|_{L_1} \leq 1, \|g_n - f_n\|_{L_1} \leq \frac{1}{n}, \|\log g_n - \log f_n\|_{L_1} \leq \frac{1}{n},$$

but  $\|g_n^+ - f_n^+\|_{H_2} \geq 2 - 1/n$ .

Nevertheless, one can still obtain an estimate for  $\|g^+ - f^+\|_{H_2}$  if one takes into account that for each  $f \in L_1(\mathbb{T})$  there exists an Orlicz space  $L_\Psi(\mathbb{T})$  such that  $f \in L_\Psi(\mathbb{T})$  (see, e.g., [17, Sect. 8]). One can show that there exists a function  $\Pi_\Psi : [0, +\infty) \rightarrow [0, +\infty)$  such that  $\lim_{t \rightarrow 0} \Pi_\Psi(t) = 0$  and

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + \|f\|_\Psi \Pi_\Psi(\|\log f - \log g\|_{L_1}) \quad (7)$$

(see Theorem 3 below). The estimate becomes particularly simple if  $f \in L_\infty(\mathbb{T})$ .

**Theorem 2** *Let  $f$  and  $g$  be arbitrary spectral densities for which  $f^+$  and  $g^+$  exist. Then,*

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + 2.5\|f\|_{L_\infty} \|\log f - \log g\|_{L_1}.$$

The paper is organized as follows. In Sect. 2, we prove (7) and Theorem 2. Theorem 1 is proved in Sect. 3.

This paper is a preliminary step toward the investigation of similar problems in the more complicated matrix case, which is going to be the subject of a forthcoming paper.

## 2 Positive results

Let  $K$  be the best constant in Kolmogorov's weak type (1, 1) inequality

$$m\{\vartheta \in [-\pi, \pi) : |\tilde{\psi}(\vartheta)| \geq \lambda\} \leq \frac{K}{\lambda} \int_{-\pi}^{\pi} |\psi(\vartheta)| d\vartheta, \quad \lambda > 0, \quad \psi \in L_1[-\pi, \pi),$$

where  $m$  stands for the Lebesgue measure on the real line. It is known that  $K = (1 + 3^{-2} + 5^{-2} + \dots)/(1 - 3^{-2} + 5^{-2} - \dots) \approx 1.347$  (see [9]).

**Lemma 1** *Let  $G : [0, +\infty) \rightarrow [0, +\infty)$  be a bounded absolutely continuous function that attains its maximum at point  $a \in (0, +\infty)$  (and possibly elsewhere) and is nondecreasing on  $[0, a]$ . Suppose  $G(0) = 0$  and*

$$I(G) := \int_0^a G'(\lambda) \frac{d\lambda}{\lambda} < +\infty.$$

Then,

$$\int_{-\pi}^{\pi} G(|\tilde{\psi}(\vartheta)|) d\vartheta \leq KI(G)\|\psi\|_{L_1}.$$

*Proof* Let

$$\mu_{\psi}(\lambda) := |\{\vartheta \in [-\pi, \pi) : |\tilde{\psi}(\vartheta)| \geq \lambda\}|.$$

Using Kolmogorov's weak type (1, 1) estimate with constant  $K$ , one gets

$$\begin{aligned} \int_{-\pi}^{\pi} G(|\tilde{\psi}(\vartheta)|) d\vartheta &\leq \int_{|\tilde{\psi}(\vartheta)| < a} G(|\tilde{\psi}(\vartheta)|) d\vartheta + G(a)\mu_{\psi}(a) \\ &= \int_{|\tilde{\psi}(\vartheta)| < a} \int_0^{|\tilde{\psi}(\vartheta)|} G'(\lambda) d\lambda d\vartheta + G(a)\mu_{\psi}(a) \\ &= \int_0^a G'(\lambda)(\mu_{\psi}(\lambda) - \mu_{\psi}(a)) d\lambda + G(a)\mu_{\psi}(a) \\ &= \int_0^a G'(\lambda)\mu_{\psi}(\lambda) d\lambda - G(a)\mu_{\psi}(a) + G(a)\mu_{\psi}(a) \\ &= \int_0^a G'(\lambda)\mu_{\psi}(\lambda) d\lambda \leq K\|\psi\|_{L_1} \int_0^a G'(\lambda) \frac{d\lambda}{\lambda} \\ &= KI(G)\|\psi\|_{L_1}. \end{aligned}$$

□

We need some notation from the theory of Orlicz spaces (see [17, 18]). Let  $\Phi$  and  $\Psi$  be mutually complementary  $N$ -functions, i.e.,

$$\Phi(x) = \int_0^{|x|} u(t) dt \quad \text{and} \quad \Psi(x) = \int_0^{|x|} v(t) dt,$$

where  $u : [0, \infty) \rightarrow [0, \infty)$  is a right continuous, nondecreasing function with  $u(0) = 0$  and  $u(\infty) := \lim_{t \rightarrow \infty} u(t) = \infty$ , and  $v$  is defined by the equality  $v(x) = \sup_{u(t) \leq x} t$ . Let  $(\Omega, \Sigma, \mu)$  be a measure space, and let  $L_\Phi(\Omega), L_\Psi(\Omega)$  be the corresponding Orlicz spaces, i.e.,  $L_\Phi(\Omega)$  is the set of measurable functions on  $\Omega$  for which either of the following norms

$$\|f\|_\Phi = \sup \left\{ \left| \int_\Omega fg d\mu \right| : \int_\Omega \Psi(g) d\mu \leq 1 \right\}$$

or

$$\|f\|_{(\Phi)} = \inf \left\{ \kappa > 0 : \int_\Omega \Phi \left( \frac{f}{\kappa} \right) d\mu \leq 1 \right\}$$

is finite. Note that these two norms are equivalent, namely (see, e.g., [17, (9.24)] or [18, Sect. 3.3, (14)])

$$\|f\|_{(\Phi)} \leq \|f\|_\Phi \leq 2\|f\|_{(\Phi)}, \quad \forall f \in L_\Phi(\Omega).$$

We will use the following Hölder inequality (see, e.g., [17, (9.27)] or [18, Sect. 3.3, (16)])

$$\left| \int_\Omega fg d\mu \right| \leq \|f\|_\Psi \|g\|_{(\Phi)}. \tag{8}$$

For an  $N$ -function  $\Phi$ , let

$$\Lambda_\Phi(s) := \inf \left\{ t > 0 : \frac{1}{t} \Phi' \left( \frac{1}{t} \right) \leq \frac{1}{s} \right\}, \quad s > 0. \tag{9}$$

If  $\Phi'$  is continuous, the above definition of  $\Lambda_\Phi$  can be rewritten in terms of inverse functions, because  $\Phi'$  is nondecreasing. For an arbitrary  $N$ -function  $\Phi$ , one has

$$\tau \Phi'(\tau) \leq \int_\tau^{2\tau} \Phi'(x) dx = \Phi(2\tau) - \Phi(\tau) < \Phi(2\tau).$$

Hence,

$$\Lambda_\Phi(s) \leq \frac{2}{\Phi^{-1} \left( \frac{1}{s} \right)}, \quad s > 0.$$

It is clear that

$$\Lambda_\Phi(s) \rightarrow 0 \quad \text{as } s \rightarrow 0+. \tag{10}$$

Also,

$$\Phi(\tau) \equiv \tau^q/q, \quad 1 < q < \infty \implies \Lambda_\Phi(s) \equiv s^{1/q}. \tag{11}$$

**Lemma 2** *For every  $N$ -function  $\Phi$ , the following estimate holds*

$$\|1 - \cos \tilde{\psi}\|_{(\Phi)} \leq 2\Lambda_\Phi(K_0\|\psi\|_{L_1}), \quad \forall \psi \in L_1, \tag{12}$$

where

$$K_0 := \frac{K}{2} \int_0^\pi \frac{\sin \lambda}{\lambda} d\lambda < 1.25 \tag{13}$$

and  $K$  is the same as in Lemma 1.

*Proof* We will use Lemma 1 with

$$G(\lambda) = \Phi \left( \frac{1 - \cos \lambda}{\kappa} \right)$$

and  $a = \pi$ . We have

$$I(G) = \frac{1}{\kappa} \int_0^\pi \Phi' \left( \frac{1 - \cos \lambda}{\kappa} \right) \sin \lambda \frac{d\lambda}{\lambda} \leq \frac{1}{\kappa} \Phi' \left( \frac{2}{\kappa} \right) \int_0^\pi \frac{\sin \lambda}{\lambda} d\lambda.$$

Hence,

$$\int_{-\pi}^\pi \Phi \left( \frac{1 - \cos \tilde{\psi}(\vartheta)}{\kappa} \right) d\vartheta \leq \frac{2}{\kappa} \Phi' \left( \frac{2}{\kappa} \right) \frac{K}{2} \int_0^\pi \frac{\sin \lambda}{\lambda} d\lambda \|\psi\|_{L_1}. \tag{14}$$

Taking  $\kappa > 2\Lambda_\Phi (K_0 \|\psi\|_{L_1})$ , we observe that the right-hand side of inequality (14) does not exceed 1 by virtue of the definition (9). Thus, (12) follows.  $\square$

**Lemma 3**

$$\|1 - \cos \tilde{\psi}\|_{L_1} \leq 2K_0 \|\psi\|_{L_1}, \quad \forall \psi \in L_1, \tag{15}$$

where  $K_0$  is defined by (13).

*Proof* We will use Lemma 1 with

$$G(\lambda) = 1 - \cos \lambda$$

and  $a = \pi$ . We have

$$I(G) = \int_0^\pi \frac{\sin \lambda}{\lambda} d\lambda$$

and

$$\|1 - \cos \tilde{\psi}\|_{L_1} = \int_{-\pi}^\pi (1 - \cos \tilde{\psi}(\vartheta)) d\vartheta \leq K \int_0^\pi \frac{\sin \lambda}{\lambda} d\lambda \|\psi\|_{L_1},$$

which implies (15).  $\square$

**Theorem 3** For every pair  $\Phi$  and  $\Psi$  of mutually complementary  $N$ -functions, the following estimate holds

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + 4\|f\|_\Psi \Lambda_\Phi \left( \frac{K_0}{2} \|\log f - \log g\|_{L_1} \right),$$

where  $K_0$  is the same as in Lemma 2.

*Proof*

$$\begin{aligned}
 \|f^+ - g^+\|_{H_2}^2 &= \|f^+\|_{H_2}^2 + \|g^+\|_{H_2}^2 \\
 &\quad - 2\operatorname{Re} \int_{-\pi}^{\pi} f^{1/2}(\vartheta)g^{1/2}(\vartheta) \exp\left(\frac{i}{2}(\log f(\vartheta) - \log g(\vartheta))\right) d\vartheta \\
 &= \|f^{1/2}\|_{L_2}^2 + \|g^{1/2}\|_{L_2}^2 - 2 \int_{-\pi}^{\pi} f^{1/2}(\vartheta)g^{1/2}(\vartheta) d\vartheta \\
 &\quad + 2 \int_{-\pi}^{\pi} f^{1/2}(\vartheta)g^{1/2}(\vartheta) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))\right)\right) d\vartheta \\
 &= \|(f^{1/2} - g^{1/2})^2\|_{L_1} + 2 \int_{-\pi}^{\pi} f^{1/2}(\vartheta) \\
 &\quad \times (g^{1/2}(\vartheta) - f^{1/2}(\vartheta)) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))\right)\right) d\vartheta \\
 &\quad + 2 \int_{-\pi}^{\pi} f(\vartheta) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))\right)\right) d\vartheta \\
 &\leq \int_{-\pi}^{\pi} (g^{1/2}(\vartheta) - f^{1/2}(\vartheta))^2 d\vartheta \\
 &\quad + 4 \int_{-\pi}^{\pi} f^{1/2}(\vartheta) (g^{1/2}(\vartheta) - f^{1/2}(\vartheta))_+ d\vartheta \\
 &\quad + 2 \int_{-\pi}^{\pi} f(\vartheta) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))\right)\right) d\vartheta.
 \end{aligned}$$

Here and below, for  $x \in \mathbb{R}$ , we define  $x_+ := \max(x, 0)$ .

Using the Hölder inequality (8) and the elementary inequality

$$(a^{1/2} - b^{1/2})^2 + 2b^{1/2} (a^{1/2} - b^{1/2})_+ \leq |a - b|, \quad \forall a, b \geq 0,$$

which is easily proved by considering the cases  $a \geq b$  and  $a < b$  separately, one gets

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + 2\|f\|_{\Psi} \left\| 1 - \cos\left(\frac{1}{2}(\log f - \log g)\right) \right\|_{(\Phi)}.$$

It is now left to apply Lemma 2 with  $\psi = \frac{1}{2}(\log f - \log g)$ .  $\square$

Since every integrable function belongs to a certain Orlicz space (see [17, Sect. 8]), Theorem 3 with an appropriate pair  $\Phi$  and  $\Psi$  of mutually complementary  $N$ -functions applies to any nonnegative integrable function  $f$  with an integrable logarithm. The condition (10) is fulfilled as well.



**Corollary 1** For every  $p \in (1, \infty)$ , there exists a constant  $C(p)$  such that

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + C(p)\|f\|_{L_p}\|\log f - \log g\|_{L_1}^{\frac{p-1}{p}}.$$

One can take  $C(p) = 2^{\frac{p+1}{p}} K_0^{\frac{p-1}{p}} \left(\frac{p}{p-1}\right)^{\frac{p-1}{p}}$ , where  $K_0$  is defined by (13).

*Proof* It is sufficient to take  $\Psi(t) \equiv t^p/p, \Phi(t) \equiv t^q/q, 1 < p < \infty, q = p/(p-1)$  in Theorem 3 and to apply (11) and [17, (9.7)]. □

**Corollary 2** There exists a constant  $C$  such that

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + C\|f\|_{L_\infty}\|\log f - \log g\|_{L_1}.$$

One can take  $C = 2K_0 < 2.5$ , where  $K_0$  is defined by (13).

*Proof* It follows from the proof of Theorem 3 that

$$\|f^+ - g^+\|_{H_2}^2 \leq 2\|f - g\|_{L_1} + 2\|f\|_{L_\infty} \left\| 1 - \cos\left(\frac{1}{2}(\log f - \log g)^\sim\right) \right\|_{L_1}.$$

It is now left to apply Lemma 3. □

### 3 Negative results

In this section, we prove Theorem 1.

*Proof* It follows from the proof of Theorem 3 that for any  $f, g \geq 0$  one has

$$\begin{aligned} \|f^+ - g^+\|_{H_2}^2 &= \|(f^{1/2} - g^{1/2})^2\|_{L_1} \\ &+ 2 \int_{-\pi}^{\pi} f^{1/2}(\vartheta) \left(g^{1/2}(\vartheta) - f^{1/2}(\vartheta)\right) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))^\sim\right)\right) d\vartheta \\ &+ 2 \int_{-\pi}^{\pi} f(\vartheta) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))^\sim\right)\right) d\vartheta \\ &\geq -4 \int_{-\pi}^{\pi} f^{1/2}(\vartheta) \left|g^{1/2}(\vartheta) - f^{1/2}(\vartheta)\right| d\vartheta \\ &+ 2 \int_{-\pi}^{\pi} f(\vartheta) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))^\sim\right)\right) d\vartheta \\ &\geq 2 \int_{-\pi}^{\pi} f(\vartheta) \left(1 - \cos\left(\frac{1}{2}(\log f(\vartheta) - \log g(\vartheta))^\sim\right)\right) d\vartheta - 4\|f - g\|_{L_1}. \end{aligned}$$

Let  $w_n$  be a conformal mapping of the unit disk onto the ellipse with the axes

$$[-\varepsilon_n, 0] \quad \text{and} \quad -\varepsilon_n/2 + i[-2\pi, 2\pi],$$

such that  $w_n(0) = -\varepsilon_n/2$ , where  $\varepsilon_n = \frac{1}{2\pi n}$ . Let  $h_n := \exp(\operatorname{Re} w_n)$ . Then  $(\log h_n)^\sim = (\operatorname{Re} w_n)^\sim = \operatorname{Im} w_n$  and therefore  $\|1 - \cos(\frac{1}{2}(\log h_n)^\sim)\|_{L_\infty} = 2$ .

Due to duality considerations, there exists  $f_n^0 \geq 0$  such that  $\|f_n^0\|_{L_1} = 1$  and

$$\begin{aligned} & \int_{-\pi}^{\pi} f_n^0(\vartheta) \left( 1 - \cos\left(\frac{1}{2}(\log h_n(\vartheta))^\sim\right) \right) d\vartheta \\ & \geq \left(1 - \frac{\varepsilon_n}{2}\right) \left\| 1 - \cos\left(\frac{1}{2}(\log h_n)^\sim\right) \right\|_{L_\infty}. \end{aligned}$$

If  $\log f_n^0 \in L_1$ , we take  $f_n = f_n^0$ . Otherwise, we define  $f_n = (1 - \varepsilon_n/2)f_n^0 + \varepsilon_n/4\pi$ . Then,  $\|f_n\|_{L_1} = 1$ , and  $(1 - \varepsilon_n/2)^2 > 1 - \varepsilon_n$  implies that

$$\begin{aligned} & \int_{-\pi}^{\pi} f_n(\vartheta) \left( 1 - \cos\left(\frac{1}{2}(\log h_n(\vartheta))^\sim\right) \right) d\vartheta \\ & \geq (1 - \varepsilon_n) \left\| 1 - \cos\left(\frac{1}{2}(\log h_n)^\sim\right) \right\|_{L_\infty}. \end{aligned}$$

Finally, let  $g_n = h_n f_n$ . Then,  $0 \leq g_n \leq f_n$ ,  $\|g_n\|_{L_1} \leq 1$ ,

$$\|f_n - g_n\|_{L_1} = \|f_n(1 - h_n)\|_{L_1} \leq \|1 - h_n\|_{L_\infty} \leq 1 - e^{-\varepsilon_n} \leq \varepsilon_n < \frac{1}{2n},$$

$$\|\log f_n - \log g_n\|_{L_1} = \|\log h_n\|_{L_1} \leq 2\pi \|\log h_n\|_{L_\infty} = 2\pi \varepsilon_n = \frac{1}{n},$$

and

$$\begin{aligned} \|f_n^+ - g_n^+\|_{H_2}^2 & \geq 2(1 - \varepsilon_n) \left\| 1 - \cos\left(\frac{1}{2}(\log h_n)^\sim\right) \right\|_{L_\infty} - 4\|f - g\|_{L_1} \\ & > 4(1 - \varepsilon_n) - \frac{2}{n} > 4\left(1 - \frac{1}{4n}\right) - \frac{2}{n} \geq \left(2 - \frac{1}{n}\right)^2. \end{aligned}$$

□

*Remark* The norms  $\|\log f_n\|_{L_1}$  and  $\|\log g_n\|_{L_1}$  might not be bounded in Theorem 1. Let  $f_n^0$  be the function from the above proof. Changing the definition of  $f_n$  in the proof to  $f_n = f_n^0 + 1$ , one can change the estimates  $\|f_n\|_{L_1}, \|g_n\|_{L_1} \leq 1$  in the theorem for  $\|f_n\|_{L_1} = 2\pi + 1, \|g_n\|_{L_1} \leq 2\pi + 1, \|\log f_n\|_{L_1} \leq 1$ .

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

1. Anderson, B.D.O.: Continuity of the spectral factorization operation. *Mat. Apl. Comput.* **4**(2), 139–156 (1985)
2. Anderson, B.D.O., Moore, J.B.: *Linear Optimal Control*. Prentice-Hall Inc, Englewood Cliffs (1971)
3. Barclay, S.: Continuity of the spectral factorization mapping. *J. Lond. Math. Soc.* **70**(3), 763–779 (2004)
4. Bart, H., Gohberg, I., Kaashoek, M.A., Ran, A.C.M.: A state space approach to canonical factorization with applications. In: *Operator Theory: Advances and Applications*, vol. 200. Linear Operators and Linear Systems. Birkhäuser Verlag, Basel (2010)
5. Boche, H., Pohl, V.: Behavior of the spectral factorization for continuous spectral densities. *Signal Process.* **87**(5), 1078–1088 (2007)
6. Boche, H., Pohl, V.: Spectral factorization for polynomial spectral densities—impact of dimension. *IEEE Trans. Inform. Theory* **53**(11), 4236–4241 (2007)
7. Boche, H., Pohl, V.: Robustness of the spectral factorization for polynomials. In: 7th International ITG Conference on Source and Channel Coding, SCC 2008, pp. 1–6. CD-ROM (2008)
8. Daubechies, I.: Ten lectures on wavelets. In: *CBMS-NSF Regional Conference Series in Applied Mathematics*, vol. 61. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA (1992)
9. Davis, B.: On the weak type  $(1, 1)$  inequality for conjugate functions. *Proc. Am. Math. Soc.* **44**, 307–311 (1974)
10. Ephremidze, L., Janashia, G., Lagvilava, E.: On approximate spectral factorization of matrix functions. *J. Fourier Anal. Appl.* **17**(5), 976–990 (2011)
11. Fischer, R.F.H.: *Precoding and Signal Shaping for Digital Transmission*. Wiley-IEEE Press, Piscataway (2002)
12. Francis, B.A.: *A Course in  $H_\infty$  Control Theory*. Lecture Notes in Control and Information Sciences, vol. 88. Springer-Verlag, Berlin (1987)
13. Gohberg, I.C., Kreĭn, M.G.: Systems of integral equations on the half-line with kernels depending on the difference of the arguments. *Uspekhi Mat. Nauk (N.S.)* **13**(2), 3–72 (1958)
14. Jacob, B., Partington, J.R.: On the boundedness and continuity of the spectral factorization mapping. *SIAM J. Control Optim.* **40**(1), 88–106 (2001). (electronic)
15. Kailath, T., Hassibi, B., Sayed, A.H.: *Linear Estimation*. Prentice-Hall Information and System Sciences Series. Prentice-Hall, Inc., Englewood Cliffs (1999)
16. Kolmogoroff, A.N.: Stationary sequences in Hilbert's space. *Bolletin Moskovskogo Gosudarstvenogo Universiteta. Matematika* **2**, 40 (1941)
17. Krasnosel'skiĭ, M.A., Rutickiĭ, J.B.: *Convex functions and Orlicz spaces*. Translated from the first Russian edition by Leo F. Boron. P. Noordhoff Ltd., Groningen (1961)
18. Rao, M.M., Ren, Z.D.: *Theory of Orlicz spaces, Monographs and Textbooks in Pure and Applied Mathematics*, vol. 146. Marcel Dekker Inc, New York (1991)
19. Resnikoff, H.L., Wells Jr., R.O.: *Wavelet Analysis. The Scalable Structure of Information*. Springer-Verlag, New York (1998)
20. Strang, G., Nguyen, T.: *Wavelets and Filter Banks*. Wellesley-Cambridge Press, Wellesley (1996)
21. Wiener, N.: *Extrapolation, Interpolation, and Smoothing of Stationary Time Series. With Engineering Applications*. The Technology Press of the Massachusetts Institute of Technology, Cambridge; John Wiley & Sons Inc., New York; Chapman & Hall Ltd., London (1949)