



ELSEVIER

Journal of Pure and Applied Algebra 175 (2002) 187–205

JOURNAL OF
PURE AND
APPLIED ALGEBRA

www.elsevier.com/locate/jpaa

Finite preorders and Topological descent I

George Janelidze^{a,1}, Manuela Sobral^{b,*,2}

^aMathematics Institute of the Georgian Academy of Sciences, 1 M. Alexidze Str., 390093 Tbilisi, Georgia

^bDepartamento de Matemática, Universidade de Coimbra, 3000 Coimbra, Portugal

Received 28 December 2000

Communicated by A. Carboni

Dedicated to Max Kelly in honour of his 70th birthday

Abstract

It is shown that the descent constructions of finite preorders provide a simple motivation for those of topological spaces, and new counter-examples to open problems in Topological descent theory are constructed.

© 2002 Published by Elsevier Science B.V.

MSC: 18B30; 18A20; 18D30; 18A25

0. Introduction

Let $\mathcal{T}op$ be the category of topological spaces. For a given continuous map $p: E \rightarrow B$, it might be possible to describe the category $(\mathcal{T}op \downarrow B)$ of bundles over B in terms of $(\mathcal{T}op \downarrow E)$ using the pullback functor

$$p^* : (\mathcal{T}op \downarrow B) \rightarrow (\mathcal{T}op \downarrow E), \quad (0.1)$$

in which case p is called an *effective descent morphism*. There are various ways to make this precise (see [8,9]); one of them is described in Section 3.

* Corresponding author. Dept. de Matemática, Univ. de Coimbra, 3001-454 Coimbra, Portugal.

E-mail addresses: gjanel@rmi.acnet.ge (G. Janelidze), sobral@mat.uc.pt (M. Sobral).

¹ This work was pursued while the first author was visiting the University of Coimbra in October/December 1998 under the support of CMUC/FCT.

² Partial financial support by PRAXIS Project PCTEX/P/MAT/46/96 and by CMUC/FCT is gratefully acknowledged.

More generally, the same notion can still be considered when (0.1) is replaced by

$$p^* : \mathbf{A}^B \rightarrow \mathbf{A}^E, \quad (0.2)$$

where \mathbf{A} is any $\mathcal{T}op$ -indexed category, or even also the category $\mathcal{T}op$ replaced by an arbitrary category \mathcal{C} ; it is still useful to think of the objects of \mathbf{A}^B as a kind of bundles over B , possibly with additional structure.

There is also an “intermediate” level of generality, where each \mathbf{A}^B is a full subcategory in $(\mathcal{T}op \downarrow B)$ determined by a class \mathbb{E} of morphisms in $\mathcal{T}op$. The corresponding effective descent morphisms are called the *effective \mathbb{E} -descent morphisms*.

The main problem studied in *Topological descent theory* is to find out, for given classes \mathbb{D} and \mathbb{E} of continuous maps, if every $p \in \mathbb{D}$ is an effective \mathbb{E} -descent morphism.

Let us recall the main known results of this type (in chronological order):

- A continuous map $p : E \rightarrow B$ is said to be locally sectionable if every point in B has an open neighbourhood U such that the map $p^{-1}(U) \rightarrow U$ induced by p has a continuous section. Every locally sectionable map is an effective descent morphism [7].
- Every open surjective map is an effective descent morphism (Sobral, see [17]; as observed in [10] it can also be easily deduced from Moerdijk’s axioms [12]—just like it is deduced there for locales).
- Every proper map is an effective descent morphism (Moerdijk, Vermeulen [18]; see also [14]).
- Reiterman and Tholen [14] finally solved the problem of characterizing the effective descent morphisms in $\mathcal{T}op$ and gave a first example of *non-effective* descent morphism.
- Every effective descent morphism is also an effective étale-descent morphism [8]. (As T. Plewe observed later, there is a simple purely categorical proof of this fact.)
- Every triquotient map is an effective descent morphism [13], but there are counter-examples for the converse; yet, the class of triquotient maps contains all locally sectionable, all open surjections and all proper maps.
- Effective descent morphisms are stable under pullback in categories with pullbacks and coequalizers of certain naturally arising equivalence relations [17]. This result was generalized to effective \mathbb{E} -descent morphisms in [15].
- A morphism is an effective descent morphism if and only if every pullback of it is an effective bijective-descent map [16].
- There are simple examples of non-effective descent morphisms [16].

Analyzing the finite counter-examples of [15,16] we arrived at the conclusion that most of the phenomena which occur in difficult problems and proofs of Topological descent are easily detectable and easily understandable already on the level of finite topological spaces—and since those are just finite preorders, a lot of standard arguments can be used!

Accordingly, in this paper we develop the very simple descent theory of (finite) preorders, and then explain that Topological descent theory is just an infinite extension of it. We also construct new finite counter-examples to some problems of Topological descent theory.

In order to convince the readers interested in topological descent that “they must immediately interrupt their work and read our paper” let us point out the following:

The Reiterman–Tholen characterization of effective descent topological maps mentioned above says:

Theorem 0.1. *The map $p:E \rightarrow B$ is an effective descent morphism if and only if every crest of ultrafilters in B has a lifting along p (see [14] for details).*

In the case of finite topological spaces, which are exactly the finite preorders, it reduces to:

Theorem 0.2. *The map $p:E \rightarrow B$ is an effective descent morphism if and only if for every chain $b_2 \leq b_1 \leq b_0$ in B there exists $e_2 \leq e_1 \leq e_0$ in E with $p(e_i) = b_i$, for $i = 0, 1, 2$.*

The paper is organized as follows:

0. Introduction
1. Finite topological spaces
2. Quotient and Day–Kelly maps
3. Effective descent morphisms
4. Generalized descent
5. Bijective descent
6. Étale descent
7. Triquotient maps
8. Counter-examples
9. Remarks on infinite topological spaces

Note that the results of Sections 2 and 3 in some sense go back to Giraud [3], and are closely related to the similar results for categories (although they are not straightforward consequences of those). A general approach to descent constructions for internal category-like structures is developed by Gran [4] (“Maltsev case”) and Le Creurer [11] (“lexensive” case). Since the category of sets is lexensive, the results of [11] could be used here; however, we give independent proofs in order to make the paper self-contained.

The results of this paper were presented on the International Category Theory Meeting held in Coimbra in July 1999, and first appeared as the preprint [5]. In addition we are going to give an elementary characterization of effective étale-descent morphisms of finite topological spaces in [6]. Note also that the converse of Proposition 7.1 was proved by Clementino [1] providing a characterization of triquotient maps between finite spaces.

1. Finite topological spaces

Finite topological spaces have the “open closure operator”. That is, for every subset X of a finite topological space A , there is a smallest open set $\downarrow X$ containing X .

Moreover,

$$\downarrow X = \bigcup_{x \in X} \downarrow x, \tag{1.1}$$

where $\downarrow x = \downarrow \{x\}$.

We write

$$y \rightarrow x \Leftrightarrow y \in \downarrow x; \tag{1.2}$$

in classical notation our $y \rightarrow x$ would be just $y, y, \dots \rightarrow x$.

Proposition 1.1. *If A is a finite topological space, then \rightarrow is a preorder, i.e. it is reflexive and transitive:*

$$x \rightarrow x, \tag{1.3}$$

$$z \rightarrow y \rightarrow x \Rightarrow z \rightarrow x \tag{1.4}$$

for every $x, y, z \in A$. This determines an isomorphism

$$FinTop \cong FinPreord \tag{1.5}$$

between the category of finite topological spaces and the category of finite preordered sets.

It is also well-known that (1.5) extends to an isomorphism between *Preord* and the category of topological spaces for which the set of open subsets is closed under intersection.

Since

$$\downarrow x = \{y \in A \mid y \rightarrow x\}, \tag{1.6}$$

we also introduce

$$\uparrow x = \{y \in A \mid x \rightarrow y\} \tag{1.7}$$

and we have $\uparrow x = \overline{\{x\}}$, the closure of $\{x\}$.

Proposition 1.2. *Let A and A' be topological spaces with the same underlying set. The following conditions are equivalent:*

- (a) *a subset X is open in A if and only if it is closed in A' ;*
- (b) *the preorders in A and A' are opposite to each other, i.e. $y \rightarrow x$ in A if and only if $x \rightarrow y$ in A' .*

According to (1.5), a map $\alpha: A \rightarrow B$ of finite topological spaces is continuous if and only if it is a monotone map (i.e. $y \rightarrow x \Rightarrow \alpha(y) \rightarrow \alpha(x)$) of the corresponding preordered sets.

For such a map α we also have

Proposition 1.3. *The following conditions are equivalent:*

- (a) *α is a proper map;*

- (b) α is a closed map;
- (c) $\alpha(\uparrow x)$ is closed in B for every $x \in A$;
- (d) $\alpha(\uparrow x) = \uparrow \alpha(x)$ for every $x \in A$;
- (e) for every $x \in A$ and $\alpha(x) \rightarrow b$ in B , there exists $a \in A$ with $x \rightarrow a$ and $\alpha(a) = b$.

Proposition 1.4. *The following conditions are equivalent:*

- (a) α is an open map;
- (b) $\alpha(\downarrow x)$ is open in B for every $x \in A$;
- (c) $\alpha(\downarrow x) = \downarrow \alpha(x)$ for every $x \in A$;
- (d) for every $x \in A$ and $b \rightarrow \alpha(x)$ in B , there exists $a \in A$ with $a \rightarrow x$ and $\alpha(a) = b$.

Proposition 1.5. *The following conditions are equivalent:*

- (a) α is an étale map (i.e. a local homomorphism);
- (b) α is an open map and its restriction to $\downarrow x$ is injective for every $x \in A$;
- (c) the map $\downarrow x \rightarrow \downarrow \alpha(x)$ induced by α is bijective for every $x \in A$;
- (d) for every $x \in A$ and $b \rightarrow \alpha(x)$ in B there exists a unique $a \in A$ with $a \rightarrow x$ and $\alpha(a) = b$.

2. Quotient and Day–Kelly maps

Let Rel be the category of pairs $A = (A, R_A)$, where $R_A \subseteq A \times A$ is an arbitrary binary relation on A . The “quotient maps” in this category have a simple description:

Proposition 2.1. *For a morphism $\alpha: A \rightarrow B$ with $\alpha(A) = B$, the following conditions are equivalent:*

- (a) α is a regular epimorphism, i.e.

$$A \cdot_B A \begin{array}{c} \xrightarrow{\pi_1} \\ \xrightarrow{\pi_2} \end{array} A \xrightarrow{\alpha} B \tag{2.1}$$

is a coequalizer diagram in Rel ;

- (b) R_B is the smallest relation on B which makes $\alpha: A \rightarrow B$ a morphism in Rel ;
- (c) $R_B = (\alpha \times \alpha)(R_A)$, the image of $R_A \subseteq A \times A$ under the map $\alpha \times \alpha: A \times A \rightarrow B \times B$;
- (d) $b'R_B b$ if and only if there exist $a', a \in A$ with $\alpha(a') = b'$, $\alpha(a) = b$ and $a'R_A a$.

Exactly the same is true in the category $ReflRel$ of sets equipped with a reflexive relation, but not in $Preord$ —since transitivity of R_A in (2.1) does not imply transitivity of R_B . However, given such a coequalizer diagram in $ReflRel$ with transitive R_A , we obtain a coequalizer diagram in $Preord$ just by taking the transitive closure of R_B . Therefore we have

Proposition 2.2. *For a morphism $\alpha: A \rightarrow B$ in $Preord$ (or in $FinPreord$) with $\alpha(A) = B$, the following conditions are equivalent:*

- (a) α is a regular epimorphism;
- (b) R_B is the smallest relation on B which makes $\alpha: A \rightarrow B$ a morphism in $Preord$;

- (c) R_B is the transitive closure of $(\alpha \times \alpha)(R_A)$;
- (d) $b'R_B b$ if and only if there exists a (finite) sequence $(a'_1, a_1), \dots, (a'_n, a_n) \in R_A$ with $b' = \alpha(a'_1), \alpha(a_i) = \alpha(a'_{i+1}),$ for $i = 1, \dots, n - 1,$ and $\alpha(a_n) = b.$

The fact that regular epimorphisms in *Rel* and *ReflRel* are “better” than in *Preord* can also be expressed categorically:

Proposition 2.3. (a) *The regular epimorphisms in Rel and ReflRel are pullback stable, i.e. if*

$$\begin{array}{ccc}
 E \times_B A & \xrightarrow{\pi_2} & A \\
 \pi_1 \downarrow & & \downarrow \alpha \\
 E & \xrightarrow{p} & B
 \end{array} \tag{2.2}$$

is a pullback (in one of these categories) and p is a regular epimorphism, then so is π_2 ;

(b) *a morphism $p: E \rightarrow B$ in Preord is a pullback stable regular epimorphism if and only if it is a regular epimorphism in Rel (or, equivalently, in ReflRel).*

Proof. (a) follows from Proposition 2.1(a) \Leftrightarrow (c) and the fact that the regular epimorphisms in *Sets* are pullback stable.

Since every morphism in *Preord*, which is a regular epimorphism in *Rel*, must be a regular epimorphism in *Preord*, the “if” part of (b) follows from (a).

In order to prove the “only if” part of (b) we take:

- an arbitrary pair $(b', b) \in R_B$;
- $A = \{b, b'\}$ with the induced preorder;
- $\alpha: A \rightarrow B$ the inclusion map.

Since $\pi_2: E \times_B A \rightarrow A$ is a regular epimorphism, there exists a sequence

$$(x'_1, x_1), \dots, (x'_n, x_n) \in R_{E \times_B A}$$

with $b' = \pi_2(x'_1), \pi_2(x_i) = \pi_2(x'_{i+1}),$ for $i = 1, \dots, n - 1,$ and $\pi_2(x_n) = b.$ However, since there are no elements in A different from b and $b',$ this means that $b' = \pi_2(x'_k)$ and $\pi_2(x_k) = b$ for some k ($1 \leq k \leq n$). Therefore, the pair $(\pi_1(x'_k), \pi_1(x_k)) \in R_E$ has $p(\pi_1(x'_k)) = b'$ and $p(\pi_1(x_k)) = b$ as desired. \square

Remark 2.4. According to topological terminology, we say that $p: E \rightarrow B$ is a hereditary quotient map if, for every $B' \subseteq B$ with the induced preorder, the map $p^{-1}(B') \rightarrow B'$ induced by p is a quotient map (i.e. a regular epimorphism). Since in the proof of the “only if” part of Proposition 2.3(b) the morphism $\alpha: A \rightarrow B$ was an inclusion map with the induced order in $A,$ we conclude that the pullback stable regular epimorphisms in *Preord* are the same as the hereditary quotient maps.

Now we return to finite topological spaces.

A continuous map $p : E \rightarrow B$ is said to be a Day–Kelly map if for every $b \in B$ and every open covering family $(E_i)_{i \in I}$ of $p^{-1}(b)$ in E , there exists a finite set $\{i_1, \dots, i_n\}$ with

$$b \in \text{Int}(p(E_{i_1}) \cup \dots \cup p(E_{i_n})). \tag{2.3}$$

In the finite case this simplifies in the obvious way: we can just take I to be a one element set.

Since the Day–Kelly maps are known to be precisely the pullback stable regular epimorphisms of topological spaces (see [2,7])—or directly from the results above—we obtain:

Proposition 2.5. *For a morphism $p : E \rightarrow B$ in FinTop , the following conditions are equivalent:*

- (a) *p is a pullback stable regular epimorphism (in FinTop , or equivalently in Fin-Preord);*
- (b) *for every $b' \rightarrow b$ in B there exists $e' \rightarrow e$ in E with $p(e') = b'$ and $p(e) = b$;*
- (c) *p is a Day–Kelly map;*
- (d) *p is a hereditary quotient map;*
- (e) *for every $b \in B$ and open set $E' \subseteq E$ containing $p^{-1}(b)$, we have $b \in \text{Int}(p(E'))$.*

3. Effective descent morphisms

Various definitions of effective descent morphism are compared in [8,9]; one of them says that a morphism $p : E \rightarrow B$ in a category \mathcal{C} is an effective descent morphism if the pullback functor

$$p^* : (\mathcal{C} \downarrow B) \rightarrow (\mathcal{C} \downarrow E) \tag{3.1}$$

is monadic.

However, we will only need to know that the class of effective descent morphisms satisfies the following (see [8] for details):

Proposition 3.1. (a) *If \mathcal{C} has pullbacks and coequalizers (of equivalence relations), then every effective descent morphism in \mathcal{C} is a pullback stable regular epimorphism.*

(b) *If \mathcal{C} is exact, then the class of effective descent morphisms in \mathcal{C} coincides with the class of regular epimorphisms.*

Proposition 3.2. *Let \mathcal{C} and \mathcal{C}' be categories satisfying*

- (a) *\mathcal{C}' has pullbacks and coequalizers;*
- (b) *every regular epimorphism in \mathcal{C}' is an effective descent morphism;*
- (c) *\mathcal{C} is a full subcategory of \mathcal{C}' closed under pullbacks;*
- (d) *every pullback stable regular epimorphism in \mathcal{C} is a regular epimorphism in \mathcal{C}' .*

Then a pullback stable regular epimorphism $p : E \rightarrow B$ in \mathcal{C} is an effective descent morphism if and only if

$$E \times_B A \in \mathcal{C} \Rightarrow A \in \mathcal{C} \tag{3.2}$$

for every pullback (2.2) in \mathcal{C}' .

Using these two propositions, it is easy to characterize the effective descent morphisms in *Rel*, *ReflRel*, and *Preord*:

Proposition 3.3. *Every regular epimorphism in Rel is an effective descent morphism, and the same is true for ReflRel.*

Proof. An object (A, R_A) in *Rel* can be considered as a graph

$$R_A \rightrightarrows A \tag{3.3}$$

and we take $\mathcal{C} = Rel$ and \mathcal{C}' to be the category of graphs. Conditions 3.2(a)–(d) obviously hold (just note that Proposition 3.2(b) follows from Proposition 3.1(b) since now \mathcal{C} is a topos). Since implication (3.2) obviously holds as soon as p is an epimorphism in \mathcal{C}' , we conclude that every pullback stable regular epimorphism in *Rel* is an effective descent morphism—and then we apply Proposition 2.3(a).

The same arguments, but with reflexive graphs instead of graphs, can be used for reflexive relations. \square

Proposition 3.4. *For a morphism $p : E \rightarrow B$ in Preord (or in FinPreord) the following conditions are equivalent:*

- (a) p is an effective descent morphism;
- (b) p is a pullback stable regular epimorphism and, for every pullback (2.2) in *Rel* (or in *ReflRel*) with $E \times_B A$ a preorder, A also is a preorder;
- (c) for every $b_2 \rightarrow b_1 \rightarrow b_0$ in B there exists $e_2 \rightarrow e_1 \rightarrow e_0$ in E with $p(e_i) = b_i$, for $i = 0, 1, 2$.

Proof. (a) \Leftrightarrow (b) follows from the previous results. More precisely, we can apply Proposition 3.2 to $\mathcal{C} = Preord$, $\mathcal{C}' = Rel$ (or *ReflRel*) since in that case:

- Propositions 3.2(a) and 3.2(c) are obvious;
- Proposition 3.2(b) follows from Proposition 3.3;
- Proposition 3.2(d) follows from Proposition 2.3(b).

(c) \Rightarrow (b): Suppose p satisfies (c). Then p is a pullback stable regular epimorphism by Proposition 2.5(b) \Leftrightarrow (a), and we only need to show that, for every pullback (2.2) with transitive $R_E, R_B, R_{E \times_B A}$, the relation R_A is also transitive. However this is clear: given $a_2 \rightarrow a_1 \rightarrow a_0$ in A , there exists $e_2 \rightarrow e_1 \rightarrow e_0$ in E with $p(e_2) = \alpha(a_2)$, $p(e_1) = \alpha(a_1)$, $p(e_0) = \alpha(a_0)$ and hence $(e_2, a_2) \rightarrow (e_1, a_1) \rightarrow (e_0, a_0)$. Therefore $(e_2, a_2) \rightarrow (e_0, a_0)$, since $R_{E \times_B A}$ is transitive, which gives $a_2 \rightarrow a_0$ since $\pi_2 : E \times_B A \rightarrow A$ is morphism in *Rel*.

(b) \Rightarrow (c): Suppose p satisfies (b) and take:

- an arbitrary $b_2 \rightarrow b_1 \rightarrow b_0$ in B ;
- $A = \{a_0, a_1, a_2\}$ any three element set with $R_A = \Delta_A \cup \{(a_2, a_1), (a_1, a_0)\}$;
- $\alpha : A \rightarrow B$ with $\alpha(a_0) = b_0$, $\alpha(a_1) = b_1$ and $\alpha(a_2) = b_2$ (note that α need not be an injection!).

Since E and B are preorders, but A is not, $E \times_B A$ must not be a preorder. That is, there exist $x_0, x_1, x_2 \in E \times_B A$ with $(x_2, x_1), (x_1, x_0) \in R_{E \times_B A}$ but $(x_2, x_0) \notin R_{E \times_B A}$.

We have

$$E \times_B A = (p^{-1}(b_2) \times \{a_2\}) \cup (p^{-1}(b_1) \times \{a_1\}) \cup (p^{-1}(b_0) \times \{a_0\}) \tag{3.4}$$

and since $\pi_2 : E \times_B A \rightarrow A$ must be a regular epimorphism in Rel , it is easy to see that we must have $x_i \in p^{-1}(b_i) \times \{a_i\}$, for $i = 0, 1, 2$. After that, we take $e_2 = \pi_1(x_2)$, $e_1 = \pi_1(x_1)$ and $e_0 = \pi_1(x_0)$. \square

4. Generalized descent

Let \mathcal{C} be a category. Recall that a \mathcal{C} -indexed category \mathbf{A} consists of

- categories \mathbf{A}^B , defined for all objects B in \mathcal{C} ,
- functors $p^* : \mathbf{A}^B \rightarrow \mathbf{A}^E$, for all morphisms $p : E \rightarrow B$ in \mathcal{C} , and
- natural isomorphisms $\varphi_{p,q} : q^* p^* \rightarrow (pq)^*$ and $\psi_B : (1_B)^* \rightarrow 1_{\mathbf{A}^B}$, for all $q : F \rightarrow E$ and $p : E \rightarrow B$ in \mathcal{C} , with the standard coherence conditions.

For a given morphism $p : E \rightarrow B$ in a category \mathcal{C} with pullbacks and a \mathcal{C} -indexed category \mathbf{A} , the category $Des_{\mathbf{A}}(p)$ of \mathbf{A} -descent data for p is defined as a suitable 2-equalizer

$$Des_{\mathbf{A}}(p) \longrightarrow \mathbf{A}^E \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \mathbf{A}^{E \cdot_B E} \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \mathbf{A}^{E \cdot_B E \cdot_B E} \tag{4.1}$$

(described in [9] in the language of internal actions). The functor p^* has a canonical factorization

$$\begin{array}{ccc} \mathbf{A}^B & \xrightarrow{p^*} & \mathbf{A}^E \\ & \searrow K_{\mathbf{A}}^p & \nearrow U_{\mathbf{A}}^p \\ & Des_{\mathbf{A}}(p) & \end{array} \tag{4.2}$$

and p is said to be an effective \mathbf{A} -descent morphisms if K^p is a category equivalence.

In particular, any pullback stable class \mathbb{E} of morphisms in \mathcal{C} can be regarded as a \mathcal{C} -indexed category: we take

- $\mathbb{E}^B = \mathbb{E}(B)$ to be the full subcategory in $(\mathcal{C} \downarrow B)$ with objects all (A, α) with $\alpha : A \rightarrow B$ in \mathbb{E} ;
- $p^* : \mathbb{E}^B \rightarrow \mathbb{E}^E$ the pullback functor $(A, \alpha) \mapsto (E \times_B A, \pi_1)$ along $p : E \rightarrow B$;
- $\varphi_{p,q}$ and ψ_B the canonical isomorphisms $F \times_E (E \times_B (-)) \cong F \times_B (-)$ and $B \times_B (-) \cong (-)$, respectively.

The category $Des_{\mathbb{E}}(p)$ can be described as the category of triples $(C, \gamma, \xi) =$

$$E \times_B C \xrightarrow{\xi} C \xrightarrow{\gamma} E \tag{4.3}$$

such that $\gamma \in \mathbb{E}$ and the diagram

$$\begin{array}{ccc}
 E \cdot_B E \cdot_B C & \xrightarrow{1_E \cdot \zeta} & E \cdot_B C & \xleftarrow{\langle \gamma_1, 1_C \rangle} & C \\
 \langle \pi_1, \pi_3 \rangle \downarrow & & \zeta \downarrow & & \swarrow 1_C \\
 E \cdot_B C & \xrightarrow{\zeta} & C & & \\
 \pi_1 \downarrow & & \swarrow \gamma & & \\
 E & & & &
 \end{array} \tag{4.4}$$

commutes (we use the standard notation, writing π_i —here $i=1, 2$ or 3 —for all kinds of pullback projections; note also that the commutativity of the bottom triangle is already used in the square to make $1_E \times \zeta$ well defined).

If $\mathcal{C} = Rel$ or \mathcal{C} is any other concrete category considered in the previous sections, then we write

$$\zeta(e, c) = ec, \tag{4.5}$$

and the commutativity of (4.4) translates as

$$e(e'c) = ec,$$

$$\gamma(c)c = c,$$

$$\gamma(ec) = e. \tag{4.6}$$

The functor $K_{\mathbb{E}}^p : \mathbb{E}(B) \rightarrow Des_{\mathbb{E}}(p)$ is defined by

$$K_{\mathbb{E}}^p(A, \alpha) = (E \times_B A, \pi_1, \langle \pi_1, \pi_3 \rangle); \tag{4.7}$$

using the elements, $\langle \pi_1, \pi_3 \rangle : E \times_B (E \times_B A) \rightarrow E \times_B A$ would be written as

$$e(e', a) = (e, a) \tag{4.8}$$

If every $\delta : D \rightarrow E$ in \mathbb{E} gives $p\delta : E \rightarrow B$ in \mathbb{E} , then the diagram (4.2) (for $\mathbf{A} = \mathbb{E}$) can be identified with the standard diagram

$$\begin{array}{ccc}
 \mathbb{E}(B) & \xrightarrow{p^*} & \mathbb{E}(E) \\
 \text{comparison} \searrow & & \nearrow \text{forgetful} \\
 & & \mathbb{E}(E)^{\mathbb{T}}
 \end{array} \tag{4.9}$$

for the monad \mathbf{T} of the adjunction $p! \dashv p^*$.

And, of course, if \mathbb{E} is the class of all morphisms in \mathcal{C} , then an effective \mathbb{E} -descent morphism is the same as an effective descent morphism, as defined in the previous section.

5. Bijective descent

In this section \mathbb{E} denotes the class of morphisms in *Preord* which are bijections.

Proposition 5.1. *For a morphism $p: E \rightarrow B$ in *Preord*, the following conditions are equivalent:*

- (a) *p is a regular epimorphism in *Rel*;*
- (b) *for every pullback (2.2) with $\alpha \in \mathbb{E}$, the projection $\pi_2: E \times_B A \rightarrow A$ is a regular epimorphism in *Preord*.*

Proof. (a) \Rightarrow (b) follows from Proposition 2.3(b).

(b) \Rightarrow (a) can be proved with the same arguments as the “only if” part of Proposition 2.3(b), but the $\alpha: A \rightarrow B$ from Proposition 2.3(b) now has to be a bijection—and we just take $A = B$ as a set, with R_A the smallest preorder under which $\{b, b'\}$ has the preorder induced by R_B . \square

Proposition 5.2. *For a morphism $p: E \rightarrow B$ in *Preord*, the following conditions are equivalent:*

- (a) *p is an effective \mathbb{E} -descent morphism;*
- (b) *p satisfies the equivalent conditions of Proposition 5.1, and for every pullback (2.2) in *Rel* (or in *ReflRel*) with $\alpha \in \mathbb{E}$ and $E \times_B A$ a preorder, A also is a preorder;*
- (c) *p is surjective, and for every $b_2 \rightarrow b_1 \rightarrow b_0$ in B with $b_2 \neq b_0$, there exists $e_2 \rightarrow e_1 \rightarrow e_0$ in E with $p(e_i) = b_i$, for $i = 0, 1, 2$.*

Proof. (a) \Leftrightarrow (b) can be easily proved similarly to (a) \Leftrightarrow (b) of Proposition 3.4, with suitable generalizations of Propositions 3.1 and 3.2.

We can also repeat the proof of (c) \Rightarrow (b) from Proposition 3.4, since we do not need to consider the case $\alpha(a_2) = \alpha(a_0)$: if $\alpha(a_2) = \alpha(a_0)$ then $a_2 = a_0$, and then $a_2 \rightarrow a_0$ since R_A , being the image of $R_{E \times_B A}$, is reflexive.

And finally, in order to use the proof of Proposition 3.4(b) \Rightarrow (c) we just modify it as we did for Proposition 2.3(b) in order to prove Proposition 5.1(b) \Rightarrow (a). That is, we take $A = B$ as a set (so now $\alpha = 1_B$ is a bijection) with

$$R_A = \Delta_B \cup \{(b_2, b_1), (b_1, b_0)\}, \tag{5.1}$$

excluding the trivial cases $b_2 = b_1$ and $b_1 = b_0$; since $b_2 \neq b_0$, the set $\{b_2, b_1, b_0\}$ has exactly three elements as needed in the proof of Proposition 3.4(b) \Rightarrow (c). \square

Note that the same results are true in *FinPreord* or if \mathbb{E} is the class of all injections.

6. Étale descent

As follows from Proposition 1.5, a continuous map $\alpha: A \rightarrow B$ of finite topological spaces is étale if and only if it is a discrete fibration of the corresponding preorders

(considered as categories). Accordingly, in order to investigate the étale descent, we will take \mathbb{E} to be the class of discrete fibrations of preorders.

On the other hand, the discrete fibrations $A \rightarrow B$ correspond to the functors $B^{op} \rightarrow Sets$, and, moreover, the standard equivalence

$$\mathbb{E}(B) \sim Sets^{B^{op}} \tag{6.1}$$

is in fact an equivalence of *Preord*-indexed categories.

Using the equivalence (6.1) and the 2-equalizer (4.1) we can describe $Des_{\mathbb{E}}(p)$ (for a given $p: E \rightarrow B$ in *Preord*) as the 2-equalizer

$$Des_{\mathbb{E}}(p) \longrightarrow Sets^{E^{op}} \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} Sets^{(E \cdot_B E)^{op}} \begin{matrix} \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{matrix} Sets^{(E \cdot_B E \cdot_B E)^{op}} \tag{6.2}$$

and then a straightforward calculation gives

Proposition 6.1. *Let \mathcal{X} be the category of pairs (X, ζ) , where $X: E^{op} \rightarrow Sets$ is a functor, and*

$$\zeta = (\zeta_{e,e'})_{(e,e') \in E \times_B E}$$

a family of maps $\zeta_{e,e'}: X(e') \rightarrow X(e)$ such that

$$\zeta_{e,e'} \zeta_{e',e''} = \zeta_{e,e''}, \quad \zeta_{e,e} = 1_{X(e)} \tag{6.3}$$

and, for every $(e_1, e'_1) \rightarrow (e_0, e'_0)$ in $E \times_B E$, the diagram

$$\begin{array}{ccc} X(e'_0) & \xrightarrow{X(e'_0, e'_1)} & X(e'_1) \\ \zeta_{e_0, e'_0} \downarrow & & \downarrow \zeta_{e_1, e'_1} \\ X(e_0) & \xrightarrow{X(e_0, e'_0)} & X(e_1) \end{array} \tag{6.4}$$

commutes.

Let $\phi: Sets^{B^{op}} \rightarrow \mathcal{X}$ be the functor defined by $A \mapsto (p^{op}A, 1)$, where 1 is the family of identity morphisms

$$1_{e,e'} = 1_{A(p(e))}: A(p(e')) \rightarrow A(p(e)). \tag{6.5}$$

Then there exists a category equivalence $Des_{\mathbb{E}}(p) \sim \mathcal{X}$ making the diagram

$$\begin{array}{ccc} \mathbb{E}(B) & \xrightarrow{K_{\mathbb{E}}^p} & Des_{\mathbb{E}}(p) \\ \sim \downarrow & & \downarrow \sim \\ Sets^{B^{op}} & \xrightarrow{\phi} & \mathcal{X} \end{array} \tag{6.6}$$

commute, up to isomorphism.

Corollary 6.2. *A morphism $p: E \rightarrow B$ in *Preord* is an effective \mathbb{E} -descent morphism if and only if the functor ϕ described in Proposition 6.1 is a category equivalence.*

We point out that Corollary 6.2 is used to obtain an elementary characterization of effective étale-descent morphisms of finite topological spaces in [6], which itself should suggest such a characterization for all spaces.

Note also that the category \mathcal{X} of Proposition 6.1 can be described as the category of “double functors” $Eq(p) \rightarrow Sets_2$, where $Eq(p)$ is the equivalence relation

$$E \times_B E \rightrightarrows E \tag{6.7}$$

(= kernel pair of p) considered as a double category, and $Sets_2$ the double category of $Sets$, maps and commutative squares. Accordingly, there is a natural description of the functor $\phi : Sets^{B^{op}} \rightarrow \mathcal{X}$.

7. Triquotient maps

A continuous map $p : E \rightarrow B$ of topological spaces is said to be a triquotient map if there exists a map $q : Open(E) \rightarrow Open(B)$ of the sets of open subsets in E and in B , respectively, satisfying the following conditions:

- $q(U) \subseteq p(U)$, for every $U \in Open(E)$;
- $q(E) = B$;
- q is monotone, i.e. $U \subseteq V \Rightarrow q(U) \subseteq q(V)$;
- for every $U \in Open(E)$, $b \in q(U)$, and covering family $(E_i)_{i \in I}$ of $p^{-1}(b) \cap U$, there exists a finite set $\{i_1, \dots, i_n\} \subseteq I$ with

$$b \in q(E_{i_1} \cup \dots \cup E_{i_n}). \tag{7.1}$$

The q above is called a triquotient-assignment for p .

In the finite case, just like for the Day–Kelly maps, we could take I to be a one element set. That is, in the finite case, the last condition above is equivalent to

- If U and V are open subsets in E and b is an element in B , then

$$b \in q(U), \quad p^{-1}(b) \cap U \subseteq V \Rightarrow b \in q(V). \tag{7.2}$$

A surprising result is (compare with (c) \Leftrightarrow (b) in Proposition 2.5!):

Proposition 7.1. *If $p : E \rightarrow B$ is a triquotient map of topological spaces, then for every natural number n and every $b_n \rightarrow b_{n-1} \rightarrow \dots \rightarrow b_1 \rightarrow b_0$ in B there exists $e_n \rightarrow e_{n-1} \rightarrow \dots \rightarrow e_1 \rightarrow e_0$ in E with $p(e_i) = b_i$, for each $i = 0, \dots, n$.*

Proof. For a fixed $b_n \rightarrow b_{n-1} \rightarrow \dots \rightarrow b_1 \rightarrow b_0$ in B we introduce, for $i = 0, \dots, n$, the sets E_i defined by

$$E_i = \{e \in E \mid \text{there exists } e = e_i \rightarrow e_{i-1} \rightarrow \dots \rightarrow e_0 \text{ in } E \text{ with } p(e_{i-1}) = b_{i-1}, \dots, p(e_0) = b_0\} \tag{7.3}$$

and we are going to prove that each E_i is open and

$$b_i \in q(E_i) \tag{7.4}$$

for each i . This will give $b_n \in p(E_n)$, and therefore there exists $e_n \rightarrow \dots \rightarrow e_1 \rightarrow e_0$ with the required property.

The fact that each E_i is open follows from the obvious equalities

$$E_0 = E, \quad E_i = \downarrow (p^{-1}(b_{i-1}) \cap E_{i-1}) \quad (i > 0). \tag{7.5}$$

In order to prove (7.4) we use the induction by $i = 0, \dots, n$.

For $i = 0$ we have $b_0 \in B = q(E) = q(E_0)$.

Suppose $b_{i-1} \in q(E_{i-1})$. Since $b_i \in \downarrow b_{i-1}$ and $q(E_i)$ is open, in order to prove that $b_i \in q(E_i)$ it suffices to prove that $b_{i-1} \in q(E_i)$. However this follows from Condition 7.1 applied to $U = E_{i-1}$, $V = E_i$, and $b = b_{i-1}$, since $p^{-1}(b_{i-1}) \cap E_{i-1} \subseteq E_i$ by (7.5). \square


Now it is easy to construct effective descent morphisms of finite topological spaces which are not triquotient maps. Thus, the fact that the class of triquotient maps in $\mathcal{T}op$ is a proper subclass of the one of effective descent morphisms already appears for the finite spaces.

8. Counter-examples

So far we have never mentioned the (non-effective) \mathbb{E} -descent morphisms. They are those which have the comparison functor of (4.9) full and faithful. If \mathbb{E} (of (4.9)) is the class of all morphisms in the ground category \mathcal{C} , and \mathcal{C} has (pullbacks and) coequalizers of equivalence relations, then they are the same as the pullback stable regular epimorphisms. In particular, the descent morphisms in $\mathcal{T}op$ are the same as the Day–Kelly maps—which brings the following:

Problem 8.1. Is every Day–Kelly map an effective descent morphism in $\mathcal{T}op$?

The first counter-example was described in [14]; it uses ultrafilters, and the proof uses pseudotopological spaces. However, as shown in [17], there is even a finite counter-example; it can be displayed as




where B has the codiscrete topology, the non-trivial open sets in E are $\{e_{11}, e_{21}\}$, $\{e_{22}, e_{31}\}$ and their union, and p is defined by $p(e_{ij}) = b_i$.

The preorder approach of the present paper makes the whole story trivial: the Day–Kelly maps which are not effective descent morphisms are those maps $p : E \rightarrow B$ which

satisfy Proposition 2.5(b) but not Proposition 3.4(c). Briefly, they are those which are surjective on arrows, but not on composable pairs of arrows.


The preorder translation of (8.1) is



where the identity arrows are omitted. It is easy to see here that p is surjective on arrows but there is no $e'' \rightarrow e' \rightarrow e$ in E whose image in B is $b_3 \rightarrow b_1 \rightarrow b_2$ and so p is not surjective on composable pairs.


Note also that the preorder approach suggests to consider the following two (counter-) examples, the first of which is more straightforward, and the second gives the smallest possible spaces:

Example 8.2. Take



in fact this is exactly the finite version of the original counter-example from [14].

Example 8.3. Take



there is no $e'' \rightarrow e' \rightarrow e$ whose image in B is $b_1 \rightarrow b_2 \rightarrow b_1$.


- Remark 8.4.** As we see from Proposition 5.2(a) \Leftrightarrow (c) we have
- (a) $p: E \rightarrow B$ of Example 8.2 is not even an effective bijective-descent morphism (as well as the one from [16] mentioned above—see (8.2));
 - (b) $p: E \rightarrow B$ of Example 8.3 is an effective bijective-descent morphism.

Consider a further problem, which is suggested by the fact that every effective descent morphism in $\mathcal{T}op$ is an effective étale-descent morphism:

Problem 8.5. Does one of the following two classes of maps contain the other:

- Day–Kelly maps,
- effective étale-descent morphisms?

Using $p: E \rightarrow B$ of Example 8.2, consider the pullback in $RefRel$:



where again the display shows all arrows except the identities. Clearly, $E \times_B A$ is a preorder, and using Proposition 1.5(a) \Leftrightarrow (c) it is easy to see that $\pi_1: E \times_B A \rightarrow E$ is étale. Therefore, $(E \times_B A, \pi_1, \langle \pi_1, \pi_3 \rangle)$ constructed as in (4.7) belongs to $Des_{\mathbb{E}}(p)$, where \mathbb{E} is the class of étale maps in the category of topological spaces. Since p is an effective descent morphism in $RefRel$, and A is not a preorder, there is no object in $\mathbb{E}(B)$ corresponding to $(E \times_B A, \pi_1, \langle \pi_1, \pi_3 \rangle)$. That is we obtain

Proposition 8.6. *The Day–Kelly map $p: E \rightarrow B$ described in Example 8.2 is not an effective étale-descent morphism.*

Finally, let us consider

Example 8.7. Take

$$E = \left(\begin{array}{c} e_1 \\ \updownarrow \\ e_{21} \end{array} \quad \begin{array}{c} e_{22} \\ \updownarrow \\ e_3 \end{array} \right) \xrightarrow{p} \left(\begin{array}{c} b_1 \\ \updownarrow \\ b_2 \\ \updownarrow \\ b_3 \end{array} \right) = B \tag{8.6}$$

clearly this is not a Day–Kelly map, but a simple calculation using Corollary 6.2 (or directly, using the fact that B is a codiscrete space and E is a coproduct of two codiscrete spaces) shows that it is an effective étale-descent morphism.

Together with Proposition 8.6 this gives the negative answer to Problem 8.5.

9. Remarks on infinite spaces

In this section we list the questions and results of Topological descent theory, which became much more clear to us as soon as we understood their finite versions using the preorder approach.

(9.1) Our simple characterization of the effective descent morphisms of preorders, which Grothendieck and Giraud would probably consider as an obvious fact already 35 years ago (see [3]), can however be considered as a basic result whose “infinite filter generalization” is the Reiterman–Tholen complete characterization of the effective descent morphisms of topological spaces (see Theorem 0.1). Just observe that:

(a) The preorder on a finite topological space corresponds to the convergency structure on an infinite one; we will write $\mathcal{F} \rightarrow x$ when \mathcal{F} is a filter converging to a point x . In a finite space $\mathcal{F} \rightarrow x$ if and only if $y \rightarrow x$ for every y which belongs to the intersection of the elements of \mathcal{F} . Moreover, the passage from the topologies to the corresponding convergency structures determines a category isomorphism which extends the isomorphism (1.5).

(b) Since ultrafilters on a finite set are principal filters generated by the one-point subsets, the “relevant part” of a crest of ultrafilters $((\mathcal{F}_i \rightarrow b_i)_{i \in I}, \mathcal{U}, b)$ (in the sense of [14]) is the composable pair $b'' \rightarrow b' \rightarrow b$ in which $b' = b_i$ and b'' have i generating \mathcal{U} and $\{b''\}$ generating the corresponding \mathcal{F}_i .

(c) Recall that the isomorphism $FinTop \cong FinPreord$ extends to an isomorphism $FinPsTop \cong FinRefRel$, where $FinPsTop$ is the category of finite pseudotopological

spaces. And the results of [14] use the embedding $\mathcal{T}op \rightarrow PsTop$ exactly in the same way as we use $Prord \rightarrow ReflRel$.

(9.2) The three classes of morphisms in $\mathcal{T}op$ which were known to be classes of effective descent morphisms before [14], are

1. (locally) sectionable maps,
2. open surjections,
3. proper surjections.

Why? In the finite case (although “proper” reduces to “closed”) these three classes naturally occur as the three simple cases. Indeed, in order to find $e_2 \rightarrow e_1 \rightarrow e_0$ for a given $b_2 \rightarrow b_1 \rightarrow b_0$ as in Proposition 3.4(c), one could either

- use a section $B \rightarrow E$ (or a local section);
- or first find e_0 with $p(e_0) = b_0$, then e_1 using e_0 and $b_1 \rightarrow b_0$ via Proposition 1.4(d) (which is equivalent to openness), and then e_2 using $b_2 \rightarrow b_1$ via Proposition 1.4(d) again;
- or first find e_2 with $p(e_2) = b_2$, then e_1 using e_2 and $b_2 \rightarrow b_1$ via Proposition 1.3(e) (which is equivalent to closeness), and then e_0 using e_1 and $b_1 \rightarrow b_0$ via Proposition 1.3(e) again.

(9.3) See Problem 8.5; the negative answer is provided by finite counter-examples (see (8.5) and (8.6)):

(9.4) Proposition 5.2 clearly shows the difference between the (ordinary) effective descent morphisms and the effective bijective-descent morphisms: compare Propositions 5.2(c) and 3.4(c). Note also that our proof of Proposition 5.1 is, in fact, the finite version of the proof of Theorem 4.2 in [16].

References

- [1] M.M. Clementino, On finite triquotient maps, *J. Pure Appl. Algebra* 168 (2002) 387–389.
- [2] B.J. Day, G.M. Kelly, On topological quotient maps preserved by pullbacks, *Proc. Cambridge Philos. Soc.* 67 (1970) 553–558.
- [3] J. Giraud, Méthode de la descente, *Bull. Soc. Math. France Mem.* 2 (1964).
- [4] M. Gran, Central extensions for internal groupoids in Maltsev categories, Ph.D. Thesis, Louvain-la-Neuve, Belgium, 1999.
- [5] G. Janelidze, M. Sobral, Finite preorders and topological descent, Preprint 99-10, Department of Mathematics, University of Coimbra.
- [6] G. Janelidze, M. Sobral, Finite preorders and topological descent II: étale descent, *J. Pure Appl. Algebra*, to appear.
- [7] G. Janelidze, W. Tholen, How algebraic is the change-of-base functor?, *Lecture Notes in Mathematics*, Vol. 1448, Springer, Berlin, 1991, pp. 157–173.
- [8] G. Janelidze, W. Tholen, Facets of descent I, *Appl. Cat. Struct.* 2 (1994) 245–281.
- [9] G. Janelidze, W. Tholen, Facets of descent II, *Appl. Cat. Struct.* 5 (1997) 229–248.
- [10] A. Kock, I. Moerdijk, Local equivalence relations and their sheaves, *Aarhus Preprint Series*, Vol. 19, 1991.
- [11] I. Le Creurer, Descent for internal categories, Ph.D. Thesis, Louvain-la-Neuve, Belgium, 1999.
- [12] I. Moerdijk, Descent theory for toposes, *Bull. Soc. Math. Belg.* 41 (1989) 373–391.
- [13] T. Plewe, Localic triquotient maps are effective descent morphisms, *Math. Proc. Cambridge Philos. Soc.* 122 (1997) 17–43.

- [14] J. Reiterman, W. Tholen, Effective descent maps of topological spaces, *Topology Appl.* 57 (1994) 53–69.
- [15] M. Sobral, Some aspects of topological descent, *Appl. Cat. Struct.* 4 (1996) 97–106.
- [16] M. Sobral, Another approach to topological descent theory, *Appl. Cat. Struct.* 9 (2001) 505–516.
- [17] M. Sobral, W. Tholen, Effective descent morphisms and effective equivalence relations, *CMS Conference Proceedings*, Vol. 13, American Mathematical Society, Providence, RI, 1992, pp. 421–433.
- [18] J.J.C. Vermeulen, Proper maps of locales, *J. Pure Appl. Algebra* 92 (1994) 79–107.