

Short Communication

MALKHAZ ASHORDIA, GODERDZI EKHVAIA AND NESTAN KEKELIA

ON THE WELL-POSEDNESS OF GENERAL NONLINEAR BOUNDARY VALUE PROBLEMS FOR SYSTEMS OF DIFFERENTIAL EQUATIONS WITH FINITE AND FIXED POINTS OF IMPULSES

Abstract. The general nonlocal boundary value problems are considered for systems of differential equations with finite and fixed points of impulses. The sufficient conditions, among which are effective spectral ones, are established for the well-posedness of these problems.

რეზიუმე. განხილულია ზოგადი სახის არაწრფივი სასაზღვრო ამოცანები დიფერენციალურ განტოლებათა სისტემებისთვის სასრული და ფიქსირებული იმპულსების წერტილებით. დადგენილია ამ ამოცანების კორექტულობის საკმარისი, მათ შორის ეფექტური სპექტრალური პირობები.

2000 Mathematics Subject Classification: 34K10, 34K45.

Key words and phrases: Nonlocal boundary value problems, nonlinear systems, impulsive equations, solvability, unique solvability, effective conditions.

1. STATEMENT OF THE PROBLEM AND BASIC NOTATION

In the present paper, we consider the system of nonlinear impulsive equations with a finite number of impulses points

$$\frac{dx}{dt} = f(t, x) \text{ almost everywhere on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \tag{1.1}$$

$$x(\tau_l+) - x(\tau_l-) = I_l(x(\tau_l)) \quad (l = 1, \dots, m_0) \tag{1.2}$$

with the general boundary value problem

$$h(x) = 0, \tag{1.3}$$

where $a < \tau_1 < \dots < \tau_{m_0} \leq b$ (we will assume $\tau_0 = a$ and $\tau_{m_0+1} = b$, if necessary), $-\infty < a < b < +\infty$, m_0 is a natural number, $f = (f_i)_{i=1}^n$ belongs to Carathéodory class $\text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^n)$, $I_l = (I_{li})_{i=1}^n : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($l = 1, \dots, m_0$) are continuous operators, and $h : C([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ is a continuous, nonlinear in general, vector-functional.

Let x_0 be a solution of the problem (1.1), (1.2); (1.3).

Consider a sequence of vector-functions $f_k \in \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^n)$ ($k = 1, 2, \dots$), the sequences of continuous operators $I_{lk} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($k = 1, 2, \dots; l = 1, \dots, m_0$), the sequences τ_{lk} ($k = 1, 2, \dots; l = 1, \dots, m_0$) of points $a < \tau_{1k} < \dots < \tau_{m_0k} \leq b$ and the sequence of continuous vector-functionals $h_k : C([a, b], \mathbb{R}^n; \tau_{1k}, \dots, \tau_{m_0k}) \rightarrow \mathbb{R}^n$ ($k = 1, 2, \dots$).

In this paper the sufficient conditions are given guaranteeing both the solvability of the impulsive boundary value problems

$$\frac{dx}{dt} = f_k(t, x) \text{ almost everywhere on } [a, b] \setminus \{\tau_{1k}, \dots, \tau_{m_0k}\}, \quad (1.1_k)$$

$$x(\tau_{lk}+) - x(\tau_{lk}-) = I_{lk}(x(\tau_{lk})) \quad (l = 1, \dots, m_0); \quad (1.2_k)$$

$$h_k(x) = 0 \quad (1.3_k)$$

($k = 1, 2, \dots$) for any sufficient large k and the convergence of its solutions to a solution of the problem (1.1), (1.2); (1.3) as $k \rightarrow +\infty$.

As is known, the question of the well-posedness for the nonlinear impulsive boundary value problems was not investigated in earlier works. So the statement of the problems under consideration is actual.

The obtained results are analogous to ones given in [12] (see also the references therein) for the general nonlinear boundary value problems for systems of ordinary differential equations. Some results obtained in the paper are more general than already known ones even for ordinary differential case.

The analogous question is investigated in [4], [8] for linear boundary value problems for systems with impulses, and in [1]–[3], [12]–[15] for linear and nonlinear boundary value problems for systems of ordinary differential equations. Notice that the necessary and sufficient conditions are obtained for the well-posedness of the linear boundary value problem in [8] for impulsive, and in [1]–[3] for ordinary differential systems.

Quite a number of issues of the theory of systems of differential equations with impulsive effect (both linear and nonlinear) have been studied sufficiently well (for a survey of the results on impulsive systems see, e.g., [5]–[7], [9]–[11], [16], [17] and the references therein).

Throughout the paper, the following notation and definitions will be used.

$\mathbb{R} =] - \infty, +\infty[$, $\mathbb{R}_+ = [0, +\infty[$; $[a, b]$ ($a, b \in \mathbb{R}$) is a closed segment.

$\mathbb{R}^{n \times m}$ is the space of all real $n \times m$ -matrices $X = (x_{ij})_{i,j=1}^{n,m}$ with the norms

$$\|X\| = \max_{j=1, \dots, m} \sum_{i=1}^n |x_{ij}|;$$

$$|X| = (|x_{ij}|)_{i,j=1}^{n,m}, \quad [X]_+ = \frac{|X| + X}{2};$$

$$\mathbb{R}_+^{n \times m} = \left\{ (x_{ij})_{i,j=1}^{n,m} : x_{ij} \geq 0 \ (i = 1, \dots, n; j = 1, \dots, m) \right\};$$

$$\mathbb{R}^{(n \times n) \times m} = \mathbb{R}^{n \times n} \times \dots \times \mathbb{R}^{n \times n} \quad (m\text{-times}).$$

$\mathbb{R}^n = \mathbb{R}^{n \times 1}$ is the space of all real column n -vectors $x = (x_i)_{i=1}^n$; $\mathbb{R}_+^n = \mathbb{R}_+^{n \times 1}$.

If $X \in \mathbb{R}^{n \times n}$, then X^{-1} , $\det X$ and $r(X)$ are, respectively, the matrix inverse to X , the determinant of X and the spectral radius of X ; $I_{n \times n}$ is the identity $n \times n$ -matrix.

$\overset{b}{\underset{a}{V}}(X)$ is the total variation of the matrix-function $X : [a, b] \rightarrow \mathbb{R}^{n \times m}$, i.e., the sum of total variations of the latter's components; $V(X)(t) = (v(x_{ij})(t))_{i,j=1}^{n,m}$, where $v(x_{ij})(a) = 0$, $v(x_{ij})(t) = \overset{t}{\underset{a}{V}}(x_{ij})$ for $a < t \leq b$;

$X(t-)$ and $X(t+)$ are the left and the right limit of the matrix-function $X : [a, b] \rightarrow \mathbb{R}^{n \times m}$ at the point t (we will assume $X(t) = X(a)$ for $t \leq a$ and $X(t) = X(b)$ for $t \geq b$, if necessary);

$$\|X\|_s = \sup \{ \|X(t)\| : t \in [a, b] \}.$$

$BV([a, b], \mathbb{R}^{n \times m})$ is the set of all matrix-functions of bounded variation $X : [a, b] \rightarrow \mathbb{R}^{n \times m}$ (i.e., such that $\overset{b}{\underset{a}{V}}(X) < +\infty$);

$C([a, b], D)$, where $D \subset \mathbb{R}^{n \times m}$, is the set of all continuous matrix-functions $X : [a, b] \rightarrow D$;

$C([a, b], D; \tau_1, \dots, \tau_{m_0})$ is the set of all matrix-functions $X : [a, b] \rightarrow D$, having the one-sided limits $X(\tau_l-)$ ($l = 1, \dots, m_0$) and $X(\tau_l+)$ ($l = 1, \dots, m_0$), whose restrictions to an arbitrary closed interval $[c, d]$ from $[a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}$ belong to $C([c, d], D)$;

$C_s([a, b], \mathbb{R}^{n \times m}; \tau_1, \dots, \tau_{m_0})$ is the Banach space of all $X \in C([a, b], \mathbb{R}^{n \times m}; \tau_1, \dots, \tau_{m_0})$ with the norm $\|X\|_s$.

If $y \in C_s([a, b], \mathbb{R}; \tau_1, \dots, \tau_{m_0})$ and $r \in]0, +\infty[$, then

$$U(y; r) = \left\{ x \in C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) : \|x - y\|_s < r \right\};$$

$D(y, r)$ is the set of all $x \in \mathbb{R}^n$ such that $\inf \{ \|x - y(t)\| : t \in [a, b] \} < r$.

$\tilde{C}([a, b], D)$, where $D \subset \mathbb{R}^{n \times m}$, is the set of all absolutely continuous matrix-functions $X : [a, b] \rightarrow D$;

$\tilde{C}([a, b], D; \tau_1, \dots, \tau_{m_0})$ is the set of all matrix-functions $X : [a, b] \rightarrow D$, having the one-sided limits $X(\tau_l-)$ ($l = 1, \dots, m_0$) and $X(\tau_l+)$ ($l = 1, \dots, m_0$), whose restrictions to an arbitrary closed interval $[c, d]$ from $[a, b] \setminus \{\tau_k\}_{k=1}^{m_0}$ belong to $\tilde{C}([c, d], D)$.

If B_1 and B_2 are the normed spaces, then an operator $g : B_1 \rightarrow B_2$ (nonlinear, in general) is positive homogeneous if $g(\lambda x) = \lambda g(x)$ for every $\lambda \in \mathbb{R}_+$ and $x \in B_1$.

The inequalities between the matrices are understood component wise.

An operator $\varphi : C([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ is called nondecreasing if for every $x, y \in C([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0})$ such that $x(t) \leq y(t)$ for $t \in [a, b]$ the inequality $\varphi(x)(t) \leq \varphi(y)(t)$ holds for $t \in [a, b]$.

A matrix-function is said to be continuous, nondecreasing, integrable, etc., if each of its components is such.

$L([a, b], D)$, where $D \subset \mathbb{R}^{n \times m}$, is the set of all measurable and integrable matrix-functions $X : [a, b] \rightarrow D$.

If $D_1 \subset \mathbb{R}^n$ and $D_2 \subset \mathbb{R}^{n \times m}$, then $\text{Car}([a, b] \times D_1, D_2)$ is the Carathéodory class, i.e., the set of all mappings $F = (f_{kj})_{k,j=1}^{n,m} : [a, b] \times D_1 \rightarrow D_2$ such that for each $i \in \{1, \dots, l\}$, $j \in \{1, \dots, m\}$ and $k \in \{1, \dots, n\}$:

- (a) the function $f_{kj}(\cdot, x) : [a, b] \rightarrow D_2$ is measurable for every $x \in D_1$;
- (b) the function $f_{kj}(t, \cdot) : D_1 \rightarrow D_2$ is continuous for almost every $t \in [a, b]$, and

$\sup \{|f_{kj}(\cdot, x)| : x \in D_0\} \in L([a, b], \mathbb{R}; g_{ik})$ for every compact $D_0 \subset D_1$.

$\text{Car}^0([a, b] \times D_1, D_2)$ is the set of all mappings $F = (f_{kj})_{k,j=1}^{n,m} : [a, b] \times D_1 \rightarrow D_2$ such that the functions $f_{kj}(\cdot, x(\cdot))$ ($j = 1, \dots, m$; $k = 1, \dots, n$) are measurable for every vector-function $x : [a, b] \rightarrow \mathbb{R}^n$ with the bounded total variation.

$M([a, b] \times \mathbb{R}_+, \mathbb{R}_+)$ is the set of all functions $\omega \in \text{Car}([a, b] \times \mathbb{R}_+, \mathbb{R}_+)$ such that the function $\omega(t, \cdot)$ is nondecreasing, and $\omega(t, 0) \equiv 0$.

By a solution of the impulsive system (1.1), (1.2) we understand a continuous from the left vector-function $x \in \widetilde{C}([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0})$ satisfying both the system (1.1) for a.e. on $[a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}$ and the relation (1.2) for every $l \in \{1, \dots, m_0\}$.

Definition 1.1. Let $l : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ be a linear continuous operator, and let $l_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ be a positive homogeneous operator. We say that a pair $(P, \{J_l\}_{l=1}^{m_0})$, consisting of a matrix-function $P \in \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^{n \times n})$ and a finite sequence of continuous operators $J_l = (J_{li})_{i=1}^n : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($l = 1, \dots, m_0$), satisfy the Opial condition with respect to the pair (l, l_0) if:

- (a) there exist a matrix-function $\Phi \in L([a, b], \mathbb{R}_+^n)$ and constant matrices $\Psi_l \in \mathbb{R}^{n \times n}$ ($l = 1, \dots, m_0$) such that

$$|P(t, x)| \leq \Phi(t) \text{ a.e. on } [a, b], \quad x \in \mathbb{R}^n$$

and

$$|J_l(x)| \leq \Psi_l \text{ for } x \in \mathbb{R}^n \quad (l = 1, \dots, m_0);$$

- (b)

$$\det(I_{n \times n} + G_l) \neq 0 \quad (l = 1, \dots, m_0) \quad (1.4)$$

and the problem

$$\frac{dx}{dt} = A(t)x \text{ a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad (1.5)$$

$$x(\tau_l+) - x(\tau_l-) = G_l x(\tau_l) \quad (l = 1, \dots, m_0); \quad (1.6)$$

$$|l(x)| \leq l_0(x) \quad (1.7)$$

has only the trivial solution for every matrix-function $A \in L([a, b], \mathbb{R}^{n \times n})$ and constant matrices G_l ($l = 1, \dots, m_0$) for which there exists a sequence $y_k \in \tilde{C}([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0})$ ($k = 1, 2, \dots$) such that

$$\lim_{k \rightarrow +\infty} \int_a^t P(\tau, y_k(\tau)) d\tau = \int_a^t A(\tau) d\tau \quad \text{uniformly on } [a, b]$$

and

$$\lim_{k \rightarrow +\infty} J_l(y_k(\tau_l)) = G_l \quad (l = 1, \dots, m_0).$$

Remark 1.1. In particular, the condition (1.4) holds if

$$\|\Psi_l\| < 1 \quad (l = 1, \dots, m_0).$$

Below we will assume that $f = (f_i)_{i=1}^n \in \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^n)$ and, in addition, $f(\tau_l, x)$ is arbitrary for $x \in \mathbb{R}^n$ ($l = 1, \dots, m_0$).

Let x^0 be a solution of the problem (1.1), (1.2); (1.3), and r be a positive number. Let us introduce the following definition.

Definition 1.2. The solution x^0 is said to be strongly isolated in the radius r if there exist, respectively, the matrix- and the vector-functions $P \in \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^{n \times n})$ and $q \in \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^n)$, a finite sequence of continuous operators $J_l : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($l = 1, \dots, m_0$) and $h_l : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($l = 1, \dots, m_0$), a linear continuous operator $l : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$, a positive homogeneous operator $l_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$, and a continuous operator $\tilde{l} : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ such that

(a) the equalities

$$\begin{aligned} f(t, x) &= P(t, x)x + q(t, x) \quad \text{for } t \in [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad \|x - x^0(t)\| < r, \\ I_l(x) &= J_l(x)x + h_l(x) \quad \text{for } \|x - x^0(\tau_l)\| < r \quad (l = 1, \dots, m_0); \end{aligned}$$

and

$$h(x) = l(x) + \tilde{l}(x) \quad \text{for } x \in U(x^0; r)$$

are valid;

(b) the functions

$$\begin{aligned} \alpha(t, \rho) &= \max \{ \|q(t, x)\| : \|x\| \leq \rho \}, \\ \beta_l(\rho) &= \max \{ \|h_l(x)\| : \|x\| \leq \rho \} \quad (l = 1, \dots, m_0) \end{aligned}$$

and

$$\gamma(\rho) = \sup \left\{ [|\tilde{l}(x)| - l_0(x)]_+ : \|x\|_s \leq \rho \right\}$$

satisfy the condition

$$\limsup_{\rho \rightarrow +\infty} \frac{1}{\rho} \left(\gamma(\rho) + \int_a^b \alpha(t, \rho) dt + \sum_{l=1}^{m_0} \beta_l(\rho) \right) < 1;$$

(c) the problem

$$\begin{aligned} \frac{dx}{dt} &= P(t, x)x + q(t, x) \text{ a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \\ x(\tau_l+) - x(\tau_l-) &= J_l(x(\tau_l)) + h_l(x(\tau_l)) \quad (l = 1, \dots, m_0); \\ l(x) + \tilde{l}(x) &= 0 \end{aligned}$$

has no solution different from x^0 .

(d) the pair $(P, \{J_l\}_{l=1}^{m_0})$ satisfy the Opial condition with respect to the pair (l, l_0) .

Definition 1.3. We say that a sequence $(f_k, \{I_{lk}\}_{l=1}^{m_0}; h_k)$ ($k = 1, 2, \dots$) belongs to the set $W_r(f, \{I_l\}_{l=1}^{m_0}, h; x^0)$ if

(a) the equalities

$$\lim_{k \rightarrow +\infty} \int_a^t f_k(\tau, x) d\tau = \int_a^t f(\tau, x) d\tau \text{ uniformly on } [a, b]$$

and

$$\lim_{k \rightarrow +\infty} I_{lk}(x) = I_l(x) \quad (l = 1, \dots, m_0)$$

are valid for every $x \in D(x^0; r)$;

(b)

$$\lim_{k \rightarrow +\infty} h_k(x) = h(x) \text{ uniformly on } U(x^0; r);$$

(c) there exist a sequence of functions $\omega_k \in M([a, b] \times \mathbb{R}_+, \mathbb{R}_+)$ ($k = 1, 2, \dots$) and sequences of nondecreasing functions $\omega_{lk} \in C(\mathbb{R}_+, \mathbb{R}_+)$, $\omega_{lk}(0) = 0$, ($k = 1, 2, \dots; l = 1, \dots, m_0$) such that

$$\sup \left\{ \int_a^b \omega_k(t, r) dt : k = 1, 2, \dots \right\} < +\infty, \quad (1.8)$$

$$\sup \left\{ \sum_{l=1}^{m_0} \omega_{lk}(r) : k = 1, 2, \dots \right\} < +\infty; \quad (1.9)$$

$$\lim_{s \rightarrow 0+} \sup \left\{ \int_a^b \omega_k(t, s) dt : k = 1, 2, \dots \right\} = 0, \quad (1.10)$$

$$\lim_{s \rightarrow 0^+} \sup \left\{ \sum_{l=1}^{m_0} \omega_{lk}(s) : k = 1, 2, \dots \right\} = 0; \quad (1.11)$$

and

$$\begin{aligned} & \|f_k(t, x) - f_k(t, y)\| \leq \omega_k(t, \|x - y\|) \\ \text{for } t \in [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad x, y \in D(x^0; r) \quad (k = 1, 2, \dots), \\ & \|I_{lk}(x) - I_{lk}(y)\| \leq \omega_{lk}(\|x - y\|) \\ \text{for } x, y \in D(x^0; r) \quad (l = 1, \dots, m_0; \quad k = 1, 2, \dots). \end{aligned}$$

Remark 1.2. If for every natural m there exists a positive number ν_m such that

$$\begin{aligned} & \omega_k(t, m\delta) \leq \nu_m \omega_k(t, \delta) \\ \text{for } \delta > 0, \quad t \in [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\} \quad (k = 1, 2, \dots), \end{aligned}$$

then the estimate (1.8) follows from the condition (1.10); analogously, if

$$\omega_{lk}(m\delta) \leq \nu_m \omega_{lk}(\delta) \quad \text{for } \delta > 0 \quad (l = 1, \dots, m_0; \quad k = 1, 2, \dots),$$

then the estimate (1.9) follows from the condition (1.11). In particular, the sequences of the functions

$$\begin{aligned} \omega_k(t, \delta) = \max \left\{ \|f_k(t, x) - f_k(t, y)\| : x, y \in U(0, \|x^0\| + r), \|x - y\| \leq \delta \right\} \\ \text{for } t \in [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\} \quad (k = 1, 2, \dots) \end{aligned}$$

and

$$\begin{aligned} \omega_{lk}(\delta) = \max \left\{ \|I_{lk}(x) - I_{lk}(y)\| : x, y \in U(0, \|x^0\| + r), \|x - y\| \leq \delta \right\} \\ (l = 1, \dots, m_0; \quad k = 1, 2, \dots) \end{aligned}$$

have, respectively, the latters' properties.

Definition 1.4. The problem (1.1), (1.2); (1.3) is said to be $(x^0; r)$ -correct if for every $\varepsilon \in]0, r[$ and $((f_k, \{I_{lk}\}_{l=1}^{m_0}; h_k))_{k=1}^{+\infty} \in W_r(f, \{I_l\}_{l=1}^{m_0}, h; x^0)$ there exists a natural number k_0 such that the problem $(1, 1_k)$, (1.2_k) ; (1.3_k) has at least a solution contained in $U(x^0; r)$ and any such solution belongs to the ball $U(x^0; \varepsilon)$ for every $k \geq k_0$.

Definition 1.5. The problem (1.1), (1.2); (1.3) is said to be correct if it has the unique solution x^0 and it is $(x^0; r)$ -correct for every $r > 0$.

Theorem 1.1. *If the problem (1.1), (1.2); (1.3) has a solution x^0 , strongly isolated in the radius $r > 0$, then it is $(x^0; r)$ -correct.*

Theorem 1.2. *Let the conditions*

$$\begin{aligned} & \|f(t, x) - P(t, x)x\| \leq \alpha(t, \|x\|) \\ \text{a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad x \in \mathbb{R}^n, \end{aligned} \quad (1.12)$$

$$\|I_l(x) - J_l(x)x\| \leq \beta_l(\|x\|) \quad \text{for } x \in \mathbb{R}^n \quad (l = 1, \dots, m_0) \quad (1.13)$$

and

$$|h(x) - l(x)| \leq l_0(x) + l_1(\|x\|_s) \text{ for } x \in \text{BV}([a, b], \mathbb{R}^n) \quad (1.14)$$

hold, where $l : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ and $l_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ are, respectively, linear continuous and positive homogeneous continuous operators, the pair $(P, \{J_l\}_{l=1}^{m_0})$ satisfies the Opial condition with respect to the pair (l, l_0) ; $\alpha \in \text{Car}([a, b] \times \mathbb{R}_+, \mathbb{R}_+)$ is a function nondecreasing in the second variable, and $\beta_l \in C([a, b], \mathbb{R}_+)$ ($l = 1, \dots, m_0$) and $\ell_1 \in C(\mathbb{R}, \mathbb{R}_+^n)$ are, respectively, nondecreasing functions and vector-function such that

$$\limsup_{\rho \rightarrow +\infty} \frac{1}{\rho} \left(\|l_1(\rho)\| + \int_a^b \alpha(t, \rho) dt + \sum_{l=1}^{m_0} \beta_l(\rho) \right) < 1. \quad (1.15)$$

Then one-valued solvability of the problem (1.1), (1.2); (1.3) guarantees its correctness.

Theorem 1.3. Let the conditions (1.12)–(1.14),

$$P_1(t) \leq P(t, x) \leq P_2(t) \text{ a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad x \in \mathbb{R}^n,$$

and

$$J_{1l} \leq I_k(x) \leq J_{2l} \text{ for } x \in \mathbb{R}^n \quad (l = 1, \dots, m_0)$$

hold, where $P \in \text{Car}^0([a, b] \times \mathbb{R}^n, \mathbb{R}^{n \times n})$, $P_i \in L([a, b], \mathbb{R}^{n \times n})$ ($i = 1, 2$), $J_{il} \in \mathbb{R}^{n \times n}$ ($i = 1, 2; l = 1, \dots, m_0$), $l : C_s([a, b], \mathbb{R}^{n \times n}; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ and $l_0 : C_s([a, b], \mathbb{R}^{n \times n}; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ are, respectively, the linear continuous and the positive homogeneous continuous operators; $\alpha \in \text{Car}([a, b] \times \mathbb{R}_+, \mathbb{R}_+)$ is a function nondecreasing in the second variable, and $\beta_l \in C([a, b], \mathbb{R}_+)$ ($l = 1, \dots, m_0$) and $\ell_1 \in C(\mathbb{R}, \mathbb{R}_+^n)$ are, respectively, nondecreasing functions and vector-function such that the condition (1.15) holds. Let, moreover, the condition (1.4) hold and the problem (1.5), (1.6); (1.7) have only the trivial solution for every matrix-function $A \in L([a, b], \mathbb{R}^{n \times n})$ and the constant matrices $G_l \in \mathbb{R}^{n \times n}$ ($l = 1, \dots, m_0$) such that

$$P_1(t) \leq A(t) \leq P_2(t) \text{ a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad x \in \mathbb{R}^n$$

and

$$J_{1l} \leq G_l \leq J_{2l} \text{ for } x \in \mathbb{R}^n \quad (l = 1, \dots, m_0).$$

Then one-valued solvability of the problem (1.1), (1.2); (1.3) guarantees its correctness.

Remark 1.3. Theorem 1.3 is interesting only in the case where $P \notin \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^{n \times n})$, because the theorem immediately follows from Theorem 1.2 in the case where $P \in \text{Car}([a, b] \times \mathbb{R}^n, \mathbb{R}^{n \times n})$.

Theorem 1.4. *Let the conditions (1.14),*

$$|f(t, x) - P_0(t)x| \leq Q(t)|x| + q(t, \|x\|) \quad \text{a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \\ x \in \mathbb{R}^n,$$

and

$$|I_l(x) - J_{0l}x| \leq H_l|x| + h_l(\|x\|) \quad \text{for } x \in \mathbb{R}^n \quad (l = 1, \dots, m_0)$$

hold, where $P_0 \in L([a, b], \mathbb{R}^{n \times n})$, $Q \in L([a, b], \mathbb{R}_+^{n \times n})$, J_{0l} and $H_l \in \mathbb{R}^{n \times n}$ ($l = 1, \dots, m_0$) are constant matrices, $l : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ and $l_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ are, respectively, the linear continuous and positive homogeneous continuous operators; $q \in \text{Car}([a, b] \times \mathbb{R}_+, \mathbb{R}_+^n)$ is a vector-function nondecreasing in the second variable, and $h_l \in C([a, b], \mathbb{R}_+)$ ($l = 1, \dots, m_0$) and $l_1 \in C(\mathbb{R}, \mathbb{R}_+^n)$ are, respectively, the nondecreasing functions and vector-function such that the condition (1.15) holds. Let, moreover, the conditions

$$\det(I_{n \times n} + J_{0l}) \neq 0 \quad (l = 1, \dots, m_0) \quad (1.16)$$

and

$$\|H_l\| \cdot \|(I_{n \times n} + J_{0l})^{-1}\| < 1 \quad (l = 1, \dots, m_0) \quad (1.17)$$

hold, and the system of impulsive inequalities

$$\left| \frac{dx}{dt} - P_0(t)x \right| \leq Q(t)x \quad \text{a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad (1.18)$$

$$|x(\tau_l+) - x(\tau_l-) - J_{0l}x(\tau_l)| \leq H_l x(\tau_l) \quad (l = 1, \dots, m_0) \quad (1.19)$$

have only the trivial solution under the condition (1.7). Then one-valued solvability of the problem (1.1), (1.2); (1.3) guarantees its correctness.

Corollary 1.1. *Let the conditions (1.16)*

$$|f(t, x) - P_0(t)x| \leq q(t, \|x\|) \quad \text{a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad x \in \mathbb{R}^n, \quad (1.20)$$

and

$$|I_l(x) - J_{0l}x| \leq h_l(\|x\|) \quad \text{for } x \in \mathbb{R}^n \quad (l = 1, \dots, m_0) \quad (1.21)$$

hold, where $P_0 \in L([a, b], \mathbb{R}^{n \times n})$, $J_{0l} \in \mathbb{R}^{n \times n}$ ($l = 1, \dots, m_0$) are the constant matrices, $l : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ and $l_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ are, respectively, the linear continuous and positive homogeneous continuous operators; $q \in \text{Car}([a, b] \times \mathbb{R}_+, \mathbb{R}_+^n)$ is a vector-function nondecreasing in the second variable, and $h_l \in C([a, b], \mathbb{R}_+)$ ($l = 1, \dots, m_0$) and $l_1 \in C(\mathbb{R}, \mathbb{R}_+^n)$ are, respectively, the nondecreasing functions and a vector-function such that the condition (1.15) holds. Let, moreover,

$$|h(x) - \ell(x)| \leq l_1(\|x\|_s) \quad \text{for } x \in \text{BV}([a, b], \mathbb{R}^n) \quad (1.22)$$

and the impulsive system

$$\begin{aligned} \frac{dx}{dt} &= P_0(t)x \text{ a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \\ x(\tau_l+) - x(\tau_l-) &= J_{0l}x(\tau_l) \quad (l = 1, \dots, m_0) \end{aligned}$$

have only the trivial solution under the condition (1.7). Then one-valued solvability of the problem (1.1), (1.2); (1.3) guarantees its correctness.

For every matrix-function $X \in L([a, b], \mathbb{R}^{n \times n})$ and a sequence of constant matrices $Y_k \in \mathbb{R}^{n \times n}$ ($k = 1, \dots, m_0$) we introduce the operators

$$\begin{aligned} [(X, Y_1, \dots, Y_{m_0})(t)]_0 &= I_n \text{ for } a \leq t \leq b, \\ [(X, Y_1, \dots, Y_{m_0})(a)]_i &= O_{n \times n} \quad (i = 1, 2, \dots), \\ [(X, Y_1, \dots, Y_{m_0})(t)]_{i+1} &= \int_a^t X(\tau) \cdot [(X, Y_1, \dots, Y_{m_0})(\tau)]_i d\tau + \\ + \sum_{a \leq \tau_l < t} Y_l \cdot [(X, Y_1, \dots, Y_{m_0})(\tau_l)]_i &\text{ for } a < t \leq b \quad (i = 1, 2, \dots). \end{aligned} \quad (1.23)$$

Corollary 1.2. *Let the conditions (1.16), (1.20)–(1.22) hold, where*

$$\ell(x) \equiv \int_a^b dK(t) \cdot x(t),$$

$P_0 \in L([a, b], \mathbb{R}^{n \times n})$, $J_{0l} \in \mathbb{R}^{n \times n}$ ($l = 1, \dots, m_0$) are constant matrices, $K \in L([a, b], \mathbb{R}^{n \times n})$, $\ell_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ are the positive homogeneous continuous operators; $q \in \text{Car}([a, b] \times \mathbb{R}_+, \mathbb{R}_+^n)$ is a vector-function nondecreasing in the second variable, and $h_l \in C([a, b], \mathbb{R}_+)$ ($l = 1, \dots, m_0$) and $\ell_1 \in C(\mathbb{R}, \mathbb{R}_+^n)$ are, respectively, the nondecreasing functions and a vector-function such that the condition (1.15) holds. Let, moreover, there exist natural numbers k and m such that the matrix

$$M_k = - \sum_{i=0}^{k-1} \int_a^b dK(t) \cdot [(P_0, G_1, \dots, G_{m_0})(t)]_i$$

is nonsingular and

$$r(M_{k,m}) < 1,$$

where the operators $[(P_0, G_1, \dots, G_{m_0})(t)]_i$ ($i = 0, 1, \dots$) are defined by (1.23), and

$$\begin{aligned} M_{k,m} = & \left[(|P_0|, |G_1|, \dots, |G_{m_0}|)(b) \right]_m + \\ & + \sum_{i=0}^{m-1} \left[(|P_0|, |G_1|, \dots, |G_{m_0}|)(b) \right]_i \times \\ & \times \int_a^b dV(M_k^{-1}K)(t) \cdot \left[(|P_0|, |G_1|, \dots, |G_{m_0}|)(t) \right]_k. \end{aligned}$$

Then one-valued solvability of the problem (1.1), (1.2); (1.3) guarantees its correctness.

Theorem 1.5. Let the conditions (1.16), (1.17),

$$\begin{aligned} |f(t, x) - f(t, y) - P_0(t)(x - y)| & \leq Q(t)|x - y| \\ & \text{a.e. on } [a, b] \setminus \{\tau_1, \dots, \tau_{m_0}\}, \quad x, y \in \mathbb{R}^n, \\ |I_l(x) - I_l(y) - J_{0l} \cdot (x - y)| & \leq H_k \cdot |x - y| \\ & \text{for } x, y \in \mathbb{R}^n \quad (k = l, \dots, m_0) \end{aligned}$$

and

$$|h(x) - h(y) - \ell(x - y)| \leq \ell_0(x - y) \quad \text{for } x, y \in \text{BV}([a, b], \mathbb{R}^n)$$

hold, where $P_0 \in L([a, b], \mathbb{R}^{n \times n})$, $Q \in L([a, b], \mathbb{R}_+^{n \times n})$, J_{0k} and $H_l \in \mathbb{R}^{n \times n}$ ($l = 1, \dots, m_0$) are the constant matrices, $\ell : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}^n$ and $\ell_0 : C_s([a, b], \mathbb{R}^n; \tau_1, \dots, \tau_{m_0}) \rightarrow \mathbb{R}_+^n$ are, respectively, linear continuous and positive homogeneous continuous operators. Let, moreover, the system of impulsive inequalities (1.18), (1.19) have only the trivial solution under the condition (1.7). Then the problem (1.1), (1.2); (1.3) is correct.

Corollary 1.3. Let the system (1.1), (1.2); (1.3) have a unique solution x^0 defined on the whole closed interval $[a, b]$, where $h(x) \equiv x(t_0) - c_0$, and $t_0 \in [a, b]$ and $c_0 \in \mathbb{R}^n$ are such that $I_l(c_0) = 0$ if $t_0 = \tau_l$ for some $l \in \{1, \dots, m_0\}$. Let, moreover,

$$\lim_{\rho \rightarrow +\infty} \sup \left(\inf \{ \|x + I_l(y)\| : \|x\| \geq \rho, \|y\| = \rho \} \right) > \|x^0\|_s \quad (1.24)$$

for every $l \in \{1, \dots, m_0\}$ such that $\tau_l > t_0$. Then the problem (1.1), (1.2); (1.24) is correct.

Remark 1.4. It is evident that the condition (1.24) is valid if $I_l(y) \equiv 0$ for every $l \in \{1, \dots, m_0\}$ such that $\tau_l > t_0$. If the last condition is not fulfilled, i.e., $I_l(y) \not\equiv 0$ for some $l \in \{1, \dots, m_0\}$, then the strict inequality (1.24) cannot be replaced by a non-strict one for this l . Below, we will give the corresponding example.

Example. Let $n = 1$, $m_0 > 2$ be an arbitrary natural number, $\tau_l \in]a, b[$ ($l = 1, \dots, m_0$), $h(x) \equiv x(t_0) - c_0$, $t_0 = b$, $c_0 = 0$; $h_k(x) \equiv x(t_k) - c_k$ ($k = 1, 2, \dots$), $t_k \rightarrow b$ ($k \rightarrow +\infty$) and $c_k \rightarrow c_0$ ($k \rightarrow +\infty$); $f(t, x) = f_k(t, x) \equiv 0$ ($k = 1, 2, \dots$); $I_l(x) = I_{kl}(x) \equiv 0$ ($l = 1, \dots, m_0 - 1$; $k = 1, 2, \dots$);

$$I_{m_0}(x) = \begin{cases} 0 & \text{for } x < 0, \\ (1 + c_{i+1} - c_i)(i - x) - i - c_i & \text{for } x \in [i, i + 1[\text{ (} i = 0, 1, \dots \text{)} \end{cases}$$

and

$$I_{km_0}(x) = \begin{cases} I_{m_0}(x) & \text{for } x \in] - \infty, k - 1[\cup] k + 1, +\infty[, \\ (1 - c_{k-1} - c_k)(k - x) + c_k - k & \text{for } x \in [k - 1, k[, \\ (1 + c_{k+1} + c_k)(k - x) + c_k - k & \text{for } x \in [k, k + 1[\end{cases} \\ (k = 1, 2, \dots).$$

Then $x^0(t) \equiv 0$, $((f_k, \{I_{lk}\}_{l=1}^{m_0}; h_k))_{k=1}^{+\infty} \in W_r(f, \{I_l\}_{l=1}^{m_0}, h; x^0)$. Moreover, the problem (1.1_k), (1.2_k); (1.3_k) has the unique solution

$$x_k(t) = \begin{cases} c_k & \text{for } a \leq t \leq \tau_{m_0}, \\ k & \text{for } \tau_{m_0} < t \leq b \end{cases}$$

for every natural k . As to the condition (1.24), it is transformed into the equality for $t = \tau_{m_0}$ only.

ACKNOWLEDGEMENT

The present work was supported by the Shota Rustaveli National Science Foundation (Grant No. FR/182/5-101/11).

REFERENCES

1. M. ASHORDIA, On the stability of solutions of linear boundary value problems for a system of ordinary differential equations. *Georgian Math. J.* **1** (1994), No. 2, 115–126.
2. M. ASHORDIA, Criteria of correctness of linear boundary value problems for systems of generalized ordinary differential equations. *Czechoslovak Math. J.* **46(121)** (1996), No. 3, 385–404.
3. M. ASHORDIA, Criteria of correctness of linear boundary value problems for systems of ordinary differential equations. *Rep. Enlarged Sess. Semin. I. Vekua Appl. Math.* **15** (2000), No. 1-3, 40–43.
4. M. T. ASHORDIA AND N. A. KEKELIA, On the well-posedness of the Cauchy problem for linear systems of generalized ordinary differential equations on an infinite interval. (Russian) *Differ. Uravn.* **40** (2004), No. 4, 443–454, 574; translation in *Differ. Equ.* **40** (2004), No. 4, 477–490.
5. M. ASHORDIA, On the general and multipoint boundary value problems for linear systems of generalized ordinary differential equations, linear impulse and linear difference systems. *Mem. Differential Equations Math. Phys.* **36** (2005), 1–80.
6. M. ASHORDIA, On the two-point boundary value problems for linear impulsive systems with singularities. *Georgian Math. J.* **19** (2012), No. 1, 19–40.

7. SH. AKHALAIA, M. ASHORDIA, AND N. KEKELIA, On the necessary and sufficient conditions for the stability of linear generalized ordinary differential, linear impulsive and linear difference systems. *Georgian Math. J.* **16** (2009), No. 4, 597–616.
8. M. ASHORDIA AND G. EKHVAIA, Criteria of correctness of linear boundary value problems for systems of impulsive equations with finite and fixed points of impulses actions. *Mem. Differential Equations Math. Phys.* **37** (2006), 154–157.
9. M. ASHORDIA AND G. EKHVAIA, On the solvability of a multipoint boundary value problem for systems of nonlinear impulsive equations with finite and fixed points of impulses actions. *Mem. Differential Equations Math. Phys.* **43** (2008), 153–158.
10. D. D. BAĪNOV AND P. S. SIMEONOV, Systems with impulse effect. Stability, theory and applications. Ellis Horwood Series: Mathematics and its Applications. *Ellis Horwood Ltd., Chichester; Halsted Press [John Wiley & Sons, Inc.], New York*, 1989.
11. M. BENCHOHRA, J. HENDERSON, AND S. NTOUYAS, Impulsive differential equations and inclusions. Contemporary Mathematics and Its Applications, 2. *Hindawi Publishing Corporation, New York*, 2006.
12. I. T. KIGURADZE, Boundary value problems for systems of ordinary differential equations. (Russian) Translated in *J. Soviet Math.* **43** (1988), No. 2, 2259–2339. Itogi Nauki i Tekhniki, *Current problems in mathematics. Newest results, Vol. 30 (Russian)*, 3–103, 204, *Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow*, 1987.
13. I. KIGURADZE, The initial value problem and boundary value problems for systems of ordinary differential equations. Vol. I Linear theory. (Russian) “*Metsniereba*”, *Tbilisi*, 1997.
14. M. A. KRASNOSEL'SKIĪ AND S. G. KREĪN, On the principle of averaging in nonlinear mechanics. (Russian) *Uspehi Mat. Nauk (N.S.)* **10** (1955), No. 3(65), 147–152.
15. YA. KURCVEĪL' AND Z. VOREL, Continuous dependence of solutions of differential equations on a parameter. (Russian) *Czechoslovak Math. J.* **7 (82)** (1957), 568–583.
16. V. LAKSHMIKANTHAM, D. D. BAĪNOV, AND P. S. SIMEONOV, Theory of impulsive differential equations. Series in Modern Applied Mathematics, 6. *World Scientific Publishing Co., Inc., Teaneck, NJ*, 1989.
17. N. A. PERESTYUK, V. A. PLOTNIKOV, A. M. SAMOILENKO, AND N. V. SKRIPNIK, Differential equations with impulse effects. Multivalued right-hand sides with discontinuities. de Gruyter Studies in Mathematics, 40. *Walter de Gruyter & Co., Berlin*, 2011.

(Received 21.03.2013)

Authors' addresses:

Malkhaz Ashordia

1. A. Razmadze Mathematical Institute of I. Javakhishvili Tbilisi State University, 6 Tamarashvili St., Tbilisi 0177, Georgia.

2. Sukhumi State University, 9 Politkovskaia St., Tbilisi 0186, Georgia.

E-mail: ashord@rmi.ge

Goderdzi Ekhvaia, Nestan Kekelia

Sukhumi State University, 9 Politkovskaia St., Tbilisi 0186, Georgia.

E-mail: goderdzi.ekhvaia@mail.ru; nest.kek@mail.ru