

# Memoirs on Differential Equations and Mathematical Physics

VOLUME 99, 2026, 1–13

---

Mokhtar Naceri

EXISTENCE RESULTS FOR NONLINEAR  
WEIGHTED ANISOTROPIC ELLIPTIC SYSTEMS  
WITH VARIABLE EXPONENTS

**Abstract.** This paper establishes the existence of distributional solutions for a class of anisotropic elliptic  $\vec{p}(\cdot)$ -systems. These systems are weighted by a positive function that belongs to the  $\vec{p}(\cdot)$ -Sobolev space and involves  $L^1(\Omega; \mathbb{R}^m)$ -data associated with the  $L^1(\Omega)$ -coefficient of the zero order term.

**2020 Mathematics Subject Classification.** 35J60, 35J66, 35J67.

**Key words and phrases.** Nonlinear elliptic systems, Weight function,  $\vec{p}(\cdot)$ -Sobolev spaces, Variable exponents, Distributional solution, Existence.

# 1 Introduction

This study focuses on establishing the existence of at least one distributional solution  $u = (u_1, \dots, u_m)^\top$  for a particular class of weighted elliptic  $\vec{p}(\cdot)$ -systems expressed in the form

$$-\sum_{i=1}^N \partial_i(v(x)\sigma_i(x, \partial_i u)) + A(x)g(x, u) = f \text{ in } \Omega, \quad (1.1)$$

$$u = 0 \text{ on } \partial\Omega.$$

Here,  $\Omega \subset \mathbb{R}^N$  ( $N \geq 2$ ) is an open bounded Lipschitz domain,  $\partial_i u = \frac{\partial u}{\partial x_i}$ ,  $i = 1, \dots, N$ ,  $f \in L^1(\Omega; \mathbb{R}^m)$ ,  $A(\cdot) \in L^1(\Omega)$ , and  $v(\cdot)$  is in  $\overset{\circ}{W}^{1, \vec{p}(\cdot)}(\Omega)$  such that there exists  $\alpha, \beta > 0$ :

$$|f(x)| \leq \alpha A(x), \quad (1.2)$$

$$v(x) \geq \beta. \quad (1.3)$$

$\sigma_i : \Omega \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ ,  $i = 1, \dots, N$ , are the Carathéodory functions such that, for almost every  $x \in \Omega$  and every  $s, s' \in \mathbb{R}^m$  ( $s, s' \neq (0, 0)$ ), we have

$$\sigma_i(x, s) \cdot s \geq c_1 |s|^{p_i(x)}, \quad (1.4)$$

$$|\sigma_i(x, s)| \leq c_2 (|s|^{p_i(x)} + |h|)^{1 - \frac{1}{p_i(x)}}, \quad h \in L^1(\Omega) \quad (1.5)$$

$$(\sigma_i(x, s) - \sigma_i(x, s')) \cdot (s - s') \geq \begin{cases} c_3 |s - s'|^{p_i(x)}, & \text{if } p_i(x) \geq 2, \\ c_4 \frac{|s - s'|^2}{(|s| + |s'|)^{2-p_i(x)}}, & \text{if } 1 < p_i(x) < 2, \end{cases} \quad (1.6)$$

where  $c_1, c_2, c_3, c_4$  are the positive constants.

The function  $g : \Omega \times \mathbb{R}^m \rightarrow \mathbb{R}^m$  is a Carathéodory function and meets the following conditions almost everywhere for  $x \in \Omega$ :

$$g(x, s) \cdot (s - s') \geq 0, \quad \forall s, s' \in \mathbb{R}^m, \quad |s| = |s'|, \quad (1.7)$$

$$\sup_{|s| \leq t} |g(x, s)| \in L^1(\Omega; \mathbb{R}^m), \quad \forall s \in \mathbb{R}^m, \text{ and } \forall t > 0, \quad (1.8)$$

$$g(x, s) \cdot s \geq \sum_{i=1}^N |\rho|^{p_i(x)+1}, \quad \forall s \in \mathbb{R}^m. \quad (1.9)$$

As a typical example, consider the following:

$$-\sum_{i=1}^N \partial_i(v(x)|\partial_i u|^{p_i(x)-2} \partial_i u) + A(x)u \sum_{i=1}^N |u|^{p_i(x)-1} = f \text{ in } \Omega,$$

$$u = 0 \text{ on } \partial\Omega,$$

where  $f, A(\cdot), v(\cdot)$ , and  $p_i(\cdot)$ ,  $i = 1, \dots, N$ , are restricted as in Theorem 3.1.

In this work, we focus on the  $\vec{p}(x)$ -anisotropic differential operator, which has a wide range of applications in applied sciences. For example, such operators are frequently employed in the study of electro-rheological fluids and image processing, as highlighted in references [10, 17, 27]. Various existence results for systems involving these operators under different conditions and data have been established, as detailed in [5, 19, 24–26]. For the anisotropic scalar case, the related results can be found in [1–4, 13]. Furthermore, the existence of solutions for variable exponent anisotropic nonlinear weighted elliptic equations has been demonstrated in [20–23], with the corresponding isotropic scalar case discussed in [6].

In the present paper, we establish existence results for distributional solutions to a class of anisotropic nonlinear elliptic systems with variable exponents described by (1.1). These systems

are weighted by a positive function  $v(\cdot) \in \overset{\circ}{W}^{1,\vec{p}(\cdot)}(\Omega)$ , and the given datum satisfies  $f \in L^1(\Omega; \mathbb{R}^m)$ . A key assumption is the interaction described in (1.2) between the datum and the  $L^1(\Omega)$  coefficient  $A(\cdot)$  of the zero-order term inducing a regularizing effect on (1.1). Our approach to proving the existence of solutions depends critically on the requirement that the weight function belongs to the  $\vec{p}(\cdot)$ -Sobolev space.

We developed our proof based on a sequence of approximate solutions  $(u_n)$ , which requires demonstrating their existence through the main theorem on pseudo-monotone operators and the findings presented in [25]. Subsequently, we employ a priori estimates to establish the boundedness of  $(u_n)$  and the almost everywhere convergence of their partial derivatives  $\partial_i u_n$  for  $i = 1, \dots, N$ , which can be interpreted as a strong  $L^1$ -convergence. With this convergence established, we take the limit in the strong  $L^1$  sense for  $v_n(x)\sigma_i(x, \partial_i u_n)$  for  $i = 1, \dots, N$  and in  $A_n(x)g(x, u_n)$ . Ultimately, we conclude that the approximate solutions  $u_n$  converge to the solution of (1.1).

Our work is organized as follows. Section 2 focuses on the mathematical preliminaries, covering isotropic and anisotropic variable exponent Lebesgue–Sobolev spaces, along with several embedding theorems. The primary theorem, along with its proof, is presented in Section 3.

## 2 Preliminaries

This section aims to introduce fundamental definitions and properties related to isotropic and anisotropic variable exponent Lebesgue–Sobolev spaces (refer to [11, 12, 15]).

Let  $\Omega \subset \mathbb{R}^N$  ( $N \geq 2$ ) be a bounded open subset. We denote

$$\mathcal{C}_+(\overline{\Omega}) = \left\{ \text{continuous function } p(\cdot) : \overline{\Omega} \mapsto \mathbb{R}, \ / \ 1 < p^- (= p^- = \min_{x \in \overline{\Omega}} p(x)) \right\}.$$

Let  $p(\cdot) \in \mathcal{C}_+(\overline{\Omega})$ . Variable exponent Lebesgue space with  $L^{p(\cdot)}(\Omega)$  is defined by

$$L^{p(\cdot)}(\Omega) := \left\{ \text{measurable functions } u : \Omega \mapsto \mathbb{R}; \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\},$$

where

$$u \mapsto \varrho_{p(\cdot)}(u) := \int_{\Omega} |u(x)|^{p(x)} dx \text{ is called the convex modular.}$$

It forms a reflexive Banach space when equipped with the Luxemburg norm

$$u \mapsto \|u\|_{p(\cdot)} := \inf \left\{ s > 0 \mid \varrho_{p(\cdot)}\left(\frac{u}{s}\right) \leq 1 \right\}.$$

The Hölder type inequality

$$\left| \int_{\Omega} uv dx \right| \leq \left( \frac{1}{p^-} + \frac{1}{p'^-} \right) \|u\|_{p(\cdot)} \|v\|_{p'(\cdot)} \leq 2 \|u\|_{p(\cdot)} \|v\|_{p'(\cdot)}$$

holds true.

The subsequent results are presented in [11, 12].

If  $u \in L^{p(\cdot)}(\Omega)$ , then we obtain

$$\begin{aligned} \min &\leq (\varrho_{p(\cdot)}(u)^{\frac{1}{p^+}}, \varrho_{p(\cdot)}(u)^{\frac{1}{p^-}}) \leq \|u\|_{p(\cdot)} \leq \max (\varrho_{p(\cdot)}(u)^{\frac{1}{p^+}}, \varrho_{p(\cdot)}(u)^{\frac{1}{p^-}}), \\ &\min (\|u\|_{p(\cdot)}^{p^-}, \|u\|_{p(\cdot)}^{p^+}) \leq \varrho_{p(\cdot)}(u) \leq \max (\|u\|_{p(\cdot)}^{p^-}, \|u\|_{p(\cdot)}^{p^+}). \end{aligned} \quad (2.1)$$

We now proceed to introduce the  $\vec{p}(\cdot)$ -Sobolev spaces  $W^{1,\vec{p}(\cdot)}(\Omega)$ .

Let  $\vec{p}(x) = (p_1(x), \dots, p_N(x)) \in (C(\overline{\Omega}, [1, +\infty)))^N$ , and for every  $x$  in  $\overline{\Omega}$  we set

$$\begin{aligned} p_+(x) &= \max_{1 \leq i \leq N} p_i(x), \quad p_-(x) = \min_{1 \leq i \leq N} p_i(x), \\ p_-^- &= \min_{x \in \overline{\Omega}} p_-(x), \quad p_+^+ = \max_{x \in \overline{\Omega}} p_+(x), \\ \bar{p}(x) &= \frac{N}{\sum_{i=1}^N \frac{1}{p_i(x)}}, \quad \bar{p}^*(x) = \begin{cases} \frac{N\bar{p}(x)}{N - \bar{p}(x)} & \text{for } \bar{p}(x) < N, \\ +\infty & \text{for } \bar{p}(x) \geq N. \end{cases} \end{aligned}$$

The Banach space  $W^{1, \vec{p}(\cdot)}(\Omega)$  is defined by

$$W^{1, \vec{p}(\cdot)}(\Omega) = \left\{ u \in L^{p_+(\cdot)}(\Omega), \partial_i u \in L^{p_i(\cdot)}(\Omega), i = 1, \dots, N \right\}$$

under the norm

$$u \mapsto \|u\|_{\vec{p}(\cdot)} = \|u\|_{p_+(\cdot)} + \sum_{i=1}^N \|\partial_i u\|_{p_i(\cdot)}. \quad (2.2)$$

The Banach space  $\mathring{W}^{1, \vec{p}(\cdot)}(\Omega)$  is defined as follows:

$$\mathring{W}^{1, \vec{p}(\cdot)}(\Omega) = W^{1, \vec{p}(\cdot)}(\Omega) \cap W_0^{1,1}(\Omega),$$

when equipped with the norm (2.2).

The following results can be found in [14, 15].

Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain and  $\vec{p}(\cdot) \in (\mathcal{C}_+(\overline{\Omega}))^N$ .

(\*) If  $r \in \mathcal{C}_+(\overline{\Omega})$  and  $\forall x \in \overline{\Omega}$ ,  $r(x) < \max(p_+(x), \bar{p}^*(x))$ , then the embedding

$$\mathring{W}^{1, \vec{p}(\cdot)}(\Omega) \hookrightarrow L^{r(\cdot)}(\Omega)$$

is compact.

(\*) If we have

$$\forall x \in \overline{\Omega}, \quad p_+(x) < \bar{p}^*(x), \quad (2.3)$$

then the following inequality holds:

$$\|u\|_{p_+(\cdot)} \leq C \sum_{i=1}^N \|\partial_i u\|_{p_i(\cdot)}, \quad \forall u \in \mathring{W}^{1, \vec{p}(\cdot)}(\Omega),$$

where  $C > 0$  is independent of  $u$ . Thus

$$u \mapsto \sum_{i=1}^N \|\partial_i u\|_{p_i(\cdot)} \text{ is an equivalent norm to (2.2) on } \mathring{W}^{1, \vec{p}(\cdot)}(\Omega). \quad (2.4)$$

In our paper, we denote by  $L^{p(\cdot)}(\Omega, \mathbb{R}^m)$  and  $\mathring{W}^{1, \vec{p}(\cdot)}(\Omega, \mathbb{R}^m)$  ( $m \geq 1$ ) the  $\mathbb{R}^m$ -valued version of  $L^{p(\cdot)}(\Omega)$  and  $\mathring{W}^{1, \vec{p}(\cdot)}(\Omega)$ , respectively.

The space  $X = \mathring{W}^{1, \vec{p}(\cdot)}(\Omega, \mathbb{R}^m)$  becomes a Banach space, when equipped with the norm

$$\|\cdot\|_X = \|\|\cdot\|_{\vec{p}(\cdot)}\|.$$

Also  $Y = L^{p(\cdot)}(\Omega, \mathbb{R}^m)$ , when equipped with the norm

$$\|\cdot\|_Y = \|\|\cdot\|_{p(\cdot)}\|.$$

For any  $t > 0$ , the scalar truncation function  $T_t$  on  $[0, \infty)$  is defined as

$$T_t(s) := \begin{cases} s & \text{if } s \leq t, \\ t & \text{if } s > t. \end{cases}$$

For any  $t > 0$ , define the spherical truncation function  $T_t : \mathbb{R}^m \rightarrow \mathbb{R}^m$  by

$$T_t(s) := \begin{cases} s & \text{if } |s| \leq t, \\ \frac{s}{|s|} t & \text{if } |s| > t. \end{cases} \quad (2.5)$$

### 3 Statement of results and proof

**Definition.** The vector-valued function  $u = (u_1, \dots, u_m)^\top : \Omega \rightarrow \mathbb{R}^m$  is a solution of system (1.1) in the sense of distributions if and only if  $u \in W_0^{1,1}(\Omega; \mathbb{R}^m)$ , and for all  $\varphi \in C_c^\infty(\Omega; \mathbb{R}^m)$ ,

$$\int_{\Omega} \sum_{i=1}^N v(x) \sigma_i(x, \partial_i u) \cdot \partial_i \varphi \, dx + \int_{\Omega} \sum_{i=1}^N A(x) g(x, u) \cdot \varphi \, dx = \int_{\Omega} f(x) \cdot \varphi \, dx.$$

Our main Theorem is the following.

**Theorem 3.1.** Let  $p_i(\cdot)$ ,  $i = 1, \dots, N$  be in  $\mathcal{C}_+(\overline{\Omega})$  such that  $\overline{p}(\cdot) < N$  in  $\overline{\Omega}$ , and let (2.3) hold. Assume  $f$  is in  $L^1(\Omega; \mathbb{R}^m)$ ,  $A(\cdot)$  is in  $L^1(\Omega)$ , and  $v(\cdot)$  is in  $\dot{W}^{1, \overline{p}(\cdot)}(\Omega)$  such that (1.2), (1.3) and (2.3) hold.

Let  $\sigma_i$ ,  $i = 1, \dots, N$ , and  $g$  be the Carathéodory functions, where  $\sigma_i$  satisfy (1.4)–(1.6), and  $g$  satisfies (1.7)–(1.9). Then problem (1.1) has at least one solution  $u \in \dot{W}^{1, \overline{p}(\cdot)}(\Omega; \mathbb{R}^m)$  in the sense of distributions.

#### 3.1 Existence of approximate solutions

We consider the function  $\theta(\cdot)$  defined as follows:

$$\begin{aligned} \theta(x) &= \frac{nx}{x+n}, \quad x \geq 0, \\ f_n(x) &= \theta(f(x)), \quad A_n(x) = \frac{1}{\alpha} \theta(A(x)), \quad v_n(x) = \theta(v(x)), \quad n \in \mathbb{N}^*. \end{aligned} \quad (3.1)$$

Noticing the increase of  $\theta$ , (1.2) and (1.3), we can obtain

$$|f_n(x)| \leq \theta(\alpha A(x)) = \alpha A_n(x)$$

and

$$\theta(\beta) \leq v_n(x) \leq \theta(n).$$

Then we conclude that for all  $x \in \overline{\Omega}$ ,

$$|f_n(x)| \leq \alpha A_n(x) \quad (3.2)$$

and

$$\frac{\beta}{1+\beta} \leq v_n(x) \leq n. \quad (3.3)$$

**Lemma 3.1.** Let  $p_i(\cdot)$ ,  $i = 1, \dots, N$ , be in  $\mathcal{C}_+(\overline{\Omega})$  such that  $\overline{p}(\cdot) < N$  in  $\overline{\Omega}$ , and let (2.3) hold. Assume  $f$  is in  $L^1(\Omega; \mathbb{R}^m)$ ,  $A(\cdot)$  is in  $L^1(\Omega)$ , and  $v(\cdot)$  is in  $\dot{W}^{1, \overline{p}(\cdot)}(\Omega)$  such that (1.2), (1.3) and (2.3) hold.

Let  $\sigma_i$ ,  $i = 1, \dots, N$ , and  $g$  be the Carathéodory functions, where  $\sigma_i$  satisfy (1.4)–(1.6), and  $g$  satisfies (1.7)–(1.9).

Then there exists at least one weak solution  $u_n \in \mathring{W}^{1,\vec{p}(\cdot)}(\Omega; \mathbb{R}^m)$  to the approximated problems

$$\begin{aligned} - \sum_{i=1}^N \partial_i (v_n(x) \sigma_i(x, \partial_i u_n)) + A_n(x) g(x, u_n) &= f_n \text{ in } \Omega, \\ u_n &= 0 \text{ on } \partial\Omega, \end{aligned} \quad (3.4)$$

in the following sense:

$$\begin{aligned} \text{For every } \varphi \in \mathring{W}^{1,\vec{p}(\cdot)}(\Omega; \mathbb{R}^m) \cap L^\infty(\Omega; \mathbb{R}^m), \\ \sum_{i=1}^N \int_{\Omega} v_n(x) \sigma_i(x, \partial_i u_n) \cdot \partial_i \varphi \, dx + \int_{\Omega} A_n(x) g(x, u_n) \cdot \varphi \, dx &= \int_{\Omega} f_n \cdot \varphi \, dx. \end{aligned} \quad (3.5)$$

*Proof.* Consider the system

$$\begin{aligned} - \sum_{i=1}^N \partial_i (v_n(x) \sigma_i(x, \partial_i u_{n_k})) + A_n(x) T_k(g(x, u_{n_k})) &= f_n \text{ in } \Omega, \\ u_{n_k} &= 0 \text{ on } \partial\Omega. \end{aligned} \quad (3.6)$$

In a similar manner to the results obtained in [25], applying the main Theorem on the pseudo-monotone operators ((Theorem 27.A in [28], see also [8, 9, 18])), we conclude that there exists a solution  $u_{n_k} \in \mathring{W}^{1,\vec{p}(\cdot)}(\Omega; \mathbb{R}^m)$  to system (3.6), which satisfies

$$\begin{aligned} \sum_{i=1}^N \int_{\Omega} v_n(x) \sigma_i(x, \partial_i u_{n_k}) \cdot \partial_i \varphi \, dx \\ + \int_{\Omega} A_n(x) T_k(g(x, u_{n_k})) \cdot \varphi \, dx &= \int_{\Omega} f_n \cdot \varphi \, dx, \quad \forall \varphi \in \mathring{W}^{1,\vec{p}(\cdot)}(\Omega; \mathbb{R}^m). \end{aligned} \quad (3.7)$$

Now, choosing  $\varphi = u_{n_k}$  as a test function in (3.7) and using (3.3), (1.4), after dropping the nonnegative term (since  $A_n(x) T_k(g(x, u_{n_k})) \cdot u_{n_k} \geq 0$ , due to (1.9) and the fact that  $A_n(x) \geq 0$ ), we get

$$\frac{c_1 \beta}{(1 + \beta)} \sum_{i=1}^N \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} \, dx \leq n \int_{\Omega} |u_{n_k}| \, dx.$$

Using Young's inequality, for all  $\varepsilon > 0$ , we get

$$\begin{aligned} \sum_{i=1}^N \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} \, dx &\leq \frac{n(1 + \beta)}{c_1 \beta} \int_{\Omega} |u_{n_k}| \, dx \\ &\leq \frac{n(1 + \beta)}{c_1 \beta} \left( \varepsilon \int_{\Omega} |u_{n_k}|^{p^-} \, dx + C(\varepsilon) \right) \leq \frac{n(1 + \beta)}{c_1 \beta} \left( \varepsilon c \int_{\Omega} |\partial_i u_{n_k}|^{p^-} \, dx + C(\varepsilon) \right) \\ &\leq \frac{n(1 + \beta)}{c_1 \beta} \left( \varepsilon c \left( 1 + \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} \, dx \right) + C(\varepsilon) \right) \\ &\leq \frac{n(1 + \beta)}{c_1 \beta} \left( \varepsilon c \left( 1 + \sum_{i=1}^N \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} \, dx \right) + C(\varepsilon) \right). \end{aligned}$$

Choosing  $\varepsilon = \frac{c_1 \beta}{2nc(1 + \beta)}$ , we obtain

$$\sum_{i=1}^N \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} \, dx \leq c(n). \quad (3.8)$$

On the other hand, from (2.1) for all  $i = 1, \dots, N$  we have

$$1 + \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} dx \geq \| |\partial_i u_{n_k}| \|_{p_i(x)}^{p_i^-} \quad (3.9)$$

and

$$1 + \| |\partial_i u_{n_k}| \|_{p_i(x)}^{p_i^-} \geq \| |\partial_i u_{n_k}| \|_{\bar{p}_i(x)}^{p_i^-}. \quad (3.10)$$

Combining (3.9) and (3.10), we obtain

$$\sum_{i=1}^N \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} dx \geq \sum_{i=1}^N \| |\partial_i u_{n_k}| \|_{\bar{p}_i(x)}^{p_i^-} - 2N. \quad (3.11)$$

By (3.11) and (2.4) (due (2.3)), we deduce that

$$\sum_{i=1}^N \int_{\Omega} |\partial_i u_{n_k}|^{p_i(x)} dx \geq \left( \frac{1}{N} \| |u_{n_k}| \|_{\bar{p}(\cdot)} \right)^{p^-} - 2N. \quad (3.12)$$

From (3.8) and (3.12), we conclude

$$\| |u_{n_k}| \|_{\bar{p}(\cdot)} \leq C(n). \quad (3.13)$$

Through (3.13) we can conclude that there is a subsequence (still denoted by  $u_{n_k}$ )  $u_n \in \mathring{W}^{1, \bar{p}(\cdot)}(\Omega; \mathbb{R}^m)$  such that

$$u_{n_k} \rightharpoonup u_n \text{ weakly in } \mathring{W}^{1, \bar{p}(\cdot)}(\Omega; \mathbb{R}^m) \text{ and a.e in } \Omega.$$

In a similar manner to the results obtained in [25], thanks to (1.4)–(1.6) and (1.7)–(1.9), we can obtain

$$\partial_i u_{n_k} \rightarrow \partial_i u_n \text{ strongly in } L^{p_i(\cdot)}(\Omega; \mathbb{R}^m) \text{ and a.e. in } \Omega.$$

So,

$$v_n(x) \sigma_i(x, \partial_i u_{n_k}) \rightharpoonup v_n(x) \sigma_i(x, \partial_i u_n) \text{ in } L^{p'_i(\cdot)}(\Omega; \mathbb{R}^m).$$

Taking  $T_t(u_{n_k})$  as a test function in (3.7), by (1.8), (2.5), and the fact that

$$|T_t(s)| \leq M + t 1_{\{|s| > M\}}, \quad \forall s \in \mathbb{R}^m \text{ and } 0 < M < t,$$

for all  $0 < M < t$  we get

$$\int_{\{|u_{n_k}| > t\}} A_n(x) |T_k(g(x, u_{n_k}))| dx \leq \frac{M}{t} \|f\|_{L^1(\Omega; \mathbb{R}^m)} + \int_{\{|u_{n_k}| > M\}} |f_n|. \quad (3.14)$$

Let  $E \subset \Omega$  be any measurable set, we write

$$\int_E A_n(x) |T_k(g(x, u_{n_k}))| dx = \int_{E \cap \{|u_{n_k}| \leq t\}} A_n(x) |g(x, u_{n_k})| dx + \int_{E \cap \{|u_{n_k}| > t\}} A_n(x) |T_k(g(x, u_{n_k}))| dx.$$

Then, by (1.8) and the decomposition (3.14), we deduce that the sequence  $\{A_n(x) T_k(g(x, u_{n_k}))\}$  is equi-integrable in  $L^1(\Omega; \mathbb{R}^m)$ , and since  $T_k(g(x, u_{n_k})) \rightarrow g(x, u_n)$  a.e. in  $\Omega$ , Vitali's theorem implies that

$$A_n(x) T_k(g(x, u_{n_k})) \rightarrow A_n(x) g(x, u_n) \text{ in } L^1(\Omega; \mathbb{R}^m).$$

Therefore, we can obtain (3.5) by passing to the limit in (3.7).  $\square$



### 3.1.1 A priori estimates

**Lemma 3.2.** *Let  $f$ ,  $A$ ,  $v$ ,  $g$  and  $p_i$ ,  $\sigma_i$ ,  $i = 1, \dots, N$ , be restricted as in Theorem 3.1. Then*

$$|g(x, u_n)| \leq \alpha, \quad (3.15)$$

$$|u_n| \leq \left( \frac{\alpha}{N} + 1 \right)^{\frac{1}{p^-}}, \quad (3.16)$$

where  $u_n$  is the weak solution to problem (3.4).

*Proof.* After choosing  $\varphi = u_n$  in the weak formulation (3.5), and dropping the nonnegative term (since  $v_n(x)\sigma_i(x, \partial_i u_n) \cdot \partial_i u_n \geq 0$ ,  $i = 1, \dots, N$ , due (1.4), and (3.3)), we obtain

$$\int_{\Omega} A_n(x) g(x, u_n) \cdot u_n \, dx \leq \int_{\Omega} |f_n| |u_n| \, dx.$$

Using (3.2) and the fact that

$$g(x, u_n) \cdot u_n \geq |g(x, u_n)| |u_n|$$

(it is produced through the following: by (1.7), we get

$$\frac{u_n}{|u_n|} \cdot g(x, u_n) - |g(x, u_n)| = \frac{1}{|u_n|} g(x, u_n) \cdot \left( u_n - |u_n| \frac{g(x, u_n)}{|g(x, u_n)|} \right) \geq 0 \Big),$$

we obtain

$$\int_{\Omega} A_n(x) |g(x, u_n)| |u_n| \, dx \leq \alpha \int_{\Omega} A_n(x) |u_n| \, dx.$$

whence

$$\int_{\Omega} A_n(x) (|g(x, u_n)| - \alpha) |u_n| \, dx \leq 0. \quad (3.17)$$

Then (3.17) implies (3.15).

Also, by the fact that  $1 + |u_n|^{p_i(x)} \geq |u_n|^{p^-}$ ,  $i = 1, \dots, N$ , due to (1.9) and (3.15), we get (3.16).  $\square$

**Remark 3.1.** (3.15) and (3.16) imply that

$$\begin{aligned} (g(x, u_n)) & \text{ is bounded in } L^\infty(\Omega, \mathbb{R}^m), \\ (u_n) & \text{ is bounded in } L^\infty(\Omega, \mathbb{R}^m). \end{aligned} \quad (3.18)$$

**Lemma 3.3.**

$$(A_n(x)g(x, u_n)) \text{ is bounded in } L^1(\Omega, \mathbb{R}^m). \quad (3.19)$$

*Proof.* Through (3.15) and (3.1), we get

$$\int_{\Omega} |A_n(x)g(x, u_n)| \, dx \leq \alpha \int_{\Omega} |A_n(x)| \, dx \leq \|A\|_{L^1(\Omega)}. \quad (3.20)$$

So, (3.20) implies (3.19).  $\square$

**Lemma 3.4.**

$$v_n \text{ is bounded in } \mathring{W}^{1, \vec{p}(\cdot)}(\Omega) \quad (3.21)$$

and

$$v_n \rightarrow v \text{ strongly in } \mathring{W}^{1, \vec{p}(\cdot)}(\Omega). \quad (3.22)$$

*Proof.* Since, for all  $x \in \bar{\Omega}$ ,

$$\partial_i v_n(x) = \frac{\partial_i v(x)}{(1 + \frac{v(x)}{n})^2}, \quad i = 1, \dots, N,$$

we have

$$|\partial_i v_n(x)| \leq |\partial_i v(x)|$$

and, therefore,

$$v_n(\cdot) \in \mathring{W}^{1, \bar{p}(\cdot)}(\Omega).$$

From this and the fact that  $0 \leq v_n(x) \leq v(x)$ , we get (3.21) and (3.22).  $\square$

**Lemma 3.5.** *Let  $f$ ,  $A$ ,  $v$ ,  $g$  and  $p_i$ ,  $\sigma_i$ ,  $i = 1, \dots, N$ , be restricted as in Theorem 3.1. Then*

$$u_n \text{ is bounded in } \mathring{W}^{1, \bar{p}(\cdot)}(\Omega; \mathbb{R}^m), \quad (3.23)$$

where  $u_n$  is the weak solution to problem (3.4).

*Proof.* After choosing  $\varphi = u_n$  in the weak formulation (3.5), and dropping the nonnegative term (since  $A_n(x)g(x, u_n) \cdot u_n \geq 0$ , due to (1.9) and the fact that  $A_n(x) \geq 0$ ), and using (3.16), (1.4), (3.1) and (3.3), we can get

$$\frac{c_1 \beta}{(1 + \beta)} \sum_{i=1}^N \int_{\Omega} |\partial_i u_n|^{p_i(x)} dx \leq \left( \frac{\alpha}{N} + 1 \right)^{\frac{1}{p^-}} \|f\|_{L^1(\Omega, \mathbb{R}^m)}.$$

Then we have

$$\sum_{i=1}^N \int_{\Omega} |\partial_i u_n|^{p_i(x)} dx \leq c. \quad (3.24)$$

By a proof similar to the that of (3.12), we can get

$$\sum_{i=1}^N \int_{\Omega} |\partial_i u_n|^{p_i(x)} dx \geq \left( \frac{1}{N} \|u_n\|_{\bar{p}(\cdot)} \right)^{p^-} - 2N. \quad (3.25)$$

Combining (3.24) and (3.25), we obtain

$$\|u_n\|_{\bar{p}(\cdot)} \leq C, \quad (3.26)$$

where  $C > 0$  is independent of  $n$ .

Then (3.26) implies (3.23).  $\square$

**Remark 3.2.** It follows from (3.23) that there exist a function  $u \in \mathring{W}^{1, \bar{p}(\cdot)}(\Omega; \mathbb{R}^m)$  and a subsequence (still denoted by  $(u_n)$ ) such that

$$u_n \rightharpoonup u \text{ weakly in } \mathring{W}^{1, \bar{p}(\cdot)}(\Omega; \mathbb{R}^m) \text{ and a.e in } \Omega. \quad (3.27)$$

**Lemma 3.6.** *For all  $i = 1, \dots, N$ ,*

$$\partial_i u_n \rightarrow \partial_i u \text{ a.e. in } \bar{\Omega}, \quad (3.28)$$

where  $u$  is the weak limit of the sequence  $(u_n)$  in  $\mathring{W}^{1, \bar{p}(\cdot)}(\Omega; \mathbb{R}^m)$ .

*Proof.* By (3.3), we obtain

$$\frac{1}{v_n(x)} \varphi \in \mathring{W}^{1, \bar{p}(\cdot)}(\Omega, \mathbb{R}^m) \text{ for all } \varphi \in \mathring{W}^{1, \bar{p}(\cdot)}(\Omega, \mathbb{R}^m) \cap L^\infty(\Omega, \mathbb{R}^m).$$

Therefore, we can choose it as a test function in (3.5) and get

$$\sum_{i=1}^N \int_{\Omega} \sigma_i(x, \partial_i u_n) \cdot \partial_i \varphi \, dx = \int_{\Omega} \Phi_n(x) \varphi \, dx,$$

where  $\Phi_n$  is defined by

$$\Phi_n(x) = \frac{1}{v_n(x)} \left( f_n(x) - A_n(x)g(x, u_n) + \sum_{i=1}^N \sigma_i(x, \partial_i u_n) \partial_i v_n(x) \right).$$

By Young's inequality, (1.5), and since  $\partial_i u_n \in L^{p_i(\cdot)}(\Omega)$ , for all  $\varepsilon > 0$  we get

$$\begin{aligned} \int_{\Omega} |\sigma_i(x, \partial_i u_n)| \, dx &= c' + \int_{\Omega} |\partial_i u_n|^{p_i(x)-1} \, dx \\ &\leq c' + C(\varepsilon) + \varepsilon \int_{\Omega} |\partial_i u_n|^{p_i(x)} \, dx \leq c' + C(\varepsilon) + \varepsilon c = C'(\varepsilon). \end{aligned}$$

Then, for any fixed choice for  $\varepsilon > 0$ , we conclude that for all  $i = 1, \dots, N$ ,

$$(\sigma_i(x, \partial_i u_n)) \text{ is bounded in } L^1(\Omega, \mathbb{R}^m). \quad (3.29)$$

From (3.1) (implying that  $f_n \in L^1(\Omega, \mathbb{R}^m)$ ), (3.29) and (3.19), we conclude that

$$\left( f_n(x) - A_n(x)g(x, u_n) + \sum_{i=1}^N \sigma_i(x, \partial_i u_n) \partial_i v_n(x) \right) \text{ is bounded in } L^1(\Omega, \mathbb{R}^m). \quad (3.30)$$

Through (3.30) and the boundedness of the sequence  $(\frac{1}{v_n(x)})$  (due to (3.3)), we obtain

$$(\Phi_n) \text{ is bounded in } L^1(\Omega, \mathbb{R}^m).$$

So, applying the results obtained in [7] to the sequence  $(u_n)$ , we can simply obtain (3.28).  $\square$

### 3.2 Proof of Theorem 3.1

For all  $i = 1, \dots, N$ , we put

$$\sigma_i(x, \partial_i u_n) = (\sigma_i^{(1)}(x, \partial_i u_n), \dots, \sigma_i^{(m)}(x, \partial_i u_n))$$

and

$$\sigma_i(x, \partial_i u) = (\sigma_i^{(1)}(x, \partial_i u), \dots, \sigma_i^{(m)}(x, \partial_i u)).$$

By (3.28), for all  $i = 1, \dots, N$  we have

$$\sigma_i(x, \partial_i u_n) \rightharpoonup \sigma_i(x, \partial_i u) \text{ weakly in } L^{p'_i(\cdot)}(\Omega; \mathbb{R}^m).$$

Then we conclude that, for all  $i = 1, \dots, N$  and all  $j = 1, \dots, m$ ,

$$\sigma_i^{(j)}(x, \partial_i u_n) \rightharpoonup \sigma_i^{(j)}(x, \partial_i u) \text{ weakly in } L^{p'_i(\cdot)}(\Omega), \quad (3.31)$$

where  $p'_i(\cdot)$  denotes the Hölder conjugate of  $p_i(\cdot)$  in  $\overline{\Omega}$ .

By (3.22), we conclude that for all  $i = 1, \dots, N$ ,

$$v_n(\cdot) \rightarrow v(\cdot) \text{ strongly in } L^{p_i(\cdot)}(\Omega). \quad (3.32)$$

Then, from (3.31) and (3.32), for all  $i = 1, \dots, N$  and all  $j = 1, \dots, m$ , we obtain

$$v_n(x) \sigma_i^{(j)}(x, \partial_i u_n) \rightarrow v(x) \sigma_i^{(j)}(x, \partial_i u) \text{ strongly in } L^1(\Omega). \quad (3.33)$$

So, (3.33) implies that

$$v_n(x)\sigma_i(x, \partial_i u_n) \rightarrow v(x)\sigma_i(x, \partial_i u) \text{ strongly in } L^1(\Omega; \mathbb{R}^m). \quad (3.34)$$

Now, we put

$$g(x, u_n) = (g_1(x, u_n), \dots, g_m(x, u_n))$$

and

$$g(x, u) = (g_1(x, u), \dots, g_m(x, u)).$$

Through (3.18) and the fact that  $|g_j(x, u_n)| \leq |g(x, u_n)|$ ,  $j = 1, \dots, m$ , we conclude that

$$(g_j(x, u_n)) \text{ is bounded in } L^\infty(\Omega). \quad (3.35)$$

Then, as  $A_n \in L^1(\Omega)$ , from (3.35) and (3.27), for all  $j = 1, \dots, m$ , we obtain

$$A_n(x)g_j(x, u_n) \rightarrow A(x)g_j(x, u) \text{ strongly in } L^1(\Omega). \quad (3.36)$$

So, (3.36) implies that

$$A_n(x)g(x, u_n) \rightarrow A(x)g(x, u) \text{ strongly in } L^1(\Omega; \mathbb{R}^m). \quad (3.37)$$

Then, through (3.34) and (3.37), we can pass to the limit in (3.5). Thus Theorem 3.1 is proved.

## Acknowledgments

The author would like to thank the referees for their comments and suggestions.

## References

- [1] M. Bendahmane and K. H. Karlsen, Anisotropic nonlinear elliptic systems with measure data and anisotropic harmonic maps into spheres. *Electron. J. Differential Equations* **2006**, no. 46, 30 pp.
- [2] M. Bendahmane and K. H. Karlsen, Nonlinear anisotropic elliptic and parabolic equations in  $\mathbb{R}^N$  with advection and lower order terms and locally integrable data. *Potential Anal.* **22** (2005), no. 3, 207–227.
- [3] M. Bendahmane and K. H. Karlsen, Renormalized solutions of an anisotropic reaction-diffusion-advection system with  $L^1$  data. *Commun. Pure Appl. Anal.* **5** (2006), no. 4, 733–762.
- [4] M. Bendahmane, K. H. Karlsen and M. Saad, Nonlinear anisotropic elliptic and parabolic equations with variable exponents and  $L^1$  data. *Commun. Pure Appl. Anal.* **12** (2013), no. 3, 1201–1220.
- [5] M. Bendahmane and F. Mokhtari, Nonlinear elliptic systems with variable exponents and measure data. *Moroccan J. Pure and Appl. Anal. (MJPA)* **1** (2015), no. 2, 108–125.
- [6] L. Boccardo, P. Imparato and L. Orsina, Nonlinear weighted elliptic equations with Sobolev weights. *Boll. Unione Mat. Ital.* **15** (2022), no. 4, 503–514.
- [7] L. Boccardo and F. Murat, Almost everywhere convergence of the gradients of solutions to elliptic and parabolic equations. *Nonlinear Anal.* **19** (1992), no. 6, 581–597.
- [8] H. Brezis, Équations et inéquations non linéaires dans les espaces vectoriels en dualité. (French) *Ann. Inst. Fourier (Grenoble)* **18** (1968), fasc. 1, 115–175.
- [9] F. E. Browder, Nonlinear elliptic boundary value problems. *Bull. Amer. Math. Soc.* **69** (1963), 862–874.
- [10] Y. Chen, S. Levine and M. Rao, Variable exponent, linear growth functionals in image restoration. *SIAM J. Appl. Math.* **66** (2006), no. 4, 1383–1406.

- [11] D. Cruz-Uribe, A. Fiorenza, M. Ruzhansky and J. Wirth, *Variable Lebesgue Spaces and Hyperbolic Systems*. Selected lecture notes from the Advanced Courses on Approximation Theory and Fourier Analysis held at the Centre de Recerca Matemàtica, Barcelona, November 7–11, 2011. Edited by Sergey Tikhonov. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser/Springer, Basel, 2014.
- [12] L. Diening, P. Harjulehto, P. Hästö and M. Růžička, *Lebesgue and Sobolev Spaces with Variable Exponents*. Lecture Notes in Mathematics, 2017. Springer, Heidelberg, 2011.
- [13] G. Dolzmann, N. Hungerbühler and S. Müller, Non-linear elliptic systems with measure-valued right hand side. *Math. Z.* **226** (1997), no. 4, 545–574.
- [14] X. Fan, Anisotropic variable exponent Sobolev spaces and  $\vec{p}(x)$ -Laplacian equations. *Complex Var. Elliptic Equ.* **56** (2011), no. 7-9, 623–642.
- [15] X. Fan and D. Zhao, On the spaces  $L^{p(x)}(\Omega)$  and  $W^{m,p(x)}(\Omega)$ . *J. Math. Anal. Appl.* **263** (2001), no. 2, 424–446.
- [16] J.-L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non Linéaires*. (French) Dunod, Paris; Gauthier-Villars, Paris, 1969.
- [17] M. Mihăilescu and V. Rădulescu, A multiplicity result for a nonlinear degenerate problem arising in the theory of electrorheological fluids. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **462** (2006), no. 2073, 2625–2641.
- [18] G. J. Minty, on a “monotonicity” method for the solution of non-linear equations in Banach spaces. *Proc. Nat. Acad. Sci. U.S.A.* **50** (1963), 1038–1041.
- [19] M. Naceri, Anisotropic nonlinear elliptic systems with variable exponents, degenerate coercivity and  $L^{q(\cdot)}$  data. *Ann. Acad. Rom. Sci. Ser. Math. Appl.* **14** (2022), no. 1-2, 107–140.
- [20] M. Naceri, Existence results for anisotropic nonlinear weighted elliptic equations with variable exponents and  $L^1$  data. *Proc. Rom. Acad. Ser. A Math. Phys. Tech. Sci. Inf. Sci.* **23** (2022), no. 4, 337–346.
- [21] M. Naceri, Anisotropic nonlinear weighted elliptic equations with variable exponents. *Georgian Math. J.* **30** (2023), no. 2, 277–285.
- [22] M. Naceri, Anisotropic nonlinear elliptic equations with variable exponents and two weighted first order terms. *Filomat* **38** (2024), no. 3, 1043–1054.
- [23] M. Naceri, Nonlinear elliptic equations with variable exponents anisotropic Sobolev weights and natural growth terms. *Tatra Mt. Math. Publ.* **88** (2024), 109–126.
- [24] M. Naceri, Variable exponents anisotropic nonlinear elliptic systems with  $L^{\vec{p}(\cdot)}$ -data. *Appl. Anal.* **104** (2025), no. 3, 564–581.
- [25] M. Naceri and M. B. Benboubker, Distributional solutions of anisotropic nonlinear elliptic systems with variable exponents: existence and regularity. *Adv. Oper. Theory* **7** (2022), no. 2, Paper no. 17, 34 pp.
- [26] M. Naceri and F. Mokhtari, Anisotropic nonlinear elliptic systems with variable exponents and degenerate coercivity. *Appl. Anal.* **100** (2021), no. 11, 2347–2367.
- [27] M. Růžička, *Electrorheological Fluids: Modeling and Mathematical Theory*. Lecture Notes in Mathematics, 1748. Springer-Verlag, Berlin, 2000.
- [28] E. Zeidler, *Nonlinear Functional Analysis and its Applications*. II/B. *Nonlinear Monotone Operators*. Translated from the German by the author and Leo F. Boron. Springer-Verlag, New York, 1990.

(Received 10.04.2025; revised 05.08.2025; accepted 12.08.2025)

#### Author’s addresses:

ENS of Laghouat, BP 4033 Station post avenue of Martyrs, 03000, Laghouat, Algeria.  
*E-mails:* nasrimokhtar@gmail.com, m.naceri@ens-lagh.dz