

R. DUDUCHAVA

CONTINUING FUNCTIONS FROM A HYPERSURFACE
WITH THE BOUNDARY

We extend m -tuples of functions from the Besov spaces in both faces of a smooth hypersurface \mathcal{S} with a smooth boundary $\Gamma = \partial\mathcal{S} \neq \emptyset$ in \mathbb{R}^n into the ambient domain slit by the hypersurface $\mathbb{R}_\mathcal{S}^n := \mathbb{R}^n \setminus \overline{\mathcal{S}}$. These tuples satisfy the compatibility conditions on the boundary Γ . The traces are defined by arbitrary Dirichlet system of boundary operators and extension is performed by two different methods, one implicit and the other explicit. Explicit extension is based on the solution to the Dirichlet BVP for the poly-harmonic equation and permits the extension of distributions from the Besov space $\mathbb{B}_{p,p}^s(\mathcal{S})$ with a negative $s < 0$.

Coretractions have essential applications in boundary value problems for partial differential equations when, for example, it is necessary to reduce a BVP with non-homogeneous boundary conditions to a BVP with the homogeneous boundary conditions.

The extended version of the present paper will appear in [2].

PRELIMINARIES

We distinguish two faces \mathcal{S}^- and \mathcal{S}^+ of the hypersurface and the normal vector field ν is directed, as usual, from the face \mathcal{S}^+ to \mathcal{S}^- . Let $\mathbb{R}_\mathcal{C}^n := \mathbb{R}^n \setminus \overline{\mathcal{C}}$ be the space, “cut” by a hypersurface \mathcal{S} . Let \mathcal{S} be a subsurface of an equally smooth surface \mathcal{C} without boundary $\partial\mathcal{C} = \emptyset$, whose borders is a compact inner domain Ω^+ . Let $\Omega^- = \mathbb{R}^3 \setminus \overline{\Omega^+}$ be the complemented domain and ν -the outer unit normal vector field, which extends the existing field from the subsurface \mathcal{S} .

For the definition of the Bessel potential $\mathbb{H}_p^\theta(\mathcal{C})$, $\widetilde{\mathbb{H}}_p^\theta(\mathcal{S})$ and the Besov $\mathbb{B}_{p,p}^\theta(\mathcal{C})$, $\widetilde{\mathbb{B}}_{p,p}^\theta(\mathcal{S})$ spaces used below, we refer, e.g., to [5].

For a pair of Besov spaces we introduce the following shortcut $\mathbb{B}_{p,p}^s(\mathcal{S}) := \mathbb{B}_{p,p}^s(\mathcal{S}) \otimes \widetilde{\mathbb{B}}_{p,p}^s(\mathcal{S})$. The notation $[s]^- \in \mathbb{Z}$ is used for the largest positive or negative integers, less than s , i.e., $s - 1 \leq [s]^- < s$.

2010 *Mathematics Subject Classification*: 46E35, 35J30, 35C15.

Key words and phrases. Hypersurface, trace, retraction, coretraction, dirichlet system, dirichlet problem, extension from a hypersurface, poly-harmonic equation, admissible matrix operator.

A partial differential operator with matrix coefficients $\mathbf{A}(x, D)$ is called **normal** on \mathcal{S} if its homogeneous principal symbol is non-degenerated on the normal vector field $\inf |\det \mathcal{A}_{\text{pr}}(x, \nu(x))| \neq 0$ for all $x \in \mathcal{S}$. Obviously, the class of elliptic operators $\inf |\det \mathcal{A}_{\text{pr}}(x, \xi)| \neq 0$ for all $x \in \mathcal{S}$, and all $|\xi| = 1$, is a subclass of the class of normal operators. For operators with constant symbols these classes coincide.

It is well known (see [5]) that the trace operators $\gamma_{\mathcal{S}}^{\pm} \mathbf{B}(x, D)$ map the following spaces continuously

$$\gamma_{\mathcal{S}}^{\pm} \mathbf{B}(x, D) : \mathbb{H}_{p, \text{loc}}^s(\overline{\mathbb{R}_{\mathcal{S}}^n}) \longrightarrow \mathbb{B}_{p, p}^{s - \frac{1}{p} - r}(\mathcal{S}), \quad 1 < p < \infty,$$

provided $s > r + 1/p$ and the ‘‘jump’’ function meets the compatibility condition:

$$[\gamma_{\mathcal{S}} \mathbf{B}(x, D) \Phi] := \gamma_{\mathcal{S}}^+ \mathbf{B}(x, D) \Phi - \gamma_{\mathcal{S}}^- \mathbf{B}(x, D) \Phi \in \widetilde{\mathbb{B}}_{p, p}^{s - \frac{1}{p} - r}(\mathcal{S}), \quad \forall \Phi \in \mathbb{H}_{p, \text{loc}}^s(\overline{\mathbb{R}_{\mathcal{S}}^n}).$$

The trace operator can be viewed as a continuous mapping by the operator pairs

$$\mathcal{R}_{\mathcal{S}}(\mathbf{B}) \Phi := \{ \gamma_{\mathcal{S}}^+ \mathbf{B}(x, D) \Phi + \gamma_{\mathcal{S}}^- \mathbf{B}(x, D) \Phi, \gamma_{\mathcal{S}}^+ \mathbf{B}(x, D) \Phi - \gamma_{\mathcal{S}}^- \mathbf{B}(x, D) \Phi \} \quad (1)$$

$$\mathcal{R}_{\mathcal{S}}(\mathbf{B}) : \mathbb{H}_{p, \text{loc}}^s(\overline{\mathbb{R}_{\mathcal{S}}^n}) \longrightarrow \mathbb{B}_{p, p}^{s - \frac{1}{p} - r}(\mathcal{S}) \otimes \widetilde{\mathbb{B}}_{p, p}^{s - \frac{1}{p} - r}(\mathcal{S}). \quad (2)$$

1. AN IMPLICIT EXTENSION FROM A HYPERSURFACE

For a pair of spaces we introduce the following shortcut: $\overset{2}{\mathbb{B}}_{p, p}^s(\mathcal{S}) := \mathbb{B}_{p, p}^s(\mathcal{S}) \otimes \widetilde{\mathbb{B}}_{p, p}^s(\mathcal{S})$. The notation $[s]^- \in \mathbb{Z}$ is used for the largest positive or negative integers less than s , i.e., $s - 1 \leq [s]^- < s$.

Theorem 1.1. *Let $\mathbf{A}(x, D)$ be a PDO of order $k \in \mathbb{N}_0$ and of normal type, $1 < p < \infty$, $k \in \mathbb{N}_0$, $s > 0$, $k \leq s + 1$ and denote $p_s := p$ if $s \neq 0, \pm 1, \dots$, $p_s < p$ if $s = 0, \pm 1, \dots$*

Further, let $\vec{\mathbf{B}}^{(k)}(x, D) := \{\mathbf{B}_0(x, D), \dots, \mathbf{B}_{k-1}(x, D)\}^{\top}$ be a Dirichlet system of boundary operators and $\{\varphi_j^{\pm}\}_{j=0}^{k-1}$ be vector functions such that

$$\Phi_j := (\varphi_j^+ + \varphi_j^-, \varphi_j^+ - \varphi_j^-) \in \overset{2}{\mathbb{B}}_{p, p}^{s-j}(\mathcal{S}), \quad \text{for all } j = 0, 1, \dots, k-1. \quad (3)$$

Then there exists a continuous linear operator

$$\mathcal{P}_{\mathbf{A}} : \overset{\otimes}{\underset{j=0}{k-1}} \overset{2}{\mathbb{B}}_{p, p}^{s-j}(\mathcal{S}) \rightarrow \mathbb{H}_{p_s, \text{loc}}^{s+1/p_s}(\mathbb{R}_{\mathcal{S}}^n) \quad (4)$$

such that $\gamma_{\mathcal{S}^{\pm}} \mathbf{B}_j \mathcal{P}_{\mathbf{A}} \Phi = \varphi_j^{\pm}$ for $j = 0, 1, \dots, k-1$ and $\mathbf{A} \mathcal{P}_{\mathbf{A}} \Phi \in \widetilde{\mathbb{H}}_{p_s, \text{loc}}^{s-k+1/p_s}(\mathbb{R}_{\mathcal{S}}^n)$, where $\Phi := \{\Phi_j\}_{j=0}^{k-1}$.

Proof. First we extend the pairs $(\varphi_j^+ + \varphi_j^-, \varphi_j^+ - \varphi_j^-)$ to the closed hypersurface \mathcal{C} , which contains \mathcal{S} . The second components are extended by 0. The obtained system is then extended into Ω^+ and Ω^- by coretraction for closed surfaces (see [5]) and, due to the conditions $\varphi_j^+ - \varphi_j^- = 0$ on the

complement surface $\mathcal{S}^c := \mathcal{C} \setminus \overline{\mathcal{S}}$, we prove that this particular extension is “continuous” across the complement surface \mathcal{S}^c . \square

2. PERTURBED POLY-HARMONIC EQUATIONS

The perturbed poly-harmonic (poly-Laplacian) operator

$$\Delta_\omega^m \varphi := \Delta^m \varphi + (-\omega^2)^m \varphi = (\operatorname{div} \nabla)^m \varphi + (-\omega^2)^m \varphi, \quad m = 1, 2, \dots, k > 0, \quad (5)$$

is elliptic, self-adjoint $(\Delta_\omega^m)^* = \Delta_\omega^m$ and is sign-definite in the following sense:

$$\begin{aligned} ((-1)^m \Delta_\omega^m \varphi, \varphi) &= \|\nabla^m \varphi\|_{L_2(\mathbb{R}^n)}^2 + \omega^{2m} \|\varphi\|_{L_2(\mathbb{R}^n)}^2 \geq \\ &\geq c \|\varphi\|_{\mathbb{H}^m(\mathbb{R}^n)}^2 \end{aligned} \quad (6)$$

for some $c > 0$. Then $\Delta_\omega^m : \mathbb{H}^m(\mathbb{R}^n) \rightarrow \mathbb{H}^{-m}(\mathbb{R}^n)$ has the trivial kernel $\operatorname{Ker} \Delta_\omega^m = \{0\}$ and cokernel $\operatorname{Coker} \Delta_\omega^m = \{0\}$.

If $\mathcal{K}_{\Delta_\omega^m}(x)$ denotes the fundamental solution of Δ_ω^m , it decays at infinity fast

$$x^\beta \partial^\alpha \mathcal{K}_{\Delta_\omega^m}(x) = \mathcal{O}\left(|x|^{|\alpha|+|\beta|} e^{-\omega|x|}\right) = \mathcal{O}(1) \quad \text{as } |x| \rightarrow \infty, \quad \forall \alpha, \beta \in \mathbb{N}^n. \quad (7)$$

Let

$$\mathbf{g}_j \in \mathbb{B}_{p,p}^{s-j-1/p}(\mathcal{C}), \quad j = 0, \dots, m-1, \quad f \in \widetilde{\mathbb{H}}_p^{s-2m}(\Omega), \quad 1 < p < \infty, \quad s \in \mathbb{R} \quad (8)$$

and look for a solution $\varphi \in \mathbb{H}_p^s(\Omega)$ of the boundary value problem

$$\begin{cases} (\Delta_\omega^m \varphi)(x) = f(x), & x \in \Omega, \\ (\gamma_{\mathcal{C}} \mathbf{D}_j(x, D)\varphi)(x) = \mathbf{g}_j(x), & x \in \mathcal{C}, \quad j = 0, \dots, m-1, \end{cases} \quad (9)$$

where

$$\mathbf{D}_0(x, D) := I, \quad \mathbf{D}_j(x, D) := \begin{cases} -\Delta^r & \text{if } j = 2r, \\ \partial_\nu \Delta^r & \text{if } j = 2r + 1, \end{cases} \quad (10)$$

$$j = 1, \dots, 2m-1$$

are the natural boundary operators from the Green formula below (11) and (12).

The proof of the next Lemma 2.1 is standard, based on the Gauss formula.

Lemma 2.1. *For the operator Δ_ω^m in the domain $\Omega \subset \mathbb{R}^n$ with the smooth boundary $\mathcal{C} := \partial\Omega$ the following I and II Green formulae are valid:*

$$(\Delta_\omega^m \varphi, \psi)_\Omega = \sum_{j=[m/2]}^{2m-1} ((\mathbf{D}_j(x, D)\varphi)^+, (\mathbf{C}_j(x, D)\psi)^+)_\mathcal{C} + \mathbb{A}(\varphi, \psi), \quad (11)$$

$$(\Delta_\omega^m \varphi, \psi)_\Omega - (\varphi, \Delta_\omega^m \psi)_\Omega = \sum_{j=0}^{2m-1} ((\mathbf{D}_j(x, D)\varphi)^+, (\mathbf{C}_j(x, D)\psi)^+)_\mathcal{C}. \quad (12)$$

for arbitrary $\varphi, \psi \in \mathbb{X}^{2m}(\Omega)$, where $\mathbb{A}(\varphi, \psi)$ is the bilinear form:

$$\mathbb{A}(\varphi, \psi) := \begin{cases} (\Delta^\ell \varphi, \Delta^\ell \psi)_\Omega + \omega^{2m}(\varphi, \psi)_\Omega, & m = 2\ell, \\ -(\nabla \Delta^\ell \varphi, \nabla \Delta^\ell \psi)_\Omega - \omega^{2m}(\varphi, \psi)_\Omega, & m = 2\ell + 1. \end{cases} \quad (13)$$

For a solution $\varphi \in \mathbb{X}_p^m(\Omega)$ of the equation $\Delta_\omega^m \varphi = f$ with $f \in \widetilde{\mathbb{X}}_p^{-m}(\Omega)$ the traces $\gamma_{\mathcal{C}} \Delta^{\ell+j} \varphi$ and $\gamma_{\mathcal{C}} \partial_\nu \Delta^{\ell+j} \varphi$, $j = 0, 1, \dots, [(m+1)/2]$ exist and

$$\gamma_{\mathcal{C}} \Delta^j \varphi \in \mathbb{B}_{p,p}^{m-2j-\frac{1}{p}}(\mathcal{C}), \quad \gamma_{\mathcal{C}} \partial_\nu \Delta^j \varphi \in \mathbb{B}_{p,p}^{m-2j-1-\frac{1}{p}-1}(\mathcal{C}). \quad (14)$$

Corollary 2.2. A solution $\varphi \in \mathbb{H}_p^m(\Omega)$ to the boundary value problem (8)-(9) is represented as follows:

$$\varphi(x) = (\mathbf{N}_\Omega f)(x) + \sum_{j=0}^{2m-1} \mathbf{V}_j(\mathbf{D}_j(x, D)\varphi)^+(x), \quad x \in \Omega, \quad (15)$$

where

$$(\mathbf{N}_\Omega \varphi)(x) := \int_{\Omega} \mathcal{K}_{\Delta_\omega^m}(x-y)\varphi(y) dy, \quad (16)$$

$$(\mathbf{V}_j \psi)(x) := \oint_{\mathcal{C}} (\mathbf{C}_j(x, D)\mathcal{K}_{\Delta_\omega^m})(x-x)\psi(x) d\sigma \quad (17)$$

are, respectively, the volume (Newton) potential and the layer potentials (cf. [1]).

Theorem 2.3. If a solution $\varphi \in \mathbb{H}^m(\Omega)$ to BVP (9)-(8) for $p = 2$ and $s = m$ exists, it is unique.

The Dirichlet problem (9)-(8) for $p = 2$ and $s = m$ has a unique solution $\varphi \in \mathbb{H}^m(\Omega)$ for an arbitrary right-hand side $\mathbf{g}_j \in \mathbb{H}^{m-j-\frac{1}{2}}$, $j = 0, \dots, m-1$ and for $f \in \widetilde{\mathbb{H}}^{-m}(\Omega)$.

Lemma 2.4. \mathbf{N}_Ω is a Ψ DO of order $-2m$ with a proper symbol, while \mathbf{V}_j is the potential operator of order $-j-1/p$ and maps the spaces

$$\begin{aligned} \mathbf{N}_\Omega &: \widetilde{\mathbb{H}}_p^\theta(\Omega) \longrightarrow \mathbb{H}_p^{\theta-2m}(\Omega), \\ \mathbf{V}_j &: \mathbb{B}_{p,p}^\theta(\mathcal{C}) \longrightarrow \mathbb{H}_p^{\theta+j+\frac{1}{p}}(\Omega) \cap \mathbb{S}(\Omega) \quad j = 0, 1, \dots, 2m-1 \end{aligned} \quad (18)$$

continuously for arbitrary $1 < p < \infty$ and $\theta \in \mathbb{R}$.

The following Plemelj formulae are valid:

$$\begin{aligned} [(\mathbf{D}_j \mathbf{V}_j \psi)]^\pm(x) &= \pm \frac{1}{2} \psi(x) + (\mathbf{V}_{jj} \psi)(x), \\ [(\mathbf{D}_j \mathbf{V}_k \psi)]^-(x) &= (\mathbf{V}_{jk} \psi)(x) = [(\mathbf{D}_j \mathbf{V}_k \psi)]^+(x), \\ x \in \mathcal{C}, \quad j, k &= 0, 1, \dots, 2m-1, \quad j \neq k, \quad \psi \in C(\mathcal{C}), \end{aligned} \quad (19)$$

where $\mathbf{V}_{jk}(x, D)$ is the direct value of the potential operator $\mathbf{D}_j(t, \partial)\mathbf{V}_k$ on the boundary \mathcal{C} . It represents a Ψ DO of order $\text{ord } \mathbf{V}_{jk}(x, D) = -2m + 1 + j + 2m - 1 - k = j - k$, $j, k = 0, 1, \dots, 2m-1$.

Proof. For the standard proofs we refer to [1, 3]. \square

Theorem 2.5. *BVP (9) has a solution for an arbitrary right-hand side f and data $\mathbf{g}_0, \dots, \mathbf{g}_{m-1}$ which satisfy the conditions (8), represented by the formula*

$$\varphi(x) = (\mathbf{N}_\Omega f)(x) + \sum_{j=0}^{m-1} (\mathbf{V}_j g_j)(x) + \sum_{j=m}^{2m-1} (\mathbf{V}_j \psi_j)(x), \quad x \in \Omega, \quad (20)$$

where the vector-function $\Psi := (\psi_0, \dots, \psi_{m-1})^\top \in \overset{m-1}{\otimes}_{j=0} \mathbb{B}_{p,p}^{s-m-j-1/p}(\mathcal{C})$ is a unique solution to the following boundary pseudodifferential system:

$$\mathbf{W}(x, D)\Psi(x) = \frac{1}{2}\mathbf{G}(x) - \vec{\mathbf{D}}^{(m)}(x, D)\mathbf{N}_\Omega f(x) - \mathbf{W}_0\mathbf{G}(x), \quad x \in \mathcal{C}, \quad (21)$$

$$\mathbf{G} := (\mathbf{g}_0, \dots, \mathbf{g}_{m-1})^\top \in \overset{m-1}{\otimes}_{j=0} \mathbb{B}_{p,p}^{s-j-1/p}(\mathcal{C}), \quad (22)$$

$$\vec{\mathbf{D}}^{(m)}(x, D) := (\mathbf{D}_0(x, D), \dots, \mathbf{D}_{m-1}(x, D))^\top$$

and \mathbf{W}, \mathbf{W}_0 are the matrix operators

$$\mathbf{W} := \begin{bmatrix} \mathbf{V}_{0m} & \cdots & \mathbf{V}_{0(2m-1)} \\ \mathbf{V}_{1m} & \cdots & \mathbf{V}_{1(2m-1)} \\ \cdots & \cdots & \cdots \\ \mathbf{V}_{(m-1)m} & \cdots & \mathbf{V}_{(m-1)(2m-1)} \end{bmatrix},$$

$$\mathbf{W}_0 := \begin{bmatrix} \mathbf{V}_{00} & \cdots & \mathbf{V}_{0(m-1)} \\ \mathbf{V}_{10} & \cdots & \mathbf{V}_{1(m-1)} \\ \cdots & \cdots & \cdots \\ \mathbf{V}_{(m-1)0} & \cdots & \mathbf{V}_{(m-1)(m-1)} \end{bmatrix}.$$

If $s \geq m$, any solution φ to BVP (9) is represented by formula (20) and the solution is unique.

Proof. With the help of the Green formulae (11) and (12), invoking Lax-Milgram lemma, the result is proved first for $p = 2$ (see similar proofs in [3, 4]).

Then, using the representation formulae (18), Plemelj formulae (19), we derive equivalent boundary Ψ DOs and prove their solvability for general p and s . \square

3. EXTENSION WITH POTENTIALS FROM A HYPERSURFACE

Based on Theorem 2.5, we can prove similar result for the BVP with general boundary conditions

$$\begin{cases} (\Delta_\omega^m \varphi)(x) = f(x), & x \in \Omega, \\ (\gamma_{\mathcal{C}} \vec{\mathbf{B}}^{(m)}(x, D)\varphi)(x) = \mathbf{G}(x), & x \in \mathcal{C}, \end{cases} \quad (23)$$

$$\mathbf{G} := (g_0, \dots, g_{m-1})^\top \in \overset{m-1}{\otimes}_{j=0} \mathbb{B}_{p,p}^{s-j-1/p}(\mathcal{C}),$$

where $\vec{\mathbf{B}}^{(m)}(\mathbf{B}_0(x, D), \dots, \mathbf{B}_{m-1}(x, D))^\top$ is a Dirichlet system of order m , ord $\mathbf{B}_j = j$, $j = 0, 1, \dots, m-1$, and conditions (8) hold. Let

$$\vec{\mathbf{B}}^{(m)}(x, D) = \mathbf{M}_{\mathbf{B}, \mathbf{D}}^{(m \times m)}(x, \mathcal{D}) \vec{\mathbf{D}}^{(m)}(x, D), \quad (24)$$

where $\vec{\mathbf{D}}^{(m)}(x, D)$ is defined in (21) and $\mathbf{M}_{\mathbf{B}, \mathbf{D}}^{(m \times m)}(x, \mathcal{D})$ is a lower triangular invertible matrix operator (called **admissible**).

Let us explain what is meant under the *generalized traces* $\gamma_{\mathcal{C}}^\pm \psi = \psi^\pm \in \mathbb{B}_{p,p}^{\theta - \frac{1}{p}}(\mathcal{C})$ for distributions $\psi \in \mathbb{H}_p^\theta(\Omega^\pm)$ when $-\infty < \theta \leq \frac{1}{p}$. Towards this end, in a vicinity $\Omega_{\mathcal{C}}^\pm \subset \Omega^\pm$ of the surface \mathcal{C} we consider a local coordinate system $x = (\mathcal{x}, t) \in \Omega_{\mathcal{C}}^\pm$, where $\mathcal{x} \in \mathcal{C}$ and $0 \leq \pm t < \varepsilon$ is the transverse variable, equal to the signed distance from the reference point x to the surface \mathcal{C} . Then we define

$$\psi^\pm := \gamma_{\mathcal{S}^\pm} \psi \quad \text{if} \quad \lim_{0 < \pm t \rightarrow 0} \|\psi(\cdot, t) - \psi^\pm(\cdot)\|_{\mathbb{B}_{p,p}^{\theta - \frac{1}{p}}(\mathcal{C})}. \quad (25)$$

Lemma 3.1. *The Plemelj formulae (19) are valid for $\psi \in \mathbb{B}_{p,p}^\theta(\mathcal{C})$ when $1 < p < \infty$ and $\theta \in \mathbb{R}$ are arbitrary.*

Theorem 3.2. *BVP (23) has a solution for arbitrary f and data \mathbf{G} , represented by the formula*

$$\varphi(x) = (\mathbf{N}_\Omega f)(x) + \sum_{j=0}^{m-1} (\mathbf{V}_j \tilde{g}_j)(x) + \sum_{j=m}^{2m-1} (\mathbf{V}_j \psi_{j-m})(x), \quad x \in \Omega. \quad (26)$$

Here the vector $\Psi := (\psi_0, \dots, \psi_{m-1})^\top \in \bigotimes_{j=0}^{m-1} \mathbb{B}_{p,p}^{s-m-j-1/p}(\mathcal{C})$ is a unique solution to the following boundary pseudodifferential system:

$$\mathbf{W}(x, D)\Psi(x) = \frac{1}{2} \tilde{\mathbf{G}}(x) - \vec{\mathbf{D}}^{(m)}(x, D) \mathbf{N}_\Omega f(x) - \mathbf{W}_0 \tilde{\mathbf{G}}(x), \quad x \in \mathcal{C}, \quad (27)$$

with the Ψ DO $\mathbf{W}(x, D)$ from (21)

$$\tilde{\mathbf{G}} := (\tilde{g}_0, \dots, \tilde{g}_{m-1})^\top = \left[\mathbf{M}_{\mathbf{B}, \mathbf{D}}^{m \times m}(x, \mathcal{D}) \right]^{-1} \mathbf{G} \in \bigotimes_{j=0}^{m-1} \mathbb{B}_{p,p}^{s-j-1/p}(\mathcal{C}). \quad (28)$$

If $s \geq m$, any solution φ to BVP (23) is represented by formula (26) and the solution is unique.

As we see, the space setting for the BVP (23) is also general and arbitrary $s \in \mathbb{R}$ can be taken, involving the Bessel potential spaces of distributions for $s < 0$.

Applying this result, we can prove the following result on the (explicit) continuation of the set of functions from a hypersurface.

Theorem 3.3. *Let $m \in \mathbb{N}_0$, $\{\varphi_j^\pm\}_{j=0}^{m-1}$ be two m -tuples of functions (distributions, if $s < 0$)*

$$\Phi_j := (\varphi_j^+ + \varphi_j^-, \varphi_j^+ - \varphi_j^-) \in \mathbb{B}_{p,p}^{s-j}(\mathcal{S}), \quad j = 0, 1, \dots, m-1, \quad (29)$$

$$1 < p < \infty, \quad s \in \mathbb{R}$$

and denote $p_s := p$ if $s \neq 0, \pm 1, \dots$, $p_s < p$ if $s = 0, \pm 1, \dots$

For a given Dirichlet system of boundary operators

$$\vec{\mathbf{B}}^{(m)}(x, D) := \{\mathbf{B}_0(x, D), \dots, \mathbf{B}_{m-1}(x, D)\}^\top \quad (30)$$

and the preassigned boundary data (29) there exists an operator

$$\mathcal{P} : \bigotimes_{j=0}^{m-1} \mathbb{B}_{p,p}^{s-j}(\mathcal{S}) \rightarrow \mathbb{H}_{p_s}^{s+1/p_s}(\mathbb{R}_S^n) \cap \mathbb{S}(\mathbb{R}_S^n) \quad (31)$$

which is a co-retraction $\gamma_{\mathcal{S}^\pm} \mathbf{B}_j \mathcal{P} \Phi = \varphi_j^\pm$, $j = 0, 1, \dots, m-1$ to the trace operator $\gamma_{\mathcal{C}}(\vec{\mathbf{B}}^{(m)}) := \{\gamma_{\mathcal{C}} \mathbf{B}_0(x, D), \dots, \gamma_{\mathcal{C}} \mathbf{B}_{m-1}(x, D)\}^\top$.

ACKNOWLEDGEMENT

The investigation was supported by the grant of the Georgian National Science Foundation GNSF/ST07/3-175.

REFERENCES

1. R. Duduchava, The Green formula and layer potentials. *Integral Equations and Operator Theory* **41** (2001), No. 2, 127–178.
2. R. Duduchava, Continuation of functions from hypersurfaces. *Complex Analysis and Differential Equations*, (to appear).
3. R. Duduchava, D. Natroshvili, E. Shargorodsky, Basic boundary value problems of thermoelasticity for anisotropic bodies with cuts. I-II, *Georgian Math. J.* **2** (1995), No. 2, 123–140.
4. R. Duduchava, F.-O. Speck, Pseudodifferential operators on compact manifolds with Lipschitz boundary. *Math. Nachr.* **160** (1993), 149–191.
5. H. Triebel, Interpolation theory, function spaces, differential operators. Second edition. *Johann Ambrosius Barth, Verlag, Heidelberg*, 1995.

Author's address:

A. Razmadze Mathematical Institute
 I. Javakhishvili Tbilisi State University
 2, University Str., Tbilisi 0186
 Georgia