

# Enchanted Topological Spaces

Yu-Lin Chou\*

## Abstract

This article introduces and studies in particular a new topological invariant — enchantedness; the enchanted spaces include every Euclidean space of dimension at least two. Via enchantedness one may gain some insight into Euclidean topologies, and local enchantedness in particular furnishes a new, simple, and intrinsic way to “topologically differentiate” topological 1-manifolds from higher-dimensional ones. A generic construction of enchanted spaces, with some additional nice properties, is given. This construction also induces a notion of multiple-point connectification, generalizing the existing, well-developed notion of one-point connectification, and a dual result pertaining to multiple-point connectification.

**MSC 2020:** 54D99; 54D05; 54B99; 54D35

**Keywords:** connectedness; enchanted spaces; Euclidean topologies; generalized Sierpiński spaces; local enchantedness; multiple-point connectification; topological manifolds

Like compactness, connectedness is not hereditary. However, compactness is at least closed hereditary.

Connectedness, on the other hand, seems not as fortunate; here we certainly have omitted the trivial cases such as singletons. Although an

---

\*freelance author, Hsinchu County, Taiwan (R.O.C.); email: [chou.y.l.edu@gmail.com](mailto:chou.y.l.edu@gmail.com).

abstract nontrivial characterization of connected subspaces is available (e.g. Theorem 6.1.7 in Engelking [2]), it is unclear whether there is a simple, easily recognizable property, “comparable” to closedness in the compactness case, such that every subset, with this property, of a connected space inherits connectedness; this is arguably an unsettled old “problem”.

We develop in this paper some fundamental properties of some new concepts, initially motivated by the problem, which, however, turn out to be worth studying independently of the initially motivating problem. From their fruits, these new topological “infrastructures”, revolving around the new topological invariant — enchantedness, seem to admit respectively an unexpected, significant position; and they also serve as tools that complement connectedness theory and that may contribute to constructing counterexamples.

Throughout, a topological space is assumed to possess a property if and only if a corresponding declaration is made.

To begin with, a preliminary observation is that one can easily construct topological spaces with rich subspace properties where there is some open proper subset, receiving a nontrivial subspace topology, such that every superset of this set is connected:

**Theorem 1.** *Let  $X$  be a topological space; let  $F$  be a finite nonempty set with  $X \cap F = \emptyset$ . If  $Y := X \cup F$ , then there exists some connected topology of  $Y$  such that i) every superset of  $F$  in  $Y$  is connected and ii) the set  $F$  is open in  $Y$ , and receives a nontrivial subspace topology from  $Y$  whenever  $F$  has at least two elements; moreover, the given space  $X$*

is embeddable into  $Y$  as a closed subspace.

*Proof.* Without loss of clarity, we present a proof slightly more general with respect to the purpose.

Represent  $F \equiv \{p_1, \dots, p_n\}$ , and let  $\mathcal{T}_X$  be the given topology of  $X$ .

Give  $Y$  the topology

$$\mathcal{T}_Y := \{\emptyset\} \cup \bigcup_{1 \leq i \leq n} \{G \cup \{p_1, \dots, p_i\} \mid G \in \mathcal{T}_X\}.$$

Then, since every nonempty open set in  $Y$  contains  $p_1$ , the topology  $\mathcal{T}_Y$  is connected.

Likewise, as every nonempty open set in any given subspace  $\{p_1, \dots, p_i\}$  of  $Y$  contains  $p_1$  by construction, it follows that each  $\{p_1, \dots, p_i\}$  is connected. Moreover, each  $\{p_1, \dots, p_i\}$  is dense in  $Y$ ; for, the only closed superset of  $\{p_1, \dots, p_i\}$  in  $Y$  is by construction the space  $Y$  itself.

Given any  $1 \leq i \leq n$ , fix any  $A \supset \{p_1, \dots, p_i\}$  in  $Y$ . Since  $\{p_1, \dots, p_i\}$  is connected and dense in  $Y$ , it follows that  $A$  is connected. This proves i).

Upon remarking that each set  $\{p_1, \dots, p_i\}$  is open in  $Y$  and, as a subspace of  $Y$ , receives the topology  $\{\emptyset\} \cup \{\{p_1, \dots, p_j\} \mid 1 \leq j \leq i\}$ , and upon referring to the definition of  $\mathcal{T}_Y$ , we have completed the proof.  $\square$

**Remark 1.** Theorem 1 evidently allows for nearly arbitrary choices of  $X$  and  $F$  to deliver the desired conclusion, and so the collection  $\{A \mid F \subset A \subset Y\} = \{S \cup F \mid S \subset X\}$  of sets in  $Y$  can thus be

in particular made nontrivial in a wild sense. Moreover, the set  $X$  is closed and nonopen in  $Y$ , and the given topology of  $X$  is precisely the subspace topology of  $X$  received from  $Y$ .

The setting in Theorem 1 is reminiscent of that of extension topologies, but is essentially different in that  $F$ , as a subspace of  $Y$ , receives a connected topology.

We add that, although the space  $Y$  in Theorem 1 is manifestly  $T_0$  whenever  $X$  is  $T_0$ , it cannot be  $T_1$  (and hence not Hausdorff) as long as  $F$  has at least two elements.  $\square$

The subspace topology that  $F$ , when it has at least two elements, receives in Theorem 1 does not play a main role in this paper; but, in view of its conceptual importance, it may deserve a name:

**Definition 1.** A topological space is called a *generalized Sierpiński space* if and only if it is a finite set  $\{p_1, \dots, p_n\}$  having at least two elements and receiving a topology of the form  $\{\emptyset\} \cup \bigcup_{1 \leq i \leq n} \{q_1, \dots, q_i\}$  with  $\{q_1, \dots, q_n\} = \{p_1, \dots, p_n\}$ .  $\square$

The position of a generalized Sierpiński space is seen in the following

**Proposition 2.** *i) Every generalized Sierpiński topology is nontrivial and nondiscrete, and is well-ordered by the inclusion relation. ii) Every set in any given generalized Sierpiński space is connected.*

*Proof.* The statement i) is evident.

For ii), fix a generalized Sierpiński space  $S \equiv \{p_1, \dots, p_n\}$ , and, without loss of generality, let  $G_i := \{p_1, \dots, p_i\}$  be a nonempty open set in the given topology of  $S$  for all  $1 \leq i \leq n$ .

If  $A \subset S$  is disconnected, choose some open sets  $G_i, G_j$  in  $S$  such that  $G_i \cap A$  and  $G_j \cap A$  are nonempty and form a partition of  $A$ . Since then  $G_i \neq G_j$ , assume that  $G_i \subsetneq G_j$  without loss of generality. It follows that  $A \subset G_j \setminus G_i$ ; for, if  $p \in A$  is also contained in  $G_i$ , then  $p \in G_i \cap G_j \cap A$ .

We in turn have  $A \subset \{p_{i+1}, \dots, p_j\}$ . If  $k$  is the smallest element of  $\{i+1, \dots, j\}$  with  $p_k \in A$ , then every nonempty set open in  $A$  contains  $p_k$ ; and so  $A$  receives a connected subspace topology, a contradiction.  $\square$

**Remark 2.** For the notion of a generalized Sierpiński space, requiring finiteness is not essential; with more notation, one can easily extend the notion to countable sets (with at least two elements) without losing Proposition 2.

However, for our purposes, considering finite sets suffices.  $\square$

**Convention.** By the Sierpiński space we will always mean the topological space  $(\{0, 1\}, \{\emptyset, \{0\}, \{0, 1\}\})$ .  $\square$

We exemplify Theorem 1 by constructing strictly increasing sequences of connected subspaces in some spaces with “increasing complexity” of their connected components and topological structures:

**Example 1.** i) Consider the discrete space  $X := \mathbb{N} := \omega \setminus \{\emptyset\}$  of positive integers, and, given any  $n \in \mathbb{N}$ , let  $F := \{1/(n+1), \dots, n/(n+1)\}$ . Upon giving  $Y := X \cup F$  the specific topology as in Theorem 1, it follows by Theorem 1 that each of the subspaces  $\{1, \dots, m\} \cup F$  of  $Y$  is connected.

ii) For each  $n \in \mathbb{N}$ , let  $K_n$  be the remaining closed set in  $[0, 1]$  obtained from the  $n$ th removal stage of the usual Cantor ternary operation, so that  $X := \bigcap_{n \in \mathbb{N}} K_n$  is the classical Cantor set. Then  $X$  is totally disconnected, nondiscrete, and nontrivial.

For each  $n \in \mathbb{N}$ , choose some  $x_n \in \bigcap_{m \geq n} K_m$ . If  $F$  is a finite set of positive prime numbers, then the subspaces  $\{x_1, \dots, x_n\} \cup F$  of  $Y := X \cup F$  topologized as in Theorem 1 are all connected by Theorem 1.

iii) As another choice of  $X$  as a nondiscrete, nontrivial, and totally disconnected space, we might consider an uncountable ordinal number taken as an ordinal space. Upon choosing a finite set  $F$  that does not meet  $X$ , e.g. the choice of  $F$  in the subexample i), the union of  $F$  and each of the initial segments included in  $X$  is by Theorem 1 connected as a subspace of  $Y := X \cup F$  topologized as in Theorem 1.

iv) We now consider a disconnected space  $X$  with infinitely many connected components each of which has continuum-many elements. Let  $X := \bigcup_{n \in \omega} ]n, n + 1[$  be a subspace of the Euclidean space  $\mathbb{R}$ . Then  $X$  is a desired choice. If  $F$  is a finite set of negative even integers, and if  $Y := X \cup F$  is topologized in the same way as in Theorem 1, then the subspaces  $(\bigcup_{n=0}^m ]n, n + 1[) \cup F$  of  $Y$  are all connected by Theorem 1. □

Thanks to Theorem 1, we may introduce a new class of topological spaces:

**Definition 2.** Let  $Y$  be a topological space; let  $E \subset Y$ . The space  $Y$  is

said to be *enchanted by  $E$*  if and only if i) the set  $E$  is a proper subset of  $Y$  and ii) every proper subset  $A$  of  $Y$  with  $A \supset E$  is connected. In this case, the set  $E$  is called an *enchancing set* in  $Y$ .

The space  $Y$  is said to be an *enchanted space* if and only if there is some enchancing set in  $Y$ .  $\square$

Thus every enchancing set in an enchanted space is connected.

**Remark 3.** In other words, Theorem 1 asserts (in particular) that, given any finite nonempty set  $F$ , one can always construct “many” connected spaces enchanted by  $F$  with  $F$  being an open nontrivial subspace; and so Theorem 1 indeed gives nontrivial examples of an enchanted space that is connected.

But there are natural and simple enchanted (and connected) spaces: The Sierpiński space  $\{0, 1\}$  (receiving the topology  $\{\emptyset, \{0\}, \{0, 1\}\}$ ) is enchanted by  $\{0\}$ . More generally, we might also consider the generalized Sierpiński spaces: By Proposition 2, every proper subset, empty or not, of any given generalized Sierpiński space  $S$  is an enchancing set in  $S$ ; in particular, every generalized Sierpiński space is an enchanted space.

In passing, every given indiscrete (i.e. trivial) space is enchanted by each of its proper subsets.  $\square$

**Remark 4.** It follows immediately from the definition of an enchancing set that, in any given enchanted space, every union of enchancing sets is an enchancing set provided that the union is a proper subset of the given space.

However, a union of enchanted subspaces, even with one point in common, is not necessarily an enchanted subspace: The interval  $]0, 2]$  as a subspace of the Euclidean space  $\mathbb{R}$  is enchanted by  $]0, 2[$ , and the interval  $[1, 3[$  as a subspace of  $\mathbb{R}$  is enchanted by  $]1, 3[$ ; but, since every connected subspace of  $]0, 3[$  evidently has some disconnected superset in  $]0, 3[$ , the interval  $]0, 3[ = ]0, 2] \cup [1, 3[$  is not an enchanted subspace of  $\mathbb{R}$ .

In addition, the notion of an enchanting set and that of an enchanted subspace, while they sometimes agree, are logically independent: In the subspace  $[0, 1]$  of the Euclidean space  $\mathbb{R}$ , the set  $]0, 1]$  is both an enchanting set in and an enchanted subspace of  $[0, 1]$ ; the set  $]0, 1[$  is an enchanting set in  $[0, 1]$  but not an enchanted subspace of  $[0, 1]$ ; and the singleton  $\{0\}$  is an enchanted subspace (by the empty set) of  $[0, 1]$  but not an enchanting set in  $[0, 1]$ .  $\square$

Enchantedness is in particular a topological invariant:

**Theorem 3.** *Every continuous bijective image of an enchanted space is enchanted.*

*Proof.* Let  $Y$  be an enchanted space; fix a topological space  $Z$  and a continuous bijection  $f : Y \rightarrow Z$ .

Choose by assumption some enchanting set  $E$  in  $Y$ ; then the  $f$ -image  $f^1(E)$  of  $E$  is from assumption a proper subset of  $Z$ . If  $A$  is a proper subset of  $Z$  that includes  $f^1(E)$ , then, since the  $f$ -preimage  $f^{-1}(A)$  of  $A$  is a proper subset of  $Y$  that includes  $E$ , the set  $f^{-1}(A)$  is connected from assumption. But then, as  $f$  is by assumption continuous, the set

$A = f^1 \circ f^{-1}(A)$  is connected.

Now that  $Z$  is enchanted by  $f^1(E)$ , the proof is complete.  $\square$

Enchantedness is not hereditary and does not carry over to Cartesian products:

**Proposition 4.** *i) A subspace of an enchanted space need not be enchanted. ii) A Cartesian product of enchanted spaces need not be enchanted.*

*Proof.* i) Take any topological space  $X$  that is not enchanted (e.g. any discrete space with at least three elements), and consider the space  $X \cup F$  enchanted by some chosen finite nonempty  $F$  that does not meet  $X$  in the way described in Theorem 1; but then  $X$  is by construction a subspace of  $X \cup F$ .

ii) The discrete space  $\{0, 1\}$  is enchanted by, e.g.  $\{0\}$ ; but the Cartesian product  $\{0, 1\} \times \{0, 1\}$ , also being a discrete space, is evidently not enchanted.  $\square$

Nevertheless, for enchantedness there is a relatively easily recognizable simple property such that every subset of an enchanted space with this property inherits enchantedness:

**Theorem 5.** *If a subset  $Z$  of an enchanted space  $Y$  properly includes some enchanting set  $E$  in  $Y$ , then  $Z$  is an enchanted subspace of  $Y$  with  $E$  being an enchanting set in  $Z$ .*

*Proof.* The enchanting set  $E$  in  $Y$  is from assumption a proper subset of  $Z$  and hence connected in  $Z$ . But every proper subset  $A$  of  $Z$  with

$A \supset E$  is from assumption connected in  $Y$ , and hence connected in  $Z$ ; so  $E$  enchants  $Z$ .  $\square$

Regarding enchanted subspaces, we also have

**Proposition 6.** *If  $A, Z$  are subsets of a topological space  $Y$  with  $A \supset Z$ , it follows that  $Z$  is an enchanted subspace of  $Y$  if and only if  $Z$  is an enchanted subspace of the subspace  $A$  of  $Y$ .*

*Proof.* From assumption, the subspace topology of  $Z$  received from  $Y$  coincides with that of  $Z$  received from  $A$ .  $\square$

**Theorem 7.** *Let  $Y$  be a topological space; let  $\Theta$  be a nonempty set; let  $Z_\theta$  be an enchanted subspace of  $Y$  for all  $\theta \in \Theta$ . If  $E_\theta$  is an enchanting set in  $Z_\theta$  for all  $\theta \in \Theta$ , and if  $\bigcap_{\theta \in \Theta} E_\theta \neq \emptyset$ , then  $\bigcup_{\theta \in \Theta} Z_\theta$  is an enchanted subspace of  $Y$  with  $\bigcup_{\theta \in \Theta} E_\theta$  being an enchanting set in  $\bigcup_{\theta \in \Theta} Z_\theta$ .*

*Proof.* From assumption, every  $E_\theta$  is also connected in  $Y$ , so that  $\bigcup_{\theta} E_\theta$  is connected in  $Y$  and hence in  $\bigcup_{\theta} Z_\theta$ .

Choose by assumption some point  $x$  of  $\bigcap_{\theta} E_\theta$ , and let  $A$  be a proper subset of  $\bigcup_{\theta} Z_\theta$  that includes  $\bigcup_{\theta} E_\theta$ . Then  $A = \bigcup_{\theta} (A \cap Z_\theta)$ , with each  $A \cap Z_\theta \supset E_\theta$  being connected in  $Z_\theta$ . In particular, the set  $A$  is a union of connected sets  $A \cap Z_\theta$  in  $Y$ . Since  $x \in \bigcap_{\theta} (A \cap Z_\theta)$ , it follows that  $A$  is connected in  $Y$  and hence in  $\bigcup_{\theta} Z_\theta$ .

We have proved that  $\bigcup_{\theta} E_\theta$  enchants  $\bigcup_{\theta} Z_\theta$ .  $\square$

Moreover, regarding Cartesian products we do at least have

**Proposition 8.** *Let  $\Theta$  be a nonempty set; let  $Y_\theta$  be a topological space for all  $\theta \in \Theta$ ; let  $\gamma \in \Theta$ . If  $E_\gamma$  is a proper subset of  $Y_\gamma$ , and if  $\prod_{\theta \in \Theta} Y_\theta$  is enchanted by the natural projective preimage  $\pi_\gamma^{-1}(E_\gamma)$  in  $\prod_{\theta \in \Theta} Y_\theta$ , then  $Y_\gamma$  is enchanted by  $E_\gamma$ .*

*Proof.* We might first remark that the proof of Theorem 3 also serves continuous surjections that carry some enchanting set to a proper subset. But then the natural projection  $\pi_\gamma : \prod_{\theta} Y_\theta \rightarrow Y_\gamma, (y_\theta)_\theta \mapsto y_\gamma$  is, from assumption, such a continuous surjection.  $\square$

It follows from Theorem 3 that the cardinal of the collection of all enchanting sets in an enchanted space is also a topological invariant:

**Corollary 9.** *Let  $Y, Z$  be enchanted spaces. If  $Y$  is homeomorphic to  $Z$ , then the collection of all the enchanting sets in  $Y$  is equinumerous to that of all the enchanting sets in  $Z$ , corresponding ones being homeomorphic.*

*Proof.* This follows immediately from the proof of Theorem 3 by choosing a homeomorphism  $f$  acting between  $Y$  and  $Z$ ; for then the  $f$ -image map  $f^1$  is a desired choice.  $\square$

**Remark 5.** We now have a new view on the topological differences between compact intervals and a bounded half-closed interval in the Euclidean space  $\mathbb{R}$ : The compact interval  $[-1, 1]$ , as an enchanted subspace of  $\mathbb{R}$ , has more than one enchanting sets, e.g.  $[-1, 1[$  or  $] -1, 1]$ ; but the bounded half-closed interval  $] -1, 1]$ , as an enchanted subspace of  $\mathbb{R}$ , has exactly one enchanting set, e.g.  $] -1, 1[$  (since every connected

proper subset of  $] -1, 1]$  other than  $] -1, 1[$  evidently has some disconnected superset in  $] -1, 1]$ ; and so Corollary 9 gives a new proof that a compact interval is never homeomorphic to a bounded half-closed interval.

Moreover, we obtain a new view on the extended real line  $\overline{\mathbb{R}}$  topologized as usual: Since  $[-1, 1]$  as a subspace of  $\mathbb{R}$  evidently has exactly three enchanting sets, and since  $[-1, 1]$  is homeomorphic to  $\overline{\mathbb{R}}$ , it holds by Theorem 3 and Corollary 9 that  $\overline{\mathbb{R}}$  has exactly three enchanting sets.  $\square$

Although an enchanted space is sometimes also a connected space, the two notions are logically independent:

**Proposition 10.** *i) There exists some enchanted space that is disconnected. ii) There exists some connected space that is nonenchanted.*

*Proof.* For a quick justification of i), we might consider once more the discrete space  $\{0, 1\}$ .

There are also less trivial examples; we give a general construction of enchanted disconnected spaces with at least three elements: Let  $(X, \mathcal{T}_X)$  be a connected space with cardinality at least two; let  $p \notin X$ . Topologize the set  $Y := X \cup \{p\}$  by the topology  $\{\emptyset\} \cup \mathcal{T}_X \cup \{G \cup \{p\} \mid G \in \mathcal{T}_X\}$ . Then, since  $X$  and  $\{p\}$  are clopen in  $Y$ , the space  $Y$  is disconnected.

Moreover, the given topology of  $X$  coincides with the subspace topology of  $X$  in  $Y$ ; and so the set  $X$  enchants  $Y$ .

On the other hand, for ii), the Euclidean space  $\mathbb{R}$  cannot be en-

chanted; for, every connected proper subset of  $\mathbb{R}$ , which has to be a singleton or an interval, evidently has some disconnected superset in  $\mathbb{R}$ . This completes the proof.  $\square$

**Remark 6.** Thus, from the proof of the statement i) in Proposition 10, an enchanting set in an enchanted space can also be closed.  $\square$

In many cases, no connected component of a topological space can be an enchanting set:

**Proposition 11.** *If  $Y$  is a topological space with every connected component having at least two elements, then no connected component of  $Y$  is an enchanting set in  $Y$ .*

*Proof.* If  $Y$  is connected, then the only connected component, being  $Y$  itself, is not a proper subset of  $Y$ .

Suppose  $Y$  is disconnected. If  $C$  is a connected component of  $Y$ , then, due to the maximality of connected components, every set in  $Y$  properly including  $C$  is disconnected. But, from assumption, we can choose some element  $x$  of another connected component of  $Y$  such that the set  $C \cup \{x\}$ , which properly includes  $C$ , is a proper subset of  $Y$ ; and therefore  $C$  cannot enchant  $Y$ .  $\square$

There is a simple sufficient condition for a topological space to be an enchanted connected space:

**Proposition 12.** *If  $Y$  is a topological space, and if  $E$  is a connected proper subset of  $Y$  that is dense in  $Y$ , then  $Y$  is connected and is enchanted by  $E$ .*

*Proof.* Since  $Y = \text{cl}(E)$  by assumption, if  $A \subset Y$  includes  $E$  then  $A$  is connected.  $\square$

**Example 2.** Let  $E$  be the graph of the function  $x \mapsto \sin(1/x)$  on  $]0, 1]$ , considered as a subspace of the Euclidean space  $\mathbb{R}^2$ , so that  $Y := E \cup \{(0, z) \mid -1 \leq z \leq 1\} = \text{cl}(E)$  as a subspace of the Euclidean space  $\mathbb{R}^2$  is the “topologist’s sine curve”. Thus  $Y$  is by Proposition 12 in particular an enchanted space with  $E$  being an enchanting set.  $\square$

**Remark 7.** The converse of Proposition 12 is not true: The Sierpiński space  $\{0, 1\}$  can be enchanted by  $\{1\}$ , but  $\{1\}$  is closed therein.

Apart from Example 2, the assumption of Proposition 12 is by Theorem 1 far from vacuous.  $\square$

**Corollary 13.** *The closure of every given nonclosed connected set in a topological space  $Y$  is an enchanted subspace of  $Y$ .*

*Proof.* If  $Q \subset Y$  is nonclosed and connected, then  $Q$  is a proper subset of the subspace  $\text{cl}(Q)$  of  $Y$  and is connected in  $\text{cl}(Q)$ ; the desired conclusion follows from Proposition 12.  $\square$

For our purposes, we adopt the definition that an interval in the real field  $\mathbb{R}$  is precisely a subset  $I$  of  $\mathbb{R}$  such that if  $c \in \mathbb{R}$  and if  $a < c < b$  for some  $a, b \in I$  then  $c \in I$ . Then there is a characterization for (nonempty) enchanted connected subspaces of the Euclidean space  $\mathbb{R}$ :

**Theorem 14.** *A nonempty enchanted connected subspace of the Euclidean space  $\mathbb{R}$  is precisely a subspace of  $\mathbb{R}$  that is an interval nonopen in  $\mathbb{R}$ .*

*Proof.* A singleton in  $\mathbb{R}$  is enchanted by the empty set; and a compact interval  $[a, b]$  in  $\mathbb{R}$  is enchanted by, for example, the subinterval  $[a, b[$ . Every half-closed interval  $[a, b[$  in  $\mathbb{R}$ , with  $a, b \in \mathbb{R}$ , is evidently enchanted by the subinterval  $]a, b[$ ; the same observation shows that every half-closed interval  $]a, b]$  in  $\mathbb{R}$  is an enchanted subspace, and that all closed rays  $[a, +\infty[$  or  $] -\infty, a]$  in  $\mathbb{R}$  are enchanted subspaces. This proves one of the assertions.

Suppose the converse is false, and choose one such nonempty enchanted connected subspace  $Y$  of  $\mathbb{R}$ . Then  $Y$  is an interval in  $\mathbb{R}$  that is open in  $\mathbb{R}$ . Since every connected proper subset of a nonempty interval open in  $\mathbb{R}$  evidently has, by the properties of the real numbers, some disconnected superset being a proper subset of the interval, it follows that  $Y$  is not enchanted; this contradiction completes the proof.  $\square$

**Remark 8.** From Theorems 14 and 3, although less information is really needed, a new view on the fact that every half-closed interval and every closed ray in the Euclidean space  $\mathbb{R}$  are not homeomorphic to  $\mathbb{R}$  is readily available.  $\square$

For (nonempty) open sets in the Euclidean space  $\mathbb{R}$ , we have

**Proposition 15.** *Every nonempty open subset of the Euclidean space  $\mathbb{R}$  is nonenchanted.*

*Proof.* Let  $G \subset \mathbb{R}$  be open. If  $G$  is connected, then by Theorem 14 we are through. If  $G$  is disconnected, fix a connected proper subset  $Q$  of  $G$ . Since every open set in  $\mathbb{R}$  is a disjoint union of intervals open in  $\mathbb{R}$ ,

choose some partition  $\mathcal{S}$  of  $G$  by intervals open in  $\mathbb{R}$ . Then  $\mathcal{S}$  has at least two elements, and we can choose some  $I \in \mathcal{S}$  such that  $Q \subset I$ .

But, upon choosing some point  $x$  of some  $J \in \mathcal{S}$  with  $J \neq I$ , we see that  $Q \cup \{x\}$  is a disconnected superset of  $Q$  that is a proper subset of  $G$ .  $\square$

Even though the Euclidean space  $\mathbb{R}$  is not enchanted, for higher-dimensional Euclidean spaces we have

**Theorem 16.** *Every Euclidean space  $\mathbb{R}^n$  with  $n \geq 2$  is enchanted.*

*Proof.* If  $x \in \mathbb{R}^n$ , then the set  $E := \mathbb{R}^n \setminus \{x\}$ , being a co-countable set in  $\mathbb{R}^n$ , is a connected subspace of  $\mathbb{R}^n$ ; so  $E$  enchants  $\mathbb{R}^n$ .  $\square$

**Remark 9.** The proof of Theorem 16 implies also that every Euclidean space of dimension at least two has at least continuum-many enchanting sets.

In addition, a new aspect of the fact that every Euclidean space  $\mathbb{R}^n$  with  $n \geq 2$  is not homeomorphic to  $\mathbb{R}$  is obtained immediately from Proposition 15, Theorem 16, and Theorem 3.  $\square$

Enchanted open subspaces are “abundant” in the Euclidean spaces  $\mathbb{R}^n$  with  $n \geq 2$ :

**Theorem 17.** *Let  $n \geq 2$  be an integer; let  $\Theta$  be a nonempty set. If  $\{Z_\theta\}_{\theta \in \Theta}$  is a collection of subsets of  $\mathbb{R}^n$  homeomorphic to  $\mathbb{R}^n$  with nonempty intersection, then  $\cup_{\theta \in \Theta} Z_\theta$  is an enchanted connected open subspace of  $\mathbb{R}^n$ .*

*Proof.* That the union  $\cup_{\theta} Z_{\theta}$  is connected is evident from assumption.

By invariance of domain, every  $Z_{\theta}$  is open in  $\mathbb{R}^n$ ; so  $\cup_{\theta} Z_{\theta}$  is open in  $\mathbb{R}^n$ .

To show the enchantedness of  $\cup_{\theta} Z_{\theta}$ , choose by assumption some  $x \in \cap_{\theta} Z_{\theta}$ . For every  $\theta$ , choose some  $z_{\theta} \in Z_{\theta}$  such that  $z_{\theta} \neq x$ . Then, from assumption, the set  $Z_{\theta} \setminus \{z_{\theta}\}$  enchants  $Z_{\theta}$  for all  $\theta$ . Since  $x \in \cap_{\theta} (Z_{\theta} \setminus \{z_{\theta}\})$ , it follows by Theorem 7 that  $\cup_{\theta} Z_{\theta}$  is enchanted. This completes the proof.  $\square$

**Example 3.** Thus the union of a collection of open balls in the Euclidean space  $\mathbb{R}^2$  containing the origin is an enchanted subspace of  $\mathbb{R}^2$ .  $\square$

It would be natural to consider the following “local version” of enchantedness:

**Definition 3.** Let  $Y$  be a topological space. The space  $Y$  is called *locally enchanted* if and only if there exists some basis of  $Y$  consisting of enchanted subspaces.  $\square$

It is straightforward to show that local enchantedness is also a topological invariant:

**Theorem 18.** *Every homeomorphic copy of a locally enchanted space is locally enchanted.*

*Proof.* Let  $Y$  be a locally enchanted space; let  $Z$  be a topological space; let  $f : Y \rightarrow Z$  be a homeomorphism.

If  $G$  is open in  $Z$ , choose by assumption some collection  $\mathcal{V}$  of enchanted open subspaces of  $Y$  such that  $\cup \mathcal{V} = f^{-1}(G)$ . Then, since  $f^1(V)$ , which is open in  $Z$  for all  $V \in \mathcal{V}$  from assumption, is for all  $V \in \mathcal{V}$  enchanted as a subspace of  $Z$  by Theorem 3, and since  $G = \cup_{V \in \mathcal{V}} f^1(V)$ , we are through.  $\square$

Like enchantedness, local enchantedness is not hereditary and does not carry over to Cartesian products:

**Proposition 19.** *i) A subspace of a locally enchanted space need not be locally enchanted. ii) A Cartesian product of locally enchanted spaces need not be locally enchanted.*

*Proof.* i) From Theorem 16, every open ball in the Euclidean space  $\mathbb{R}^2$ , being homeomorphic to  $\mathbb{R}^2$ , is an enchanted open subspace of  $\mathbb{R}^2$ ; so  $\mathbb{R}^2$  is locally enchanted.

However, the Euclidean space  $\mathbb{R}$  cannot be locally enchanted; indeed, every nonempty enchanted subspace of  $\mathbb{R}$  is not open by Proposition 15. But then, from Theorem 18, the subspace  $\mathbb{R} \times \{(0, 0)\}$  of  $\mathbb{R}^2$  is not locally enchanted.

ii) Consider an infinite collection of discrete spaces  $Y_\theta$ , each of which has at least two elements. Then no set in  $\prod_\theta Y_\theta$  with at most two elements is open in  $\prod_\theta Y_\theta$ .

Fix any  $\theta$  and any nonempty open  $G_\theta$  in  $Y_\theta$ . If  $V \subset \pi_\theta^{-1}(G_\theta)$  is an enchanted subspace of  $\prod_\theta Y_\theta$ , then, since  $\prod_\theta Y_\theta$  is totally disconnected, the set  $V$  has at most two elements; but then  $V$  is not open in  $\prod_\theta Y_\theta$ . Thus  $\prod_\theta Y_\theta$  is not locally enchanted.  $\square$

But local enchantedness is open hereditary:

**Theorem 20.** *Every open set in a locally enchanted space  $Y$  is a locally enchanted subspace of  $Y$ .*

*Proof.* Fix any open set  $G$  in  $Y$ . Since every open set in  $G$  is from assumption open in  $Y$ , every open set in  $G$  is a union of enchanted open subspaces of  $Y$ .

Fix any nonempty  $V \subset G$  such that  $V$  is open in  $G$ . Then for every point of  $V$  we can choose some enchanted open subspace  $W$  of  $Y$  such that  $W$  contains the point and  $W \subset V$ , and  $W$  is also open in  $G$ .

But since  $W \subset G$ , the set  $W$  is also an enchanted subspace of  $G$  by Proposition 6. The proof is complete.  $\square$

For reference, given any  $n \in \mathbb{N}$ , by a topological  $n$ -manifold we mean precisely a (not necessarily second-countable or Hausdorff) locally Euclidean  $n$ -space, i.e. a topological space where every point has some neighborhood homeomorphic to some open subset of  $\mathbb{R}^n$ . To justify the introduction and evince the position of local enchantedness, we have the following results:

**Theorem 21.** *i) Every topological  $n$ -manifold with  $n \geq 2$  is locally enchanted. ii) Every topological 1-manifold is not locally enchanted.*

*Proof.* Let  $n \in \mathbb{N}$ , and fix a topological  $n$ -manifold  $M$ .

To prove i), suppose  $n \geq 2$ . If  $G$  is open in  $M$ , and if  $x \in G$ , then we can choose some coordinate neighborhood  $V$  of  $x$ , and choose in turn a neighborhood  $W$  of  $x$  that is included in  $V \cap G$  and homeomorphic to

$\mathbb{R}^n$ . Since  $W$  is then by Theorems 3 and 16 also enchanted, the desired conclusion follows.

For ii), suppose  $n = 1$ , and assume that  $M$ , now being a topological 1-manifold, is locally enchanted. Fix a point  $x$  of  $M$ , and choose some chart  $(G, \varphi)$  of  $M$  about  $x$ . And we can further choose by assumption some enchanted neighborhood  $V$  of  $x$  such that  $V \subset G$ .

But then  $\varphi^1(V)$  is by Theorem 3 an enchanted open subspace of  $\mathbb{R}$ , which contradicts Proposition 15. This completes the proof.  $\square$

**Proposition 22.** *i) There exists a topological space that is enchanted and locally enchanted. ii) There exists a topological space that is not enchanted nor locally enchanted. iii) There exists an enchanted space that is not locally enchanted. iv) There exists a locally enchanted space that is not enchanted.*

*Proof.* i) The Sierpiński space  $\{0, 1\}$  is enchanted. But the only choice of basis of this space is its topology; and both  $\{0\}$  and  $\{0, 1\}$  are enchanted open subspaces. In view of Theorems 16 and 21, one may also consider any Euclidean space  $\mathbb{R}^n$  with  $n \geq 2$ .

ii) The Euclidean space  $\mathbb{R}$  is (by Proposition 15, for concreteness) not enchanted, and is (by Theorem 21, for concreteness) not locally enchanted.

iii) Choose a topological space  $X$  that is connected and that is not locally enchanted (e.g. the Euclidean space  $\mathbb{R}$ ), and take a point  $p \notin X$ . Topologize the set  $Y := X \cup \{p\}$  in the same way as in the proof of the first statement of Proposition 10, so that  $Y$  is an enchanted space.

Now, since  $X$  is open in  $Y$ , and since a subset of  $X$  open in  $Y$  is precisely a subset of  $X$  open in the given topology of  $X$ , for  $Y$  to be locally enchanted requires  $X$  to be locally enchanted; this establishes the desired conclusion.

iv) Every discrete space with cardinality at least three is not enchanted; but, since every singleton in such a space is an enchanted open subspace of the space, the basis of the space, consisting of all the singletons in the space, is a desired choice.  $\square$

**Remark 10.** Evidently from the proof of Theorem 21, Theorem 21 holds for locally Euclidean  $n$ -spaces: Every locally Euclidean  $n$ -space with  $n \geq 2$  (resp.  $n = 1$ ) is locally enchanted (resp. not locally enchanted). In particular, from Theorems 21 and 18, we have obtained a new and simple proof that every topological  $n$ -manifold with  $n \geq 2$  is not homeomorphic to any given topological 1-manifold.

As summarized in the proof of Proposition 22, it holds that i) every Euclidean  $n$ -space with  $n \geq 2$  is both enchanted and locally enchanted and ii) the Euclidean 1-space is not enchanted nor locally enchanted. We thus have obtained a new perspective regarding the intrinsic topological differences between the Euclidean 1-space and the higher-dimensional Euclidean spaces.

Last but not least, Theorem 21 also implies that not every locally connected space is locally enchanted: We might consider the Euclidean space  $\mathbb{R}$ .  $\square$

There is an interesting result admitting a very simple proof:

**Theorem 23.** *No open set in a topological 1-manifold is homeomorphic to any given open set in a locally enchanted space.*

*Proof.* Let  $M$  be a topological 1-manifold; let  $Y$  be a locally enchanted space. If a nonempty open set  $G$  in  $M$  is homeomorphic to a nonempty open set  $W$  in  $Y$  via a homeomorphism  $f : G \rightarrow W$ , choose in  $M$  some coordinate neighborhood  $U$  of a point of  $G$ . Then  $G \cap U$  is homeomorphic to  $f^1(G \cap U)$ .

Since  $f^1(G \cap U) \subset W$  is open in  $Y$ , we can choose by assumption some nonempty enchanted open subspace  $V$  of  $Y$  with  $V \subset f^1(G \cap U)$ , and so, since  $G \cap U$  is by assumption also homeomorphic to some open subset of  $\mathbb{R}$ , the set  $f^{-1}(V) \subset G \cap U$  is homeomorphic to some open subset of  $\mathbb{R}$ .

But  $f^{-1}(V)$ , being by Theorem 3 (and Proposition 6, for concreteness) a nonempty enchanted open subspace of  $M$ , cannot be homeomorphic to any given open subset of  $\mathbb{R}$  by Theorem 3 and Proposition 15; this contradiction establishes the proof.  $\square$

For reference, a topological space is said to be locally homeomorphic to a topological space if and only if the former space has some open cover with each element being (topologically) embeddable into the latter space as an open subset; thus for every homeomorphic pair of topological spaces  $Y, Z$  it holds that  $Y$  (resp.  $Z$ ) is locally homeomorphic to  $Z$  (resp.  $Y$ ). Then we have

**Corollary 24.** *No topological 1-manifold is locally homeomorphic to any given locally enchanted space; no locally enchanted space is locally*

*homeomorphic to any given topological 1-manifold.*

*Proof.* This follows plainly from Theorem 23. □

**Remark 11.** Since every topological  $n$ -manifold with  $n \geq 2$  is by Theorem 21 locally enchanted, Theorem 23 offers a new intrinsic angle to further topologically differentiate topological 1-manifolds from higher-dimensional ones. □

We draw some additional observations. The construction introduced in the proof of Theorem 1 is motivated by the special one-point connectification given in Chou [1], which admits other independent focal points; and one-point connectifications enjoy a substantial history and rich developments in topology; one might refer to Koushesh [3], for instance, for some aspects of both the involved technical and nontechnical accounts.

Theorem 1 also inspires the following notion, which generalizes that of one-point connectification:

**Definition 4.** Let  $X, Y$  be topological spaces. The space  $Y$  is called a *multiple-point connectification* of  $X$  if and only if  $Y$  is a connected space including  $X$  as a subspace and  $Y \setminus X$  is a finite nonsingleton nonempty set. □

Here the relaxation of denseness, which is usually required in the context of space extension, in Definition 4 is not always standard.

Theorem 1 immediately admits a dual result regarding multiple-point connectifications:

**Corollary 25.** *Let  $X$  be a topological space; let  $\infty$  be a finite non-singleton nonempty set that does not meet  $X$ . Then there exists some multiple-point connectification  $X_\infty$  of  $X$ , with  $X_\infty \setminus X = \infty$ , such that*

- i) the space  $X_\infty$  is enchanted by  $\infty$  and*
- ii) the set  $\infty$  is a generalized Sierpiński space as a subspace of  $X_\infty$ .*

*Proof.* Upon choosing the specific topology for  $X_\infty$  as in Theorem 1, the desired conclusion follows plainly from Theorem 1.  $\square$

**Remark 12.** Although it is not a focus of the present work, we wish to point out that the topology introduced in Theorem 1, apart from its simplicity, would be manageable and convenient in some ways. For example, some important properties such as being second-countable or being an Alexandroff space can be easily seen to be transferable.

Moreover, we indicate once more that the topologies considered in Theorem 1 can be useful in constructing counterexamples in many contexts.  $\square$

## References

- [1] Y.-L. Chou, A special one-point connectification that characterizes  $T_0$  spaces, *Bulletin of the International Mathematical Virtual Institute* **12** (2022), 237–241.
- [2] R. Engelking, *General Topology*, revised and completed edition, Heldermann Verlag, 1989.

- [3] M.R. Koushesh, One-point connectifications, *Journal of the Australian Mathematical Society* **99** (2015), 76–84.