

ON AFFINE DIAMETERS OF FINITE SUBSETS OF THE PLANE

ALEXANDER KHARAZISHVILI

Abstract. It is shown that the total number $af(Z)$ of affine diameters of the set Z of all vertices of a convex n -gon in the plane $\mathbf{R}^2$ does not exceed $3n/2$. Also, it is established that if Z is an arbitrary n -point subset of $\mathbf{R}^2$, then $af(Z) \leq n^2/4 + 2$. Both these estimates are precise.

Let $\mathbf{R}^m$ denote the m -dimensional Euclidean space and let Z be a subset of $\mathbf{R}^m$ with $\text{card}(Z) \geq 2$.

By the standard definition, an affine diameter of Z is any line segment $[x, y]$ such that $\{x, y\} \subset Z$ and there exist two distinct parallel supporting hyperplanes G_x and G_y of Z passing through the points x and y respectively (see, e.g., [1, 3–5, 9]).

It follows from this definition that every affine diameter of Z is simultaneously an affine diameter of the convex hull of Z , denoted usually by $\text{conv}(Z)$ (the converse assertion fails to be true in general).

Any ordinary diameter of Z (with respect to the standard metric of $\mathbf{R}^m$) is an affine diameter of Z , but not conversely. If Z is a finite subset of the plane $\mathbf{R}^2$ with $\text{card}(Z) = n$, then the total number of ordinary diameters of Z does not exceed n , as a simple inductive argument shows. If $Z \subset \mathbf{R}^3$ and $\text{card}(Z) = n$, where $n \geq 2$, then the total number of ordinary diameters of Z does not exceed $2(n - 1)$ (see, for instance, [4, 6]). Various properties of affine diameters of point sets in $\mathbf{R}^m$ are considered in [1–6] and [8]. In this connection, see especially [9].

The goal of this short communication is to prove that if Z is the set of all vertices of a convex n -gon P in $\mathbf{R}^2$, then the total number of affine diameters of Z does not exceed $3n/2$. This estimate is precise, because gives the equality for the set of all vertices of a regular convex $2n$ -gon Q in $\mathbf{R}^2$ (and for the set of all vertices of the image of Q under a non-degenerate affine transformation of $\mathbf{R}^2$).

Throughout this note the symbol $Af(Z)$ will stand for the family of all affine diameters of a subset Z of the space $\mathbf{R}^m$, where $m \geq 2$. Also, we stipulate $af(Z) = \text{card}(Af(Z))$.

In the sequel, we need several auxiliary statements.

Lemma 1. *Let Z be a subset of $\mathbf{R}^m$ and let $\{Z_1, Z_2\}$ be a partition of Z . Denote by $\mathcal{A}_1$ the family of all those affine diameters of Z which are incident with at least one point from Z_1 .*

Then the inclusion $Af(Z) \subset \mathcal{A}_1 \cup Af(Z_2)$ takes place (in general, this inclusion is not reducible to the equality).

Lemma 2. *Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$, let $z \in Z$ and let z_1 and z_2 be two points of Z such that the line segments $[z, z_1]$ and $[z, z_2]$ are affine diameters of Z .*

Then any diagonal of P incident to z and lying in the convex angle formed by $[z, z_1]$ and $[z, z_2]$ is also an affine diameter of Z .

According to Lemma 2, if z is any vertex of a convex n -gon P , then there are some (maximally possible) successive vertices $z_1, z_2, \dots, z_k$ of P such that all line segments connecting these vertices with z turn out to be affine diameters of Z . Obviously, these successive vertices determine the corresponding successive sides

$$[z_1, z_2], [z_2, z_3], \dots, [z_{k-1}, z_k]$$

of P . In the sequel, we shall say that the above-mentioned successive sides constitute in P the affine support of z .

Lemma 3. *Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$, let z_1, z_2, z_3, z_4 be four distinct points from Z satisfying the following two conditions:*

2020 *Mathematics Subject Classification.* 52A10, 52B55.

Key words and phrases. Affine diameter; Convex polygon; Convex polyhedron.

- (a) the enumeration of z_1, z_2, z_3, z_4 is such that these points become successive vertices of the quadrangle $\text{conv}(z_1, z_2, z_3, z_4)$;
 (b) both line segments $[z_1, z_4]$ and $[z_2, z_3]$ are affine diameters of Z .
 Then $[z_1, z_2]$ and $[z_3, z_4]$ are two parallel sides of P .

Lemma 4. Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$ and let a point $z \in Z$ be such that there are at least five distinct affine diameters

$$[z, z_1], [z, z_2], [z, z_3], [z, z_4], [z, z_5]$$

of Z incident with z , where z_1, z_2, z_3, z_4, z_5 are some successive vertices of P .

Then $[z, z_3]$ is a unique affine diameter of Z incident with z_3 .

In connection with Lemma 4 it makes sense to notice that, for any natural number $n \geq 2$, there exists a convexly independent subset Z of $\mathbf{R}^2$ with $\text{card}(Z) = 2n + 1$, having the following properties:

- (*) each point of Z is incident with at least two affine diameters of Z ;
- (**) there is a point of Z incident with exactly four affine diameters of Z .

Lemma 5. Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$ and let one of the sides of P be an affine diameter of P .

Then the disjunction of these two assertions holds true:

- (1) P is a parallelogram;
- (2) the total number of affine diameters of Z is strictly less than $3n/2$.

Actually, in Lemma 5 the total number of affine diameters of Z does not exceed $n + 2$. Obviously, if $n > 4$, then $n + 2 < 3n/2$.

Lemma 6. Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$ and let a point $z \in Z$ be such that there are at least four distinct affine diameters $[z, z_1], [z, z_2], [z, z_3], [z, z_4]$ of Z , where z_1, z_2, z_3, z_4 are some successive vertices of P .

Then, for any vertex $z' \in Z$ distinct from z, z_1, z_2, z_3, z_4 , the side $[z_2, z_3]$ does not belong to the affine support of z' .

Lemma 7. Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$ and let n_k denote the number of all those vertices of P , each of which is incident with exactly k many affine diameters of Z .

Then the following relations are fulfilled:

- (1) $n_1 + 2n_2 + 3n_3 + \dots = 2af(Z)$;
- (2) $n_2 + 2n_3 + 3n_4 + \dots \leq 2n$.

Theorem 1. Let Z be the set of all vertices of a convex n -gon P in $\mathbf{R}^2$.

Then $af(Z) \leq 3n/2$. In addition, the equality $af(Z) = 3n/2$ is valid if and only if there exists a partition of the set of all sides of P into two-element subsets such that the sides of any member of the partition are parallel.

Remark 1. Consider any finite non-collinear convexly independent point set Z in $\mathbf{R}^2$ with $\text{card}(Z) = n$. It is easy to see that $af(Z) \geq n/2$. Therefore, in view of Theorem 1, the number $af(Z)$ is always of linear order with respect to $\text{card}(Z)$.

Remark 2. A substantially different situation can be observed for a non-coplanar convexly independent set $Z \subset \mathbf{R}^3$ with $\text{card}(Z) = n$. In this case, we obviously have

$$n/2 \leq af(Z) \leq n(n-1)/2.$$

Note also that:

- (i) if Z coincides with the set of all vertices of an n -gonal prism, then $af(Z)$ is of order n^2 ;
- (ii) if Z coincides with the set of all vertices of a convex n -gonal pyramid, then $af(Z)$ is of order n .

But it may happen that, for a finite convexly independent subset Z of $\mathbf{R}^3$ with $\text{card}(Z) = n$, the value $af(Z)$ is of order strictly between $O(n)$ and $O(n^2)$. Indeed, consider in $\mathbf{R}^3$ two distinct parallel planes H_1 and H_2 . Let M_1 be a convex m -gon in H_1 and let M_2 be a convex n -gon in H_2 . Define

$$P = \text{conv}(M_1 \cup M_2).$$

Obviously, P is a convex polyhedron in $\mathbf{R}^3$ and the total number $v(P)$ of vertices of P is equal to $m + n$. Taking into account Theorem 1, we see that

$$mn \leq af(v(P)) \leq mn + 3m/2 + 3n/2,$$

which implies that the number $af(v(P))$ is of order mn . Suppose now that n is of order $\ln(m)$. Then $af(v(P))$ is of order $m \ln(m)$. So we have the finite convexly independent set $Z = v(P)$ such that $af(Z)$ is of order strictly between $O(\text{card}(Z))$ and $O((\text{card}(Z))^2)$.

Using an analogous argument, it is not difficult to show that if α is any rational number from the closed interval $[1, 2]$, then there are infinitely many natural numbers n and convexly independent subsets Z of $\mathbf{R}^3$ such that

$$\text{card}(Z) = k = n^{\alpha-1} + n$$

and the order of $af(Z)$ is equal to k^α .

Remark 3. Let Z be a finite subset of the space $\mathbf{R}^m$ satisfying the inequality $\text{card}(Z) > 2^m$. Then one necessarily has

$$af(Z) < (1/2)\text{card}(Z)(\text{card}(Z) - 1).$$

This strong inequality follows from one nice result of L. Danzer and Grünbaum [5] (see also [1]). Note that if, for a finite set $Y \subset \mathbf{R}^m$ with $\text{card}(Y) \geq 2$, the equality

$$af(Y) = (1/2)\text{card}(Y)(\text{card}(Y) - 1)$$

holds true, then Y turns out to be the set of all vertices of a k -dimensional parallelepiped, where $k \leq m$. Some other characterizations of k -dimensional parallelepipeds in the space $\mathbf{R}^m$ can be found in [7].

Theorem 2. Let Z be a finite subset of $\mathbf{R}^2$ and let $\text{card}(Z) = n \geq 4$.

Then the inequality $af(Z) \leq n^2/4 + 2$ holds true.

Moreover, the equality $af(Z) = n^2/4 + 2$ takes place if and only if the following three conditions are simultaneously satisfied:

- (a) $n = 2k$ where $k \geq 2$ is a natural number;
- (b) $\text{conv}(Z)$ is a parallelogram in $\mathbf{R}^2$;
- (c) there are two opposite sides L_1 and L_2 of $\text{conv}(Z)$ such that k many points of Z belong to L_1 and the rest k many points of Z belong to L_2 .

Theorem 2 can be proved by using induction on n .

Remark 4. Note that if Z is a finite subset of $\mathbf{R}^3$ such that

$$\text{card}(Z) = n \in \{8, 12, 16, 20, \dots\},$$

then it may happen that $af(Z) = n^2/4 + 3n/2$. Therefore, the precise inequality of Theorem 2 does not longer hold for finite subsets of $\mathbf{R}^3$.

Remark 5. Let C be a compact convex set in $\mathbf{R}^2$ containing at least two points. As is well known (see, e.g., [3]), the following two assertions are valid:

(a) if C is not a parallelogram, then C admits a representation in the form $C = \cup\{C_1, C_2, C_3\}$ such that no affine diameter of C_i ($i = 1, 2, 3$) is an affine diameter of C ;

(b) if C is a parallelogram, then C does not admit a representation in the form $C = \cup\{C_1, C_2, C_3\}$ such that no affine diameter of C_i ($i = 1, 2, 3$) is an affine diameter of C (however, in this case C can be represented in the form $C = \cup\{C_1, C_2, C_3, C_4\}$ such that no affine diameter of C_i ($i = 1, 2, 3, 4$) is an affine diameter of C).

Note that if Z is a finite subset of $\mathbf{R}^2$ containing at least two points, then the analogs of (a) and (b) for Z can be deduced directly from Theorems 1 and 2 by using induction on $\text{card}(Z)$.

REFERENCES

1. M. Aigner, G. M. Ziegler, *Proofs from The Book*. Including illustrations by Karl H. Hofmann. Third edition. Springer-Verlag, Berlin, 2004.
2. A. D. Alexandrov, *Convex Polyhedra*. Translated from the 1950 Russian edition by N. S. Dairbekov, S. S. Kutateladze and A. B. Sossinsky. With comments and bibliography by V. A. Zalgaller and appendices by L. A. Shor and Yu. A. Volkov. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2005.
3. V. G. Boltyanskii, I. Ts. Gokhberg, *Theorems and Problems of Combinatorial Geometry*. (Russian) Izd. Nauka, Moscow, 1965 .
4. V. Boltyanskii, H. Martini, P. Soltan, *Excursions into Combinatorial Geometry*. Springer-Verlag, Berlin, 1997.
5. L. Danzer, B. Grünbaum, Über zwei Probleme bezüglich konvexer Körper von P. Erdős und V. L. Klee. (German) *Math. Z.* **79** (1962), 95–99.
6. B. Grünbaum, *Convex Polytopes*. Second edition. Prepared and with a preface by Volker Kaibel, Victor Klee and Günter M. Ziegler. Graduate Texts in Mathematics, 221. Springer-Verlag, New York, 2003.
7. A. B. Kharazishvili, Characteristic properties of a parallelepiped. (Russian) *Sakharth. SSR Mecn. Akad. Moambe* **72** (1973), no. 1, 17–19.
8. A. B. Kharazishvili, Affine diameters of convex bodies. (Russian) *Sakharth. SSR Mecn. Akad. Moambe* **90** (1978), no. 3, 541–544.
9. V. P. Soltan, Affine diameters of convex bodies—survey. *Expo. Math.* **23** (2005), no. 1, 47–63.

(Received: 03.02.2026; Published online: 10.02.2026)

A. RAZMADZE MATHEMATICAL INSTITUTE OF I. JAVAKHISHVILI TBILISI STATE UNIVERSITY, 2 MERAB ALEKSIDZE II LANE, TBILISI 0193, GEORGIA

I. VEKUA INSTITUTE OF APPLIED MATHEMATICS, 2 UNIVERSITY STR., TBILISI 0186, GEORGIA
Email address: kharaz2@yahoo.com