

HARDY–LITTLEWOOD INEQUALITIES FOR FRACTIONAL INTEGRALS IN CLASSICAL MORREY SPACES

DALI MAKHARADZE¹, ALEXANDER MESKHI^{2,3*} AND HUMBERTO RAFEIRO⁴

Abstract. We present two-weight inequalities with power-type weights for fractional integrals in classical Morrey spaces defined on the semi-axis. The obtained results involve one-sided and two-sided potentials. We treat both the Adams and Spanne type results.

1. INTRODUCTION

In our note, we present two-weight inequalities with power weights for fractional integrals in classical Morrey spaces defined on the semi-axis. We consider one-sided (Riemann–Liouville and Weyl integral transforms) and two-sided fractional integral operators. We cover both Adams and Spanne type estimates. Such type inequalities play an important role, for example, in the theory of differential and integral equations (see, e.g., [10]).

The classical Morrey spaces $L^{p,\lambda}$ were introduced by C. Morrey in 1938 (see [9]) to investigate local behavior and regularity of solutions to second order elliptic partial differential equations (see, e.g., [3, 4]). There exist two types of the boundedness of fractional integral operators in Morrey spaces: S. Spanne (unpublished) and D. R. Adams [1] type theorems.

For mapping properties of integral operators involving fractional integrals in Morrey spaces, we refer, e.g., to the monographs [7, 11] and references cited therein.

The weighted inequalities with power weights for fractional integrals in the classical Lebesgue spaces were established by Hardy and Littlewood [5] in the one-dimensional case (see also the monograph [10]). Later, similar results for the multidimensional Riesz potentials were established by Stein and Weiss [12].

2. PRELIMINARIES

For a measurable function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$, the Morrey norm is defined as follows:

$$\|f\|_{M_p^\lambda(\mathbb{R}_+)} = \|f\|_{M_p^\lambda} := \sup_{I \subset \mathbb{R}_+} \left(|I|^{-\lambda} \int_I |f(x)|^p dx \right)^{\frac{1}{p}},$$

where $0 \leq \lambda < 1$, $1 \leq p < \infty$, and the supremum is taken over all bounded intervals $I \subset \mathbb{R}_+$. If $\lambda = 0$, then $M_p^\lambda(\mathbb{R}_+)$ coincides with the Lebesgue space $L^p(\mathbb{R}_+)$, whose norm is given by

$$\|f\|_{L^p(\mathbb{R}_+)} := \left(\int_{\mathbb{R}_+} |f(x)|^p dx \right)^{1/p}.$$

Denote by I_γ , R_γ , and W_α the fractional integral operators defined on $\mathbb{R}_+$ and given by the formulas

$$I_\gamma f(x) = \int_0^\infty \frac{f(t)}{|x-t|^{1-\gamma}} dt, \quad R_\gamma f(x) = \int_0^x \frac{f(t)}{(x-t)^{1-\gamma}} dt, \quad W_\gamma f(x) = \int_x^\infty \frac{f(t)}{(t-x)^{1-\gamma}} dt,$$

where $x \in \mathbb{R}_+$ and $0 < \gamma < 1$. It is easy to check that the following pointwise estimates hold for non-negative locally integrable f :

2020 *Mathematics Subject Classification.* 26A33, 42B20.

Key words and phrases. Grand variable exponent Morrey spaces; Fractional integrals.

*Corresponding author.

(i)

$$R_\gamma f(x) + W_\gamma f(x) = I_\gamma f(x),$$

(ii)

$$I_\gamma f(x) \geq \max \{W_\gamma f(x), R_\gamma f(x)\}, \quad f \geq 0.$$

Let us recall (see Theorems A and B) the classical Hardy–Littlewood inequalities for the operators I_γ , R_γ , and W_γ , where throughout we set $p' := \frac{p}{p-1}$:

Theorem A ([5]). *Let $0 < \gamma < 1$, $1 < p \leq q < \infty$, $\alpha < \frac{1}{p'}$, $\beta < \frac{1}{q}$, $\alpha + \beta \geq 0$, and $\frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$. Then there is a positive constant C such that for all f , $\|x^\alpha f(x)\|_{L^p(\mathbb{R}_+)} < \infty$,*

$$\|x^{-\beta} I_\gamma f(x)\|_{L^q(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{L^p(\mathbb{R}_+)}.$$

Theorem B (see, e.g., [10, Theorem 5.4]). *Let $1 < p < \infty$, $0 < \gamma < \alpha + \beta + \frac{1}{p}$, $0 \leq \alpha + \beta \leq \gamma$, and $\frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$. Then the following statements are true:*

(i) *if $\alpha < \frac{1}{p'}$, then the following inequality holds for all f , $\|x^\alpha f(x)\|_{L^p(\mathbb{R}_+)} < \infty$,*

$$\|x^{-\beta} W_\gamma f(x)\|_{L^q(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{L^p(\mathbb{R}_+)};$$

(ii) *if $\beta < \frac{1}{q}$, then for all f , $\|x^\alpha f(x)\|_{L^p(\mathbb{R}_+)} < \infty$,*

$$\|x^{-\beta} R_\alpha f(x)\|_{L^q(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{L^p(\mathbb{R}_+)}$$

for the positive constant C .

The next statement is the Adams type inequality for fractional integrals which is well-known for fractional integrals, but we are interested in operator norm estimate.

In [6], the following Adams type Stein–Weiss inequality was proved on homogeneous groups, but we formulate it for the Riesz potential

$$I_\gamma^{(n)} f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\gamma}} dy, \quad 0 < \gamma < n, \quad x \in \mathbb{R}^n,$$

defined on $\mathbb{R}^n$.

Theorem C. *Let $0 < \gamma < n$, $0 \leq \alpha + \beta \leq \gamma < n$, $1 < p < \frac{n}{\gamma-\alpha-\beta}$, $\frac{1}{p} - \frac{1}{q} = \frac{\gamma-\alpha-\beta}{n-\lambda}$, $\beta < \frac{n-\lambda}{q}$, $\alpha < \frac{n}{p'}$, $0 < \lambda < n - (\gamma - \alpha - \beta)p$. Then the following weighted inequality holds for f , $\||x|^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}^n)} < \infty$,*

$$\||x|^{-\beta} I_\gamma^{(n)} f(x)\|_{M_q^\lambda(\mathbb{R}^n)} \leq C \||x|^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}^n)}.$$

Remark. Taking $\alpha = \beta = \lambda = 0$ in Theorem C, we have the classical Hardy–Littlewood–Sobolev inequality.

3. MAIN RESULTS

Now, we formulate the main statements of our paper; their proofs will appear in a forthcoming publication.

Adams-type results.

Theorem 3.1. *Let $\gamma \in (0, 1)$ and let $\alpha, \beta \in \mathbb{R}$. If $0 \leq \alpha + \beta \leq \gamma < 1$, $1 < p < \frac{1}{\gamma-\alpha-\beta}$, $\frac{1}{q} = \frac{1}{p} - \frac{\gamma-\alpha-\beta}{1-\lambda}$, $0 < \lambda < 1 - (\gamma - \alpha - \beta)p$, and $\beta < \frac{1-\lambda}{q}$, then there is a positive constant C such that the weighted inequality holds for all f , $\|x^\alpha f(x)\|_{M_p^\lambda} < \infty$:*

$$\|x^{-\beta} R_\gamma f(x)\|_{M_q^\lambda} \leq C \|x^\alpha f(x)\|_{M_p^\lambda}.$$

Theorem 3.2. Let $0 < \gamma < 1$, $0 \leq \alpha + \beta \leq \gamma < 1$, $1 < p < \frac{1}{\gamma - \alpha - \beta}$, $\frac{1}{p} - \frac{1}{q} = \frac{\gamma - \alpha - \beta}{1 - \lambda}$, $\alpha < \frac{1}{p'}$, and $0 < \lambda < 1 - (\gamma - \alpha - \beta)p$. Then the following inequality holds for all f , $\|x^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}_+)} < \infty$:

$$\|x^{-\beta} W_\gamma f(x)\|_{M_q^\lambda(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}_+)}$$

with some positive constant C .

Using Theorems 3.1 and 3.2 and the relation $I_\alpha f(x) = W_\gamma f(x) + R_\gamma f(x)$, we get the next statement.

Theorem 3.3. Let $\gamma \in (0, 1)$, $0 \leq \alpha + \beta \leq \gamma < 1$, $1 < p < \frac{1}{\gamma - \alpha - \beta}$, $\frac{1}{p} - \frac{1}{q} = \frac{\gamma - \alpha - \beta}{1 - \lambda}$, $0 < \lambda < 1 - (\gamma - \alpha - \beta)p$, $\alpha < \frac{1}{p'}$, and $\beta < \frac{1 - \lambda}{q}$. Then the following estimate holds for all f , $\|x^\alpha f(x)\|_{M_p^\lambda} < \infty$:

$$\|x^{-\beta} I_\gamma f(x)\|_{M_q^\lambda(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}_+)}$$

with some positive constant C .

Spanne type Hardy–Littlewood inequalities. Now, we discuss the Spanne type results for fractional integrals in the classical Morrey spaces.

Theorem 3.4. Let $0 < \gamma < 1$, $1 < p \leq q < \infty$, $\alpha < 1/p'$, $0 < \beta < \frac{1}{q}$, $\gamma \geq \alpha + \beta \geq 0$, $\frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$, $0 < \lambda < 1$, $0 < \mu < 1$, and $\frac{\lambda}{p} = \frac{\mu}{q}$. Then there is a positive constant C such that for all f , $\|x^\alpha f(x)\|_{M_p^\lambda} < \infty$, the inequality

$$\|x^{-\beta} I_\gamma f(x)\|_{M_q^\mu(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}_+)}$$

holds.

Theorem 3.5. Let $1 < p \leq q < \infty$, $\alpha < \frac{1}{p'}$, $\beta > 0$, $\alpha + \beta \leq \gamma < \alpha + \beta + \frac{1}{p'}$, $\frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$, $0 < \lambda < 1$, $0 < \mu < 1$, and $\frac{\lambda}{p} = \frac{\mu}{q}$. Then there is a constant C such that for all f , $\|x^\alpha f(x)\|_{M^{p,\lambda}(\mathbb{R}_+)} < \infty$, the inequality

$$\|x^{-\beta} W_\gamma f(x)\|_{M_q^\mu(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}_+)}$$

holds.

Theorem 3.6. Let $1 < p \leq q < \infty$, $0 < \beta < \frac{1}{q}$, $0 < \gamma < \alpha + \beta + \frac{1}{p'}$, $\frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$, $0 < \lambda, \mu < 1$, and $\frac{\lambda}{p} = \frac{\mu}{q}$. Then the following inequality holds for all f , $\|x^\alpha f(x)\|_{M^{p,\lambda}(\mathbb{R}_+)} < \infty$,

$$\|x^{-\beta} R_\gamma f(x)\|_{M_q^\mu(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M_p^\lambda(\mathbb{R}_+)}$$

for some positive constant C .

We can introduce a Morrey norm in a somewhat different way: we say that $f \in M^{p,s}(\mathbb{R}_+)$ if

$$\|f\|_{M^{p,s}(\mathbb{R}_+)} = \sup_{I \subset \mathbb{R}_+} |I|^{\frac{1}{s} - \frac{1}{p}} \left(\int_I |f(x)|^p dx \right)^{\frac{1}{p}}, \quad 1 < p < s < \infty.$$

It is convenient to formulate Theorems 3.4–3.6 in terms of $M^{p,s}(\mathbb{R}_+)$ norms.

Theorem 3.7. Let $0 < \gamma < 1$, $1 < p \leq q < \infty$, $p \leq s < \infty$, and $q \leq r < \infty$. Suppose that $\alpha < \frac{1}{p'}$, $0 < \beta < \frac{1}{q}$, $0 < \alpha + \beta \leq \gamma$, and $\frac{1}{s} - \frac{1}{r} = \frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$. Then there is a positive constant C such that for all f , $\|f(x)x^\alpha\|_{M^{p,s}(\mathbb{R}_+)} < \infty$,

$$\|x^{-\beta} I_\gamma f(x)\|_{M^{q,r}(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M^{p,s}(\mathbb{R}_+)}.$$

Theorem 3.8. Let $0 < \gamma < 1$, $1 < p \leq q < \infty$, $p \leq s < \infty$ and $q \leq r < \infty$. Suppose that $\alpha < \frac{1}{p'}$, $0 < \gamma < \alpha + \beta + \frac{1}{p'}$, and $\frac{1}{s} - \frac{1}{r} = \frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$. Then there is a positive constant C such that the inequality

$$\|x^{-\beta} W_\gamma f(x)\|_{M^{q,r}(\mathbb{R}_+)} \leq C \|x^\alpha f(x)\|_{M^{p,s}(\mathbb{R}_+)}$$

holds for all f , $\|x^\alpha f(x)\|_{M^{p,s}(\mathbb{R}_+)} < \infty$.

Theorem 3.9. *Let $0 < \gamma < 1$, $1 < p \leq q < \infty$, $p \leq s < \infty$, and $q \leq r < \infty$. Suppose that $0 \leq \beta < \frac{1}{q}$, $0 < \gamma < \alpha + \beta + \frac{1}{p'}$, and $\frac{1}{s} - \frac{1}{r} = \frac{1}{p} - \frac{1}{q} = \gamma - \alpha - \beta$. Then there is a positive constant C such that the inequality*

$$\|x^{-\beta} R_{\gamma} f(x)\|_{M^{q,r}(\mathbb{R}_+)} \leq C \|x^{\alpha} f(x)\|_{M^{p,s}(\mathbb{R}_+)}$$

holds for all f , $\|x^{\alpha} f(x)\|_{M^{p,s}(\mathbb{R}_+)} < \infty$.

REFERENCES

1. D. R. Adams, A note on Riesz potentials. *Duke Math. J.* **42** (1975), no. 4, 765–778.
2. A. Eridani, V. Kokilashvili, A. Meskhi, Morrey spaces and fractional integral operators. *Expo. Math.* **27** (2009), no. 3, 227–239.
3. M. Giaquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*. Annals of Mathematics Studies, 105. Princeton University Press, Princeton, NJ, 1983.
4. D. Gilbarg, N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*. Second edition. Grundlehren der mathematischen Wissenschaften, 224. Springer-Verlag, Berlin, 1983.
5. G. H. Hardy, J. E. Littlewood, Some properties of fractional integrals. I. *Math. Z.* **27** (1928), no. 1, 565–606.
6. A. Kassymov, M. A. Ragusa, M. I. Ruzhansky, D. Suragan, Stein-Weiss-Adams inequality on Morrey spaces. *J. Funct. Anal.* **285** (2023), no. 11, Paper no. 110152, 30 pp.
7. V. Kokilashvili, A. Meskhi, H. Rafeiro, S. Samko, *Integral Operators in Non-standard Function Spaces*. vol. 2. Variable exponent Hölder, Morrey-Campanato and grand spaces. Operator Theory: Advances and Applications, 249. Birkhäuser/Springer, Cham, 2016.
8. A. Meskhi, Criteria for the boundedness of potential operators in grand Lebesgue spaces. *Proc. A. Razmadze Math. Inst.* **169** (2015), 119–132.
9. C. B. Jr. Morrey, On the solutions of quasi-linear elliptic partial differential equations. *Trans. Amer. Math. Soc.* **43** (1938), no. 1, 126–166.
10. S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives*. Theory and applications. Edited and with a foreword by S. M. Nikol'skiĭ. Translated from the 1987 Russian original. Revised by the authors. Gordon and Breach Science Publishers, Yverdon, 1993.
11. Y. Sawano, G. Di Fazio, D. I. Hakim, Morrey Spaces, *Morrey Spaces—introduction and Applications to Integral Operators and PDE's*. vol. I, II. Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2020.
12. E. M. Stein, G. Weiss, Fractional integrals on n -dimensional Euclidean space. *J. Math. Mech.* **7** (1958), 503–514.

(Received: 12.02.2026; Published online: 21.04.2026)

¹DEPARTMENT OF MATHEMATICS, BATUMI SH. RUSTAVELI STATE UNIVERSITY, 32/35 RUSTAVELI/NINOSHVILI STR., BATUMI 6010, GEORGIA

²A. RAZMADZE MATHEMATICAL INSTITUTE OF I. JAVAKHISHVILI TBILISI STATE UNIVERSITY, 2 MERAB ALEKSIDZE II LANE, TBILISI 0193, GEORGIA

³G. NIKOLADZE SCIENTIFIC-RESEARCH CENTER, KUTAISI INTERNATIONAL UNIVERSITY, YOUTH AVENUE TURN 5/7, KUTAISI 4600, GEORGIA

⁴UNITED ARAB EMIRATES UNIVERSITY, DEPARTMENT OF MATHEMATICAL SCIENCES, AL AIN, UAE

Email address: dali.makharadze@bsu.edu.ge

Email address: alexander.meskhi@kiu.edu.ge

Email address: rafeiro@uaeu.ac.ae